

# ORDOVICIAN SEDIMENTARY BASINS AND PALEOBIOTAS OF THE GORNY ALTAI

*N.V. Sennikov, O.T. Obut, E.V. Lykova,  
A.V. Timokhin, T.V. Gonta, R.A. Khabibulina,  
T.A. Shcherbanenko, T.P. Kipriyanova*



TROFIMUK INSTITUTE OF PETROLEUM GEOLOGY AND GEOPHYSICS  
RUSSIAN ACADEMY OF SCIENCES  
SIBERIAN BRANCH  
NOVOSIBIRSK NATIONAL RESEARCH STATE UNIVERSITY

---

N.V. Sennikov, O.T. Obut, E.V. Lykova,  
A.V. Timokhin, T.V. Gonta, R.A. Khabibulina,  
T.A. Shcherbanenko, T.P. Kipriyanova

# ORDOVICIAN SEDIMENTARY BASINS AND PALEOBIOTAS OF THE GORNY ALTAI

Edited by  
N.V. Sennikov and A.V. Kanygin

## 13<sup>th</sup> ISOS



## NOVOSIBIRSK 2019



Novosibirsk  
Publishing House of SB RAS  
2019

УДК 551.733  
ББК 26 33  
С31

DOI 10.15372/ORDOVICIAN2019SNV

**Sennikov N.V.** Ordovician sedimentary basins and paleobiotas of the Gorny Altai / **N.V. Sennikov, O.T. Obut, E.V. Lykova, A.V. Timokhin, T.V. Gonta, R.A. Khabibulina, T.A. Shcherbanenko, T.P. Kipriyanova** ; Eds N.V. Sennikov and A.V. Kanygin ; [Translated by N.N. Mzhel'skaya, O.T. Obut, N.V. Sennikov] ; Trofimuk Institute of Petroleum Geology and Geophysics SB RAS; Novosibirsk National Research State University. – Novosibirsk : Publishing House of SB RAS, 2019. –184 p.

**Сенников Н.В.** Ордовикские седиментационные бассейны и палеобиоты Горного Алтая / **Н.В. Сенников, О.Т. Обут, Е.В. Лыкова, А.В. Тимохин, Т.В. Гонта, Р.А. Хабибулина, Т.А. Щербаненко, Т.П. Киприянова** ; Ред. Н.В. Сенников, А.В. Каныгин ; [Перевод Н.Н. Мжельской, О.Т. Обут, Н.В. Сеникова]; Институт нефтегазовой геологии и геофизики им. А.А. Трофимука СО РАН; Новосибирский национальный исследовательский университет. Новосибирск : Изд-во СО РАН, 2019. – 184 с.

### Organization



International Commission on Stratigraphy



Subcommission on Ordovician Stratigraphy



IGSP 653 “The onset of the Great Ordovician Biodiversification Event”



Trofimuk Institute of Petroleum Geology and Geophysics SB RAS



Novosibirsk National Research State University



Siberian Branch of Russian Academy of Sciences



Russian Foundation for Basic Research (RFBR)

Translated by N.N. Mzhel'skaya, O.T. Obut, N.V. Sennikov

Reviewers:  
Doctor of Sciences in Geology A.V. Dronov  
Doctor of Sciences in Geology S.V. Rozhnov

ISBN 978-5-7692-1656-5

© ИНГГ СО РАН, 2019  
© Коллектив авторов, 2019



---

---

This book is an expanded guidebook for a field excursion to the Ordovician sections of the Gorny Altai being shown to participants in the 13th International Symposium on the Ordovician System, Novosibirsk, 2019. The present guidebook concerns Ordovician stratigraphy, palaeogeography, lithology, and facies in key sections in the Gorny Altai, described bed-by-bed.

These data were previously published in part, mainly in syntheses scattered through various books and papers. This book is a synthesis of present-day knowledge with revisions of the paleontology and age implications. Updating has become possible due to discoveries of graptolite and conodont assemblages of zonal significance. The book is based on abundant biostratigraphic and geological data accumulated over the past 15 years about Ordovician deposition in the Gorny Altai. Publication of these data is a necessary step toward broad public discussion of new ideas, a prerequisite for updating the Gorny Altai Ordovician stratigraphic charts.

---

---



## CONTENTS

INTRODUCTION.....	5
1. ORDOVICIAN STRATIGRAPHY OF THE GORNY ALTAI .....	8
1.1. ORDOVICIAN REGIONAL STRATIGRAPHIC UNITS .....	—
1.1.1. REGIONAL STRATIGRAPHIC UNITS OF THE WESTERN PART OF ASFA .....	22
1.2. ORDOVICIAN GRAPTOLITE ZONES OF THE GORNY ALTAI .....	27
1.3. ORDOVICIAN CONODONT ZONES OF THE GORNY ALTAI .....	29
1.4. LOCAL ORDOVICIAN STRATIGRAPHIC UNITS OF THE GORNY ALTAI.....	30
1.4.1. OCEANIC DEPOSITION .....	—
1.4.2. SHELF DEPOSITION .....	34
1.4.2.1. THE FIRST FACIES TYPE OF SHELF DEPOSITION.....	—
1.4.2.2. THE SECOND FACIES TYPE OF SHELF DEPOSITION .....	35
1.4.2.3. THE THIRD FACIES TYPE OF SHELF DEPOSITION .....	—
2. TRACES OF GLOBAL SEDIMENTARY EVENTS IN THE ORDOVICIAN SECTIONS OF THE GORNY ALTAI .....	37
3. PALEO GEOGRAPHY OF THE WESTERN ALTAI-SAYAN FOLDED AREA (ORDOVICIAN ALTAI- SALAIR BASIN) .....	43
3.1. SEDIMENTARY TYPIES AND BIOTAS OF THE ALTAI-SALAIR ORDOVICIAN BASIN	
3.1.1. OCEANIC GENESIS.....	—
3.1.2. CONTINENTAL SLOPE GENESIS.....	—
3.1.3. SHELF GENESIS.....	—
3.2. SEDIMENTARY MODEL PROFILES FOR THE ORDOVICIAN OF THE ALTAI BASIN .....	45
3.3. PALEO GEOGRAPHIC RECONSTRUCTION OF THE ORDOVICIAN ALTAI BASIN.....	52
4. ORDOVICIAN KEY SECTIONS IN THE GORNY ALTAI.....	53
4.1. NORTH-EASTERN GORNY ALTAI (UYMEN'–LEBED' FACIES ZONE) .....	—
4.1.1. AREA OF VERKH-BIYSK VILLAGE .....	54
4.1.2. AREA OF TUROCHAK VILLAGE (LEBED' RIVER) .....	58
4.1.3. AREA OF TULOI VILLAGE.....	73
4.2. EASTERN GORNY ALTAI (TELETSKOE LAKESIDE FACIES ZONE) .....	84
4.2.1. AREA OF IOGACH VILLAGE .....	—
4.3. NORTHERN GORNY ALTAI (BIYA-KATUN' FACIES ZONE).....	87
4.3.1. AREA OF KAMLAK VILLAGE .....	—
4.4. NORTH-WESTERN GORNY ALTAI (ANUI-CHUYA FACIES ZONE) .....	90
4.4.1. AREA OF UST'-KAN VILLAGE.....	—
4.4.2. AREA OF UST'-MUTA VILLAGE .....	95
4.5. WESTERN GORNY ALTAI (CHARYSH-INYA FACIES ZONE) .....	103
4.5.1. AREA OF UST'-CHAGYRKA VILLAGE .....	—
4.5.2. AREA OF MARALIKHA VILLAGE .....	110
4.5.3. AREA OF BUGRYSHIKHA VILLAGE .....	115
4.5.4. AREA OF KRASNOSHCHIEKOV VILLAGE .....	120
4.5.5. AREA OF CHINETA VILLAGE.....	123
CONCLUSIONS .....	127
REFERENCES .....	128
PALEONTOLOGICAL PLATES .....	137

---

---

## INTRODUCTION

---

---

Ordovician strata are widespread in four major tectonic units of Siberia: in the Altai-Sayan Folded Area, upon the Siberian Craton, in the Taimyr Peninsula, and in the basement of the West Siberian Plate (Fig. 1).

The Gorny Altai lies in the western Altai-Sayan Folded Area (ASFA), a collage of terranes within the Central Asian orogen. The ASFA comprises several large geologic structures composed of Paleozoic (including Ordovician) formations of differing origins. Additional to the Gorny Altai are the Salair, Kuznetsky Alatau and Gornaya Shoriya in the west, and the West Sayan, and Tuva in the east (Fig. 2). The present tectonic framework of the ASFA (Dobretsov, 2003) has resulted from successive accretion of orogens of different ages to the Siberian craton.

In the Ordovician sediments of the Gorny Altai, all the stages (or substages or their parts) of the new Ordovician stage standard of the GSSP – Tremadocian, Floian, Dapingian, Darriwilian, Sandbian, Katian, and Hirnantian (Sennikov, Tolmacheva, 2013; Sennikov et al., 2014, 2015a, 2018a,b) are well-defined and identifiable in certain stratigraphic intervals of specific terrigenous and terrigenous carbonate sections. The application of the Ordovician stages with clear-cut lower boundaries, marked by the first appearance of the index species of the graptolite and conodont zones, and the application of the proposed “informal” substages (time slices) (Bergström et al., 2009) stimulated a major revision of the Ordovician regional subdivisions of the ASFA with the subsequent considerable specification of the chronostratigraphic extent of the previously used horizons and recognition of new ones. Also, the chronostratigraphic position of most of the Ordovician local stratigraphic units (formations) of the western ASFA has been determined more precisely. This dramatic revision has revealed many disputable problems of the Ordovician biostratigraphy of the ASFA, and the new, mainly zonal, biostratigraphic data have made it possible to propose their substantiated modern solutions. The Ordovician sections of the western ASFA are key to the resolution of the disputable questions related to the positioning of the boundaries of stratigraphic units in European Russia, not only in Asian geologic regions such as the Siberian Platform, Tuva, Kolyma, and Chukchi Peninsula, where both sedimentary and volcanosedimentary rocks occur in abundance.

The Altaian regional graptolite and conodont pelagic zonal successions can be used for precise correlations of the boundaries of local and regional units (or their parts) with the Ordovician stage boundaries of the ISC – the so-called “direct” correlations with stage boundaries based on the FAD (First Appearance Data) of the same-named index species. This shows the high efficiency of application of pelagic zonal successions in chronostratigraphy. The stratigraphic position of the ISC stage boundaries is easy to determine in the folded areas of Siberia (Gorny Altai), in which graptolites occur in terrigenous sediments and deepwater conodont assemblages are observed in carbonates.

The Ordovician sedimentary patterns of the Gorny Altai (Fig. 3) consist mainly of rhythmic alternation of terrigenous and carbonate rocks with rare volcanic intercalations. The terrigenous sections occasionally contain limestone lenses. Biohermal carbonates are frequent, mostly as algal buildups. Various facies occur dispersed through the palaeo-basin; thicknesses vary greatly.

The Gorny Altai territory is completely covered by 1:200 000 geological surveys and much of it by 1:50000 surveys undertaken during the past 60 to 70 years. Most of the surveys were carried out by people from the West Siberian Geological Surveys (currently the Zapsibgeol'syomka Prospecting Company). Large-scale (1:25000, 1:10 000, 1:5 000) geological surveys for the reference localities of the Ordovician rocks in the Gorny Altai were carried out as special stratigraphic investigations by Ermikov V.D., Gladkikh L.A., Khlebnikova T.V., Krivchikov A.V., Kuznetsov S.A., Mamlin A.N., Petrunina Z.E., Podryadchikov S.S., Puzyrev A.A., Sennikov N.V., Shokalsky S.P., Yolkin E.A., Zeifert L.L., and Zybin V.A.

Paleontological and stratigraphical data, analyzed in this monograph were collected during field studies over many years by Alekseenko A.A., Andreeva O.N., Bogashchenko E.I., Buyanova E.V., Ermikov V.D., Gabova M.F., Gladkikh L.A., Gonta T.V., Iwata K., Izokh N.G., Kalinin E.A., Khabibulina R.A., Khlebnikova T.V., Krivchikov A.V., Kul'kov N.P., Lykova E.V., Mamlina M.I., Mannik P., Mel'nikova L.M., Obut O.T., Perfil'ev E.E., Petrunina Z.E., Savitsky V.R., Sennikov N.V., Severgina L.G., Sharudo E.A., Timokhin A.V., Tokarev D.A., Tolmacheva T.Yu., Yakovleva E.V., and Yolkin E.A.







Fig. 1. Geological map of Russia and adjoining water areas.





**Fig. 2.** Russia with location of the Altai-Sayan Folded Area (ASFA) including Rudnyy Altai, Gornyy Altai, Salair, Kuznetsky Alatau, Kuznetsk basin, Minusa, West Sayan and Tuva.



**Fig. 3.** Sketch map of locations of the Altai Ordovician deposits with main structural-tectonic elements.

A – areas with the wide-spread occurrence of the Ordovician strata; B – deep faults, constraint regional blocks; C-D – boundaries: C – structural-facies zones, D – subzones (areas). 1-12 – structural-facies zones: 1 – Milovanovka; 2 – Loktevka-Batun; 3 – Suetka-Kuibyshevo; 4 – Charysh-Inya; 5 – Kharlovo; 6 – Vydrikha; 7 – Talitsa; 8 – Anui-Chuya; 9 – Biya-Katun'; 10 – Uyem'-Lebed'; 11 – Teletskoe Lakeside; 12 – Ulagan.

Faunal remains included in the present monograph were identified as follows: trilobites – Petrunina Z.E., Andreeva O.N., Yolkin E.A., Timokhin A.V.; graptolites – Obut A.M., Sennikov N.V., Lykova E.V.; conodonts – Moskalenko T.A., Izokh N.G., Obut O.T., Tolmacheva T.Yu.; ostracods – Mel'nikova L.M., Gonta T.V.; brachiopods – Severgina L.G., Andreeva O.N., Kul'kov N.P., Savitsky V.R., Modzalevskaya T.L., Shcherbanenko T.A.; stromatoporoids – Khromykh V.G.; tabulate corals – Dzyubo P.S., Galenko L.V., Khabibulina R.A.; rugose corals – Zheltonogova V.A.; bryozoans – Volkova K.N., Koromyslova A.V.; crinoids – Stukalina G.A., Dubatolova Yu.A.; gastropods – Byalyi V.I.; radiolarians – Obut O.T., Iwata K.; chitinozoans – Obut A.M., Zaslavskaya N.M., Obut O.T.

Photographs illustrated sections and outcrops have been taken by Sennikov N.V., Obut O.T., Lykova E.V., and Mannik P.

## 1. ORDOVICIAN STRATIGRAPHY OF THE GORNY ALTAI

The Ordovician stratigraphic charts of the Gorny Altai have been compiled through the past seven decades by joint efforts of prospecting, academic, and educational institutions, namely, All-Russian Geological Institute (VSEGEI, St. Petersburg), Siberian Research Institute of Geology, Geophysics & Mineral Resources (SNIIGGiMS, Novosibirsk), Institute of Geology & Geophysics (currently Trofimuk Institute of Petroleum Geology and Geophysics, Novosibirsk), Zapsibgeologiya (Novokuznetsk), Zapsibgeol'syomka (Elan' Village), Tomsk and Novosibirsk State Universities, Tomsk Technological Institute (currently Tomsk Technological University), Kuzbass Pedagogical Academy (Novokuznetsk), and others.

Among researchers who played important role in the investigation of the Ordovician stratigraphy of the Gorny Altai should be noted the following persons (in the alphabetic order): Andreeva O.N., Avrov D.P., Bartseva N.M., Bublichenko N.L., Cherepnina S.V., Dubatolova Yu.A., Dzyubo P.S., Ermikov V.D., Fedyanov V.V., Galenko L.V., Gintsinger A.B., Gladkikh L.A., Gutak Ya.M., Isaev G.D., Izokh N.G., Khalfina V.K., Khlebnikova T.V., Khromykh V.G., Kononov A.N., Krivchikov A.V., Krivchikov V.A., Kuznetsov V.A., Kul'kov N.P., Levitsky E.S., Lyakhnitsky V.N., Lykova E.V., Mamlin A.N., Modzalevskaya E.A., Modzalevskaya T.L., Moskalenko T.A., Nekhoroshev V.P., Nikonov A.A., Obut A.M., Obut O.T., Perfil'ev Yu.S., Petrunina Z.E., Podryadchikov S.S., Potapova M.S., Puzyrev A.A., Romanenko M.F., Savitsky V.P., Sennikov N.V., Sennikov V.M., Severgina L.G., Stukalina G.A., Tikhonov V.I., Usov M.A., Vyltsan I.A., Vinkman M.K., Volkov V.V., Volkova K.N., Yaroshinskaya A.M., Yolkin E.A., Zaslavskaya N.M., Zeifert L.L., Zheltonogova V.A.

### 1.1. ORDOVICIAN REGIONAL STRATIGRAPHIC UNITS

According to the usual practice in Russia, regional correlation follows chronostratigraphic subdivision into regional stages (called *horizons* in the Russian stratigraphic code). Regional stages comprise regionally spread coeval formations (or their parts) and correspond to stages of regional geological history, and especially to stages in the evolution of marine faunal groups. Regional stages are discussed and approved collectively for further use at special Russian Stratigraphic Workshops held every ten to fifteen years, or sometimes every five or seven years. The succession of regional stages, together with the respective subdivision of the International Stratigraphic Chart (Scale), make basis for correlation of local units (formations and groups) as part of regional charts and for inter-regional correlations.

There are three main steps in synthesis of stratigraphic and biostratigraphic data: (1) subdivision of local sections, (2) correlation of the sections (intra- and partly inter-regional correlations of composite sections from nearby regions), (3) age assignment (correlation among sections from geographically dispersed regions and global correlation, including correlation with the units in the International Stratigraphic Chart). The three objectives are performed with reference to different Ordovician faunal groups.

The greatest part of local Ordovician sections in the Gorny Altai are correlated mainly according to trilobites and brachiopods. These two groups of benthic organisms occur in diverse complexes of terrigenous, terrigenous-carbonate and purely carbonate compositions and, for this reason, are used in subdivision of local sections and in intra-regional correlation. Trilobites and brachiopods are also of broad use in correlation among sections from the nearby areas of Gorny Altai, Salair, Kuznetsky Alatau, Siberian Platform, Taimyr, Kazakhstan, and Urals.

Other communities besides trilobites and brachiopods used in subdivision of local sections are abundant tabulate and rugose corals, ostracods, fishes, bryozoans, crinoids, nautiloids, gastropods, stromatoporoids, radiolarians, and chitinozoans. In some cases, they may be useful for intra- or, more rarely, inter-regional correlations.

Besides the local subdivision, the succession of trilobite and brachiopod assemblages is the key to Gorny Altai regional Ordovician stages, and these, in turn, most often serve as ties in correlation among sections from proximal regions.

Succession of trilobite and graptolite assemblages is regarded as the base for the Altai Ordovician regional stages (horizons) as well as a reliable key for the intra-regional correlation. Inter-regional correlation is usually fulfilled on the bases of regional stages (horizons).

Global correlation and dating of the deposits with respect to the International Stratigraphic Chart are made proceeding from pelagic faunas of graptolites and conodonts. Although these faunal groups are less abundant in the Ordovician sections of the Gorny Altai than others.

The current Paleozoic stratigraphic charts for Central Siberia were approved at the USSR Workshops on Ordovician-Silurian Stratigraphy of 1956, 1964, 1979 and 2012 in Novosibirsk which were four milestones in the history of stratigraphic studies in the Gorny Altai and in the Altai-Sayan Folded Area as a whole (Decisions..., 1959; Documents..., 1964; Correlative..., 1964; Stratigraphy..., 1967; Decisions..., 1983; Sennikov et al., 2018a,b). The workshops concerned with unification of regional stratigraphic charts as an outcome of years-long work; the charts were then considered by the USSR Interdepartmental Stratigraphic Committee and ratified to become official guidelines for further use in geological surveys.

The Ordovician regional stages of the Gorny Altai are correlated to the units of the International Stratigraphic Chart (Scale) on the basis of the respective graptolite, conodont and chitinozoans zones (Fig. 4).

International Stratigraphic Chart (Resolutions of ISC, 2012, vol.41)			Age, Ma (The Geologic Time Scale, 2012)	Standard zonal scales (recommended by O/S commission of the ISC, 2012)				
System	Series	Stage		Graptolites (zonation: synthesis based on data from Russian regions) (O/S commission of ISC, 2012)		Conodonts (The Geologic Time Scale, 2012)		Chitinozoans (The Geologic Time Scale, 2012)
Ordovician	Upper	Hirnantian	445.2	Normalograptus persculptus Normalograptus extraordinarius / Normalograptus ojsuensis / Normalograptus mirnyensis		Amorphognathus ordovicicus		Tanuchitina oulebsiri Tanuchitina elongata
		Katian	Appendispinograptus supernus	Paraorthograptus pacificus	Ancyrochitina merga			
				Appendispinograptus supernus	Armoricochitina nigerica			
			Orthograptus quadrimucronatus	Amorphognathus superbus	Acanthochitina barbata			
					Dicranograptus clingani	Tanuchitina fistulosa		
					Diplocanthograptus caudatus	Belonechitina robusta Euconochitina tanvillensis		
		Sandbian	458.4	Climacograptus bicornis	Amorphognathus tvaerensis	Baltoniodus alobatus	?	
						Diplograptus multidentis / Diplograptus foliaceus	Baltoniodus gerdæ	Lagenochitina dalbyensis
				Nemagraptus gracilis / Oepikograptus beckeri		Baltoniodus variabilis	Lagenochitina deunffi	
						Amorph. inaequalis	Lagenochitina ponceti	
	Middle	Darriwilian	467.3	Hustedograptus teretiusculus	Pygodus anserinus	Sagittodontina kieltensis	Linochitina pissotensis	
						Pygodus serra	Laufeldochitina clavata	
				Didymograptus murchisoni / Didymograptus geminus	Eoplacognathus suecicus	Armoricochitina amorçana - Cyathochitina jenkinsi		
				No zonation		Siphonochitina formosa		
				Undulograptus (= ? Eoglyp.) dentatus	Eoplacognathus variabilis	Cyathochitina calyx - protocalix		
				Undulograptus austrodentatus		Desmochitina bulla		
				Dapingian	470.0	Expansograptus hirundo	Baltoniodus norrlandicus	Belonechitina henryi
		Isograptus gibberulus	Paroistodus originalis			Desmochitina ornensis		
			Baltoniodus navis					
		Lower	Floian	477.7	Pseudophyllograptus angustifolius elongatus / Pseudophyllograptus angustifolius tenuis	Baltoniodus triangularis	Oepikodus evae	Eremochitina brevis
	Phyllograptus densus							
	Tetragraptus phyllograptoides / Tetragraptus approximatus				Prioniodus elegans	Eremochitina baculata		
	?					Paroist. proteus	Oelandodus elongatus - Acodus deltatus	Conochitina symmetrica
	Tremadocian		485.4	Araneograptus murayi	Paracordylodus gracilis	Lagenochitina brevicollis		
					Tripodus - Drepanodus aff. amoenus	Amphorachitina confundus		
				Bryograptus ramosus / Rhabdinopora uralense / Aletograptus hyperboreus	Paltodus deltifer	Lagenochitina destombesi		
				Adelograptus tenellus / Anisograptus richardsoni	Cordylodus angulatus			
Rhabdinopora flabelliformis				Iapetognathus fluctivagus				

Underlying units

1

**Fig. 4.** Alignment of the local stratigraphic charts for the Ordovician of the Gorny Altai with the International Stratigraphic Chart based on regional stages (horizons), graptolite and conodont zones.



Stage	Regional stratigraphic units			
	Paleontological characteristic of regional units			
	Zones			Specific faunal (floral) complexes, Beds with fauna (flora)
	Graptolites	Conodonts	Chitinozoans	Graptolites
Katian	Horizon			
	Subhorizon			
	List-kyan			
Sandbian	Tekhten'			
Dartwiliian	Khankhara			
Dapingian	Bugryshikhina			
Floian	Kostinsky			
Tremadocian	Kulby-shevo			
Tremadocian	Tuloi (= Lebed')			
Tremadocian	Takoshkin (= "Upper Tayanza")			

Underlying units

2

Fig. 4. Continued.

Stage	Regional stratigraphic units				
	Horizon	Subhorizon	Paleontological characteristic of regional units		
			Specific faunal (floral) complexes, Beds with fauna (flora)		
			Conodonts	Chitinozoans	Brachiopods
Hir-nant	List- vyan	?		<i>Dalmanella testudinaria</i> (Dalm.), <i>D. dietkensis</i> Severy., <i>Zygospiraella indistincta</i> Kuik. et Severy., <i>Streptis altosinuata</i> (Holt.), <i>Hirnantia</i> aff. <i>noixella</i> Amsden, <i>Breviarmuella gromotuchaensis</i> Severy., <i>Trucizetina subrotundata</i> Havi., <i>Dedzetina</i> aff. <i>microstoma</i> Havi., <i>Siegethynchus concinnus</i> (Savage), <i>Eospirigerina gaspeensis</i> (Coop.), <i>Plectatrypa</i> aff. <i>henningsmoeni</i> Bouc. et John., <i>Giraldibella bella</i> Bergst.	
Katian	Tekhten'	<i>Amorphognathus ordovicicus</i> Br. et Mehl, <i>Scabbardella altipes</i> (Henning.) Rhodes, <i>Panderodus intermedius</i> Br., Mehl et Br., <i>P. cf. uncostatus</i> (Br. et Mehl), <i>Protopanderodus insculptus</i> (Br. et M.)	<i>Conochitina micracantha</i> Eis., <i>Tanuchitina ontariensis</i> Jans.	<i>Giraldibella bella</i> (Bergst.), <i>Thebesia thebesensis</i> Amsden, <i>Leangella septata</i> (Cooper), <i>Spirigerina mediocris</i> (Severy.)	
		<i>Phragmodus undatus</i> Br. et Mehl, <i>Panderodus</i> cf. <i>P. gracilis</i> (Br. et Mehl), <i>Belodina compressa</i> (Br. et Mehl), <i>Drepanoistodus suberectus</i> (Br. et Mehl), <i>Eraticodon</i> sp.		<i>Eospirigerina orloviensis</i> (Severy.), <i>Oxoplectra platystrophoides</i> Schuchert et Cooper, <i>Catazyga inensis</i> (Severy.), <i>Catazyga anuensis</i> (Severy.), <i>Eridorthis subinexpecta digna</i> Severy., <i>Schizophorella fallax</i> Salter	
Sandbian	Khankhara		<i>Desmochitina erinacea</i> Eis., <i>D. lecaniella</i> Eis., <i>Lagenochitina dalbyensis</i> Laufeld	<i>Eospirigerina sublevis</i> (Rozm.), <i>Austinella lebediensis</i> Severy., <i>Salopina uxunaica</i> (Severy.), <i>Glyptorthis praepulchra</i> Severy., <i>G. balclatchiensis</i> (Dav.), <i>Hesperorthis incenaria lebediensis</i> Severy., <i>Dulankarella magna</i> Ruk., <i>Catazyga salainica</i> (Severy.)	
		?		<i>Boreadorthis togaensis</i> Severy., <i>Chaulistomella amzassensis</i> (Severy.), <i>Strophomena lebediensis</i> Severy. in Rozm., <i>Rostricellula ainsliei amzassica</i> Severy., <i>Togaella grandis</i> Severy.	
		<i>Eobelodina</i> cf. <i>fornicala</i> (Stauf.)		<i>Onniella chancharica</i> Severy., <i>Plectocamera usucchiensis</i> Severy., <i>Fascifera buranensis</i> Severy., <i>Bimuria bugryschichiensis</i> Severy., <i>Chaulistomella inaquistriata</i> (Cooper), <i>Eoanastrophia lebediensis</i> (Severy.)	
Darriwilian	Bugryshikha	?		<i>Apatomorpha altaica</i> Severy., <i>Leptellina tennesseensis</i> Ulr. et Cooper, <i>Hesperorthis markovae</i> Rozman, <i>Howellites</i> cf. <i>flava</i> (Havl.)	
			<i>Cyathochitina tuloensis</i> Obut et Zasl., <i>Cyath. calix</i> (Eis.), <i>Conochitina oelandica</i> Eis., <i>C. bacillum</i> Obut et Zasl.	<i>Glyptorthis primus</i> Severy., <i>Parastrophina bilobata</i> Cooper, <i>Beloviella bugryshichiensis</i> Severy., <i>Christiania</i> aff. <i>subquadrata</i> (Hall), <i>Glyptomena karasuensis</i> Severy., <i>Hesperorthis markovae oplecis</i> Severy.	
				<i>Archaeorthis altaica</i> Severy., <i>Idiostrophia tuloviensis</i> Severy.	
Dapingian	Kuibyshevo	<i>Eoplacognathus pseudoplanus</i> (Viira), <i>Periodon aculeatus</i> Hadding, <i>Paroistodus originalis</i> (Serg.), <i>Protopanderodus rectus</i> (Lind.), <i>Scolopodus giganteus</i> Sw. et Berg, <i>Juanognathus janssoni</i> Serp.		<i>Trondorthis sibirica</i> Severy.	
		<i>Parapanderodus gracilis</i> (Barnson et Mehl), <i>Yangtzeplacognathus</i> ? sp.		?	
		?			
Floian	Tuloi (= Lebed')	<i>Paroistodus</i> cf. <i>parallelus</i> (Pand.), <i>Periodon primus</i> (?) Lofgren, <i>Paroistodus</i> cf. <i>lanceolatus</i> (Pander)		<i>Ujukites tarlykensis</i> Andr., <i>Eodalmarella</i> (?) sp.	
		<i>Oepikodus evae</i> Lindstrom, <i>Periodon</i> cf. <i>flabellum</i> (Lind.), <i>Prioniodus</i> cf. <i>P. elegans</i> Pander, <i>Drepanoistodus</i> sp.	<i>Conochitina raymondi</i> Achab, <i>Con. turgida</i> (Jenk.), <i>Con. ordinaria</i> Achab	<i>Archaeorthis sibirica</i> Severy., <i>Tritoechia orliniensis</i> Severy., <i>Orthis kozuchiensis</i> Severy., <i>Hesperonomia illovata</i> (Severy.), <i>H. korzhnevi</i> Severy., <i>H. paratylyensis</i> Severy., <i>Nanorthis multicostata</i> Ulr. et Coop.	
		<i>Oepikodus evae</i> Lindstrom, <i>Paracordylodus gracilis</i> Lind., <i>Paroistodus</i> cf. <i>proteus</i> (Lind.), <i>Paroistodus</i> cf. <i>originalis</i> (Serg.), <i>Cornuodus longibasis</i> (Lind.), <i>Oneotodus</i> sp.		<i>Nanorthis gloriosus</i> Severy., <i>Diparelasma minuta</i> Severy., <i>Rhysselasma pusilla</i> Severy., <i>Akelina akelina</i> Severy.	
Tremadocian	Takoshkin (= "Upper Tayanza")	<i>Iapetonodus</i> sp., <i>Iapetognathus</i> sp., <i>Cordylodus lindstromi</i> Druce et Jones, <i>Eoconodontus notchpeakensis</i> (Miller)	?	<i>Apheorthis lineocosta</i> Walc., <i>Nanorthis schoriensis</i> Severy., <i>Nothorthis algainensis</i> Severy., <i>Punctolira kondomiensis</i> Severy.	

Underlying units

Fig. 4. Continued.

Regional stratigraphic units									
Stage	Horizon	Subhorizon	Paleontological characteristic of regional units						
			Specific faunal (floral) complexes, Beds with fauna (flora)						
			Trilobites	Ostracods	Tabulate corals	Rugose corals	Bryozoans	Stromato- poroids	
Katian	List- yan		<i>Dalmanitina mucronata</i> (Brong.)	?	?	?	?		
			<i>Gromotuchia dilavata</i> Petrun., <i>Gr. stricta</i> Petrun., <i>Homotelus</i> sp. <i>Stenoblepharum warburgae</i> (Prib.)	?	<i>Catenipora workmanae</i> Flom., <i>C. bugryschiensis</i> Dz., <i>Rhabdotetradium</i> sp., <i>Plasmoporella convextotabulata</i> Kiaer, <i>P. bugryschiensis</i> Dz., <i>Mesofavosites subfallax</i> Dz., <i>M. dualis</i> Sok., <i>Agetolites insuetus</i> Kim., <i>Heliolites</i> sp., <i>Propora parvotabulata</i> (Kiaer), <i>Karagemia altaica</i> Dz., <i>Nyctopora altaica</i> Dz.	<i>Grewinkia lebediensis</i> Tcherepn., <i>Gr. semilunatum</i> (Scheffen), <i>Ditoechelasm altaica</i> (Tcherep.), <i>Axiphoria dietkensis</i> Tschern., <i>Paliphyllum primarium</i> Soshk., <i>Brachielasma altaica</i> Tcherepn.	?		
			?	<i>Ballardina altaica</i> Melnik., <i>Eurychilina sennikovi</i> Melnik., <i>Retiprimites formosus</i> Melnik., <i>Laccochilina</i> ( <i>Laccochilina</i> ) <i>lebediensis</i> Melnik., <i>Bolbina dubia</i> Melnik., <i>Bol. karasuensis</i> Melnik., <i>Soanella petruninae</i> Melnik., <i>Egorovella demissa</i> Melnik., <i>Bullaferum paritius</i> Melnik., <i>Pseudozygobolbina invisitata</i> Melnik., <i>Allertemmya</i> (?) <i>incerta</i> Melnik.	<i>Plasmoporella vesiculosa</i> Kiaer, <i>Nyctopora minimalis</i> (Rad.), <i>Cyrtophyllum samyshiensis</i> Tcherepn., <i>Trochiscolithus micrastes</i> (Linds.)		<i>Trematopora propria</i> Jarosh., <i>Rhinidictya lebediensis</i> Jarosh., <i>R. altaica</i> Jarosh.	<i>Clathrodiction</i> (?) <i>kirgisicum amzassensis</i> V. Khalf., <i>Lophiostroma elandense</i> V. Khalf.	
			<i>Cyrtophyllum kaniensis</i> Dz., <i>Vacuopora prisca</i> (Sok.), <i>Calapoecia anticostensis</i> Bill.						
	Khankhara		<i>Holotrachelus punctiliosus</i> (Brong.), <i>Iliaenus oviformis</i> Warb., <i>Il. cf. septentrionalis</i> Tschug., <i>Amphitichas sniatkovi</i> Web., <i>Brontecephalina huda</i> (Ang.), <i>Isocolus sjogreni</i> Ang., <i>Chasmops salinus</i> Petrun., <i>Eucrinuroides bobroviensis</i> Petrun., <i>Ceraurinus icarus</i> (Bill.), <i>Calyptaulax bellatulus</i> Petrun., <i>Paracybeloides loveni</i> (Linds.) <i>Chasmopsella unica</i> Petrun., <i>Bronteopsis gregaria</i> Raum., <i>Jaboginellus gornoaltaicus</i> Petrun., <i>Otarionella kokoriana</i> Korol., <i>Eorobergia lebediensis</i> Petrun.			?			
	Sandbian		<i>Cybelurus planifrons</i> (Web.), <i>Raymondella bugryschiensis</i> Petrun.						
	Darrwilian	Bugryshtikha		<i>Lonchodomas cf. lecturmasi</i> Web., <i>L. cf. laevisculus</i> (Bill.), <i>L. (Foliopyge) levis</i> Petrun., <i>Remopleurites longicaudatus</i> Port., <i>Eorobergia cf. uzbekiata</i> Petrun., <i>Cyberulus cf. planus</i> Lev., <i>Encrinuroides</i> sp., <i>Thaleops</i> sp., <i>Homotelus angustus</i> Petrun., <i>Amopyxella</i> ( <i>Belaxella</i> ) <i>intermicostata</i> Petrun., <i>Nidius tenghensis</i> Web., <i>Telephina mobergi</i> (Hadd.)	?	?			
Kostinsky			<i>Ceraurina cf. frequens</i> Tschug., <i>Pliomera fischeri</i> asiatica Tschug., <i>Bathyporellus nonnullus</i> Tschug., <i>Carriackia</i> sp., <i>Kolymella</i> aff. <i>plana</i> (Tschug.), <i>Pliomerellus amplexissimus</i> Petrun., <i>Pl. cf. jacuticus</i> Tschug., <i>Carolinites</i> sp., <i>Glaphurus altaicus</i> Weber, <i>Raymondaspis</i> sp.	<i>Soanella tuloica</i> Melnik., <i>?Laccoprimitia</i> sp., <i>?Quadrijugatoriidae</i> , <i>Egorovella</i> sp.					
Kuibyshevo		?							
Dapingian			<i>Eorobergia integra</i> Petrun., <i>Vogdesia?</i> <i>tuloica</i> Petrun., <i>Remopleuridella altaiensis</i> Petrun., <i>Levirobergia ojtrotica</i> Petrun.						
			<i>Pliomerops</i> sp., <i>Eorobergia</i> sp.						
Floian		Tuloi (= Lebed')	<i>Taidonurus asiaticus</i> Petrun., <i>Tersella strobilata</i> Petrun., <i>T. altaica</i> Petrun., <i>Pytine sibirica</i> Petrun., <i>Lapidaria</i> ? <i>ishpensis</i> Petrun., <i>Hypermecaspis lebediensis</i> Petrun., <i>Shumardia tagasensis</i> Petrun., <i>Seleneceme improvisa</i> Petrun.						
Tremadocian		Takoshkin (= "Upper Tayanza")	<i>Apatokephalus ex gr. serratus</i> (Boeck), <i>Amazasiella mirabilis</i> Polet., <i>Platypeltoides cf. wimani</i> (Troeds.), <i>Hysterolenus</i> sp., <i>Dikelokaphalina bidens</i> Petrun., <i>Symphysurus</i> sp., <i>Ceratopyge patula</i> Petrun., <i>Kaltykelins gracilis</i> Petrun., <i>Acrocephalina</i> sp., <i>Amzasskiella obliqua</i> Petrun., <i>Niobe zhulanica</i> Petrun.	?					

Underlying units

Fig. 4. Continued.



Correlation of local stratigraphic units									
Stage	Horizon	G o r n y   A l t a i							
		Milovanovo Zone				Loktevka - Batun Zone		Suetka - Kuibyshevo Zone	
		Shallow water inner shelf				Deep water outer shelf		Deep water outer shelf	
		Milovanovka, Gornovka rivers				Area near former Kostinsky mine, vil. Rudovozy		Watershed of Suetka and Charysh rivers, right tributaries of Charysh River	
Hir- nant -yan	List- -yan	1				2		3	
						(lower part of formation, analogue of <i>persculptus</i> Zone) Black platy mudstone. ? ----- > 10 m		(lower part of formation, analogue of <i>persculptus</i> Zone) Black platy mudstone. ? ----- > 10 m	
Katian	Tekhten'	GORNOVKA SEQUENCE				SILICEOUS-TERRIGENOUS BODY		SILICEOUS-TERRIGENOUS BODY	
		Variegated conglomerate, sandstone, siltstone, shales. <i>Sowerbyella</i> sp., <i>Bimuria</i> (?) sp., <i>Strophomena</i> sp., <i>Plectrothis</i> sp., <i>Boreadorthis</i> (?) sp., <i>Flexicalymene</i> sp., <i>Encrinurus</i> sp., <i>Calyptaulax</i> sp., <i>Cheirurina</i> sp., <i>Nileus</i> sp. ~ 800 m				Alternation of sandstone, siltstone and siliceous mudstone. <i>Appendospinograptus supernus</i> El. et W., <i>App. longispinus</i> J.T.Hall, <i>Glyptograptus amplexicaulis</i> (Hall), <i>Dictyonema</i> sp.		Alternation of sandstone, siltstone and chert with rare lenses of limy mudstone. <i>Climacograptus latus</i> El. et W., <i>App. supernus</i> El. et W., <i>Orthograptus amplexicaulis</i> (Hall), <i>Borisella subulata</i> (Web. et Bl.), <i>Secuicollata ornata</i> G., Um et I., <i>Kalimnaspheera maculosa</i> Web. et Bl., <i>Protoceraoikiscum chinocrystallum</i> G., Um. et I., <i>Inanigutta complanata</i> Naz., <i>Inanigutta</i> sp., <i>Palaeotrifidus</i> sp., <i>Periodon grandis</i> Eth., <i>Panderodus</i> sp., <i>Histoidella</i> sp., <i>Protopanderodus insculptus</i> Br. et M., <i>Decoriconus</i> sp., <i>Paroistodus</i> ? mutatus Br. et M., <i>Belodina compressa</i> Br. et M. > 100 m	
Sandbian	Khankhara							KHANKHARA FM.	
								Siltstone, mudstone, rare layers of limestone. <i>Dicellograptus intortus</i> Lapw., <i>Cryptograptus tricornis</i> (Carr.), <i>Orthograptus calcaratus</i> (Lapw.), <i>Orth. barcovaensis</i> (Obut et Sob.), <i>Dipl. caudatus</i> Lapw., <i>Cl. antiquus lineatus</i> El. et W., <i>Cl. minutus</i> (Carr.), <i>Cl. bicornis</i> (Hall), <i>Glyptograptus euglyphus</i> (Lapw.), <i>Gl. eosicatus</i> Tzaj, <i>Gl. siccatus</i> El. et W., <i>Reteograptus geinitzianus tenuis</i> Sen. 300-850 m	
Bugryshikha								BUGRYSHIKHA FM.	
								Siltstone, sandstone. <i>Dicellograptus intortus</i> Lapw., <i>Cryptograptus tricornis</i> (Carr.), <i>Orthograptus calcaratus</i> (Lapw.), <i>Orth. barcovaensis</i> (Obut et Sob.), <i>Cl. antiquus lineatus</i> El. et W., <i>Cl. minutus</i> (Carr.)  <i>Thallograptus</i> sp., <i>Acrograptus serratus</i> (Hall), <i>Acr. sagitticaulis</i> (Gurley), <i>Eoglyptograptus dentatus</i> (Brong.), <i>Cryptograptus tricornis</i> (Carr.), <i>Cryp. hopkinsoni</i> (Nichol.), <i>Amplexograptus arctus</i> El. et W., <i>Expansograptus</i> sp., <i>Pseudoclimacograptus scharenbergi</i> (Lapw.), <i>Climacograptus brevis</i> El. et W., <i>Glossograptus hincksi</i> (hopk.), <i>Glos. fimbriatus</i> (Hopk.), <i>Glos. euglyphus</i> (Lapw.), <i>Hustegograptus teretiusculus</i> (His.)  <i>Pendeograptus</i> aff. <i>pendens</i> (Ell.), <i>Amplexograptus coelatus</i> (Lapw.), <i>Pseudotrigraptus ensiformis</i> (Hall), <i>Pseudotr. aff. martelli</i> (Ross et Berry), <i>Cryptograptus tricornis</i> (Carr.), <i>Cryp. hopkinsoni</i> (Nichol.), <i>Acrograptus</i> sp., <i>Expansograptus</i> sp., <i>Corymbograptus</i> sp.	
Darriwilian	Kostinsky							600-850 m	
Kuibyshevo								VOSKRESENKA FM.	
								Dark-gray mudstone, siltstone, sandstone, at the base - conglomerate, at the top - layers and lenses of sandy limestone. <i>Undulograptus austrodentatus</i> (Har. et Keble), <i>Undulograptus dentatus</i> (Brongniart), <i>Expansograptus hirundo</i> (Salt.), <i>Ex. extensus</i> (Hall), <i>Acrograptus cognatus</i> (Har. et Th.), <i>Loganograptus logani</i> (Hall)  <i>Eotetragraptus harti</i> (Hall), <i>Isograptus gibberulus</i> (Nichol.)  <i>Isograptus forcipiformis forcipiformis</i> Ruedemann  <i>Expansograptus broggeri</i> (Monsen)  <i>Tristichograptus ensiformis</i> (Hall), <i>Phyllograptus ilicifolius glaber</i> Monsen, <i>Ph. densus densus</i> Tornquist, <i>Ph. densus opulens</i> Monsen	
Dapingian	Tuloi (= Lebed')							~300 m	
Floian									
Tremadocian	Takoshkin ("Upper Tayanza")							SUETKA FM.	
								Gray-green and lilac polymictic sandstone, siltstone, clayey shales, conglomerate.	
Underlying units		Є		Є		Є		~2500 m	
						</			

Stage		Correlation of local stratigraphic units	
		G o r n y A l t a i	
Horizon	Subhorizon	Charysh-Inya Zone	
		West region	Central region
Hir-nant	List-kyan	Central part of outer shelf with reef massifs	Central part of outer shelf with reef massifs
		The watershed of Belaya and Inya rivers, vil. Bugryshikh, vill. Chineta	Chagyryk River
Katian	Tektien'	4 a	4 b
		<p>VTORYE UTYSY FM.</p> <p>Black platy mudstone. <i>Normalograptus persculptus</i> (Salter), <i>Climacograptus longifilis</i> Manck., <i>Dalmanella testudinaria</i> (Dalm.), <i>Streptis altosinuata</i> (Holt.), <i>Brevilamnuella gromatuchaensis</i> Severg., <i>Zygospiraella indistincta</i> Kulk. et Severg., <i>Dalmanitina mucronata</i> (Bronn.)</p> <p>~10 m</p> <p>TEKHTEN' FM.</p> <p>Siltstone, sandstone, lenses and bodies of bioherm and reef limestone.</p> <p><i>Normalograptus ojsuensis</i> (Koren et Mikh.), <i>Appendispinograptus supernus</i> (El. et W.), <i>Pseudoclimacograptus</i> sp., <i>Orthograptus</i> sp., <i>Glyptograptus forrainensis</i> Rued.</p> <p><i>Dalmanella testudinaria</i> (Dalm.), <i>Streptis altosinuata</i> (Holt.), <i>Zygospiraella indistincta</i> Kulk. et Severg., <i>Brevilamnuella gromatuchaensis</i> Severg., <i>Dalmanitina mucronata</i> (Bronn.), <i>Thebesia thebesensis</i> Amsd., <i>Springerina medocis</i> (Severg.), <i>Leangella septata</i> (Coop.), <i>Giraldibella bella</i> (Bergst.), <i>Stenoblepharum warburgae</i> (Prib.), <i>Mesofavosites subfallax</i> Dz., <i>Propora parvotabulata</i> (Kiaer), <i>Catenipora bugryschichensis</i> Dz., <i>Eospiriferina orloviensis</i> (Severg.), <i>Catazyga inensis</i> Severg., <i>Oxoplectra platystrophoides</i> Schuchert et Cooper, <i>Eridorthis subinsecta digna</i> Severg., <i>Plasmoporella vesiculosa</i> Kiaer, <i>Axiphoria dietkensis</i> Tschern.</p> <p>up to 300 m</p>	<p>VTORYE UTYSY FM.</p> <p>Black platy mudstone</p> <p><i>Normalograptus persculptus</i> (Salter), <i>Nor. bohemicus</i> Marek, <i>Nor. medius</i> (Tornq.), <i>Nor. angustus</i> (Pern.), <i>Climacograptus longifilis</i> Manck., <i>Nor. scalaris</i> (His.) ~10 m</p> <p>TEKHTEN' FM.</p> <p>Siltstone, sandstone, lenses and bodies of bioherm and reef limestone.</p> <p><i>Paraorthograptus pacificus</i> (Rued.), <i>Dicellograptus ornatus minor</i> Togh., <i>D. ornatus ornatus</i> El. et W., <i>App. supernus</i> E. et W., <i>C. hastatus</i> T.S. Hall, <i>Glyptograptus posterus</i> Koren et Tzaj, <i>Conochina micracantha</i> Eis., <i>Tanuchitina ontariensis</i> Jans., <i>Amorphognathus ordovicicus</i> Br. et M., <i>Scabbardella altipes</i> (Henn.), <i>Amolodus triangulatus</i> Br. et M., <i>Panderodus infermedius</i> (Br. et M.), <i>Protopanderodus insculptus</i> (Br. et M.), <i>Secuicollacta ornata</i> G., Um. et I., <i>S. scepri</i> McDon., <i>S. silex</i> G., Um. et I., <i>Kalimnaspheera cf. maculosa</i> Web. et Bl., <i>Borisella subulata</i> (Web. et Bl.), <i>Protoceraotikiscum chinocrystalum</i> G., Um. et I., <i>Inanigutta complanata</i> Naz.</p> <p><i>Catenipora workmanae</i> Flom., <i>Rhabdotetradium</i> sp., <i>Thebesia ex gr. thebesensis</i> Amsd., <i>Hirnantia</i> aff. <i>noixella</i> Amsden, <i>Alispira praegracilis</i> Severg., <i>Brevilamnuella ex gr. Thebesensis</i> (Sav.), <i>Prostricklandia</i> ? sp.</p> <p>300 m</p>
Sandbian	Bugryshikh	Khankhara	Khankhara
		<p>Greenish-gray siltstone, sandstone, layers of limestone, at the base of formation - oolitic.</p> <p><i>Glyptorthis praepulchra</i> Severg., <i>Triplasia ainea</i> Severg., <i>Brontocephalina</i> sp., <i>Holotrachelum morinichensis</i> Petrun., <i>Pleurograptus linearis</i> (Carr.), <i>Boreadorthis togoensis</i> Severg., <i>Chaunistomella amzassensis</i> (Severg.), <i>Ceraurinus chanchianus</i> Petrun., <i>Chasmodonella</i> sp., <i>Isotoides</i> sp., <i>Climacograptus tubiliferus</i> Lapw., <i>Diplograptus compactus</i> E. et W., <i>Glyptograptus euglyphus</i> (Lapw.), <i>Dicellograptus</i> sp., <i>Leptograptus</i> sp., <i>Pseudobelodina</i> sp., <i>Onniella chanchianica</i> Severg., <i>Bimuria bugryschichensis</i> Severg., <i>Plectacamara yuscughensis</i> Severg., <i>Paurorthis altaica</i> Severg., <i>Plectorthis altaica</i> Severg., <i>Bronteopsis altaica</i> Severg., <i>Retiograptus geinitzianus</i> Hall, <i>Tretaspis</i> sp.</p> <p>up to 350 m</p>	<p>Gray quartz sandstone, siltstone, clayey shales, limestone.</p> <p><i>Orthograptus quadrimucronatus</i> (Hall), <i>Diplocanthograptus caudatus</i> (Lapw.), <i>Anaptombonites ex gr. grayae sibirica</i> Severg., <i>Leptograptus</i> sp., <i>Cryptograptus tricornis</i> (Carr.), <i>Cl. bicornis</i> (Hall), <i>Eobolodina cf. fomicata</i> (Staaf.), <i>Hesperorthis</i> sp., <i>Leptellina</i> sp.</p> <p>160 m</p>
Darriwilian	Kostinsky	Bugryshikh	Bugryshikh
		<p>BUGRYSHIKHA FM.</p> <p>Dark-gray sandstone and siltstone, in lower part - conglomerate with quartz pebbles.</p> <p><i>Cl. antiquus lineatus</i> E. et W., <i>Dip. multident</i> E. et W.</p> <p><i>Reteograptus geinitzianus</i> Hall, <i>Cybelurus planifrons</i> (Web.), <i>Raymondella bugryschichensis</i> Petrun., <i>Apatomorpha altaica</i> Severg., <i>Leptellina tennesseensis</i> Coop., <i>Isophragma extensum</i> Coop., <i>Homotelus angustus</i> Petrun., <i>Lonchodomas (Foliopyge) levis</i> Petrun., <i>Ampyxella (Belaxella) infermicostata</i> Petrun., <i>Nileus tengrensis</i> Web., <i>Telephina mobergi</i> (Hadd.), <i>Hustedograptus teretiusculus</i> (His.), <i>Leptograptus</i> sp., <i>Beloviella bugryschichensis</i> Severg., <i>Glyptorthis primus</i> Severg., <i>Cybelurus planus</i> Lev.</p> <p>&gt;1200 m</p> <p><i>Amplexograptus coelatus</i> (Lapw.), <i>Expansograptus jakovlevi</i> (Keller)</p>	<p>BUGRYSHIKHA FM.</p> <p>Dark-colored sandstone, siltstone, in lower part - conglomerate with quartz pebbles.</p> <p><i>Isophragma</i> sp., <i>Rostricellula</i> sp., <i>Diplograptus multident</i> E. et W.</p> <p><i>Acrograptus serratus</i> (Hall), <i>Glossograptus hincksi</i> (Hopk.), <i>Glos. fimbriatus</i> (Hopk.), <i>Climacograptus aff. brevis</i> E. et W., <i>Am. perexcavatus</i> (Lapw.), <i>Pseudoclimacograptus scharenbergi</i> (Lapw.), <i>Hustedograptus teretiusculus</i> (His.)</p> <p><i>Glyptograptus tricornis insectiformis</i> Rued., <i>Amplexograptus coelatus</i> (Lapw.), <i>Glossograptus fimbriatus</i> (Hopk.)</p> <p>137 m</p>
Dapingian	Kuibyshevo	Tuloi (= Lebed')	Tuloi (= Lebed')
		<p>SUETKA FM.</p> <p>Gray-green and purple polymictic sandstone, siltstone, clayey shales, conglomerate.</p> <p>~2500 m</p>	<p>KOSTINSKY BEDS</p> <p>Siltstone, limestone with <i>Kolymella cf. plana</i> (Tchug.), <i>Eccoptochile tchagryica</i> Petr., <i>Ceraurina cf. frequens</i> Tchug., <i>Ateleasma batunensis</i> Sev., <i>Eorobergia cf. bipunctata</i> Tchug., <i>Idiostrophia costata</i> Coop., <i>Plomerrulus amplissimus</i> Petrun., <i>P. parasensis</i> Petr., <i>Bathyrulius nonnullus</i> Tchug., <i>Glaphurus altaicus</i> Web., <i>Eccoptochile tchagryica</i> Petr., <i>Plectacamara cf. constata</i> Coop., <i>Archaeorthis altaica</i> Severg., <i>Juanognathus jannussoni</i> Serp., <i>Parapanderodus striatus</i> (Gr. et El.), <i>Coopernathus cf. nyini</i> (Cooper), <i>Acodus eletsicus</i> Tolm. et al.</p> <p>~10 m</p> <p>VOSKRESENKA FM.</p> <p>Gray-colored and variegated siltstone, sandstone, rare mudstone, in basal layers - lenses of gravelite, conglomerate.</p> <p><i>Desmochitina minor</i> Eis., <i>D. rhenana</i> Eis., <i>Conochitina</i> sp., <i>Iliaenus</i> sp., <i>Temnodiscus</i> sp.</p> <p>70-225 m</p>
Tremadocian	Takoshkin (= "Upper Tayanza")	GORNY ALTAI SERIES (upper part)	GORNY ALTAI SERIES (upper part)
		<p>SUETKA FM.</p> <p>Gray-green and purple polymictic sandstone, siltstone, clayey shales, conglomerate.</p> <p>~2500 m</p>	<p>SUETKA FM.</p> <p>Variegated polymictic sandstone, siltstone, clayey shales, gravelite.</p> <p>~3000 m</p>

Fig. 4. Continued.

Correlation of local stratigraphic units														
G o r n y   A l t a i														
Stage	Horizon	Subhorizon	Charysh-Inya Zone				Kharlov Zone	Vydrikhha Zone	Talitsa Zone					
			South region						Central region					
			Deep water inner shelf						Deep water oceanic basin		Deep water oceanic basin			
			Gromotukha R.						Kamyshenka R.		Vydrikhha, Vyatchikhha riv.		Marcheta, Marchetenok rivers	
			4 c						5		6		7 a	
Hir- nant	List- yan	<div>TEKHTE' FM. Massive bedded and clayey limestone, siltstone and mudstone. <i>Mesofavosites dualis</i> Sok., <i>Agatolites</i> sp., <i>Heliolites</i> sp., <i>Gromotuchia dilavata</i> Petr., <i>Gr. stricta</i> Petr., <i>Homotelus</i> sp., <i>Trucizetina</i> <i>subrotundata</i> Havi., <i>Dedzetina</i> aff. <i>microstoma</i> Havi., <i>Dalmanella dietkensis</i> Sev., <i>Stegerhynchus concinnus</i> (Savage), <i>Giraldibella bella</i> Bergstrom, <i>Eospirigirina gaspeensis</i> (Coop.), <i>Brevilamnulella</i> <i>gromotuchaensis</i> (Sev.), <i>Plectatrypa</i> aff. <i>henningsmoeni</i> Boucot et Johnson   </div>												





Correlation of local stratigraphic units									
Stage	Horizon	G o r n y   A l t a i							
		A n u i - C h u y a   Z o n e							
		North region				North district of Central region			
		Shallow water inner shelf				Central part of outer shelf with reef massifs			
Hir- nant	List- yan	Bulukhta, Sarasa rivers				Kelei, Muta rivers, Tekhten' Brook			
		8 c				8 d			
Katian	Tekhten'	<p>BULUKHTA FM.</p> <p>Gray limy sandstone, limestone, conglomerate. <i>Glyptorthis balclatchiensis</i> (Dav.), <i>Rostricellula ainsliei</i> amzassica Severg., <i>Triplasia ex gr. ainsa</i> Severg., <i>Onniella</i> sp., <i>Calapoecia</i> sp., <i>Brachyelasma</i> sp., <i>Sibiriolites</i> sp., <i>Procheliolites</i> sp. up to 600 m</p>				<p>VTORYE UTYSY FM.</p> <p>Basal layers, black platy mudstone, analogues of layers with graptolites of <i>persculptus</i> Zone.   </p>			

Fig. 4. Continued.

Stage		Correlation of local stratigraphic units									
		G o r n y   A l t a i									
		A n u i - C h u y a   Z o n e									
		South district of Central region					South region				
Horizon	Subhorizon	Inner part of outer shelf with reef massifs					Inner part of outer shelf with reef massifs				
		Chakyr, Elanda, Ebogon rivers					B. and M. Yaloman, Nizh. and Verkh. Karasu, Saldzhar				
Hir- nant	List- vyan-	8 e					8 f				
							VTORYE UTYOSY FM. Basal layers, limy siltstone, analogues of Beds with graptolites of <i>persculptus</i> Zone. ----- ? ----- 10-20 m				
Katian	Tekhten'	TEKHTEN' FM. Limestone, sandstone, siltstone, mudstone. <i>Pseudoclimacograptus</i> sp., <i>Onniella</i> cf. <i>chancharica</i> Severg., <i>Sowerbyella</i> sp., <i>Nyctopora altaica</i> Dz., <i>N. tchakyrensis</i> Dz., <i>Catenipora</i> sp., <i>Lyopora altaica</i> Dz., <i>Eofletcheria kovalevskiyi</i> Dz., <i>Calapocia baragashiensis</i> Dz., <i>C. lebediensis</i> Dz., <i>Hedstroemia</i> sp., <i>Clathrodiction</i> (?) <i>kirgisicum amzassensis</i> V. Khalf., <i>Lophiostroma elandense</i> V. Khalf. (чакырские слои). Песчаники, алевролиты. <i>Sceptaspis</i> cf. <i>unica</i> Petrun., <i>Lonchodomas</i> sp., <i>Homotelus</i> sp., <i>Ceraurinus</i> sp., <i>Eorobergia</i> cf. <i>crassilimbata</i> Petrun., <i>Dactylogonia subgeniculata</i> Severg., <i>Rostricellula</i> sp. up to 350 m					TEKHTEN' FM. Sandstone, siltstone, clayey and reefal limestone. <i>Palaeofavosites</i> cf. <i>legibilis</i> Sok., <i>Stelodictyon</i> cf. <i>mamilatum</i> (F. Sch.), <i>Mesofavosites shivertiensis</i> Dz., <i>Agetolites</i> <i>insuetus</i> Kim, <i>Plasmoporella convexotabulata</i> Kiaer., <i>Plasmoporella</i> cf. <i>vesiculosa</i> Kiaer., <i>Cyrtophyllum samyshiensis</i> Dz., <i>Wormsipora karasuensis</i> Dz., <i>Grewingia semilunatum</i> (Scheffen), <i>Brachielasma altaica</i> Tcherepn., <i>Ditocholasma canica</i> Tcherepn., <i>Lepidocyloides</i> ex gr. <i>insignis</i> (Sev.), <i>Bellimurina</i> ex gr. <i>incommode</i> Will., <i>Zygospiraella</i> cf. <i>indistincta</i> Kulk. et Severg. > 600 m				
		Sandbian	Khankhara	KHANKHARA FM. Green- and dark-gray siltstone and sandstone, gray limestone. <i>Dicranograptus clingani</i> (Carr.), <i>Pseudoclimacograptus sharenbergi</i> (Lapw.), <i>Bronteopsis gregaria</i> Raym., <i>Jaboganellus gornoaltaicus</i> Petr., <i>Amplocularia detera</i> Petrun., <i>Lonchodomas tchakyrensis</i> Petr., <i>L. tardus</i> Petrun., <i>Homotelus collatatus</i> Petrun., <i>Ceraurinus</i> <i>frontonis</i> Petrun., <i>Remopleurides</i> cf. <i>warburgae</i> Dean, <i>Staurocephalus</i> sp., <i>Paracybeloides</i> cf. <i>loveni</i> (Linnr.), <i>Otarionellina koksoriana</i> Korol., <i>Isoteloides</i> sp., <i>Sivorthis</i> cf. <i>friendsvillensis</i> (Coop.), <i>S. jabogonicum</i> (Severg.), <i>Isophragma</i> cf. <i>extensum</i> Coop., <i>Strophomena</i> sp., <i>Chaulistomella</i> sp., <i>Apatomorpha altaica</i> Severg. up to 700 m					KHANKHARA FM. Siltstone, mudstone with layers and lenses of limestone. <i>Cyrtophyllum kaniensis</i> Dz., <i>Karagomia altaica</i> Dz., <i>Strophomena</i> cf. <i>lebediensis</i> Severg., <i>Catazyga altaica</i> (Severg.), <i>Hesperorthis</i> cf. <i>lebediensis</i> Severg., <i>Bellimurina</i> ex gr. <i>incommode</i> Will., <i>Titambonites</i> cf. <i>elandicus</i> Severg., <i>Onniella</i> <i>chancharica</i> Severg., <i>Chaulistomella amzassensis</i> (Severg.), <i>Jaboganellus parrectilimbus</i> Petr., <i>Sceptaspis</i> ? <i>katunica</i> Petr., <i>Onduella</i> sp., <i>Phaenoporella</i> sp., <i>Polygonograptus</i> aff. <i>marinae</i> Obut, <i>Dictyonema</i> sp. > 700 m		
Darrivilian	Bugryshikha			BUGRYSHIKHA FM. Gray and dark-gray sandstone, siltstone, in the lower part - small-pebble conglomerates.  <i>Telephina möbergi</i> (Hadd.), <i>Lonchodomas (Foliopyge) levis</i> Petr., <i>L. laevisculus</i> (Bill.), <i>Homotelus angustus</i> Petr., <i>Cybelurus</i> <i>planus</i> Lev.; <i>Sivorthis friendsvillensis</i> (Coop.), <i>Leptellina semilunata</i> Will.  <							

Fig. 4. Continued.



Correlation of local stratigraphic units															
Stage	Horizon	G o r n y   A l t a i													
		Anui-Chuya Zone					Biya-Katun' Zone								
		South-east region					West region			Central region					
		Central part of outer shelf with reef massifs					Shallow water inner shelf			Deep water part of intra-island arc basin					
		Chuya, Achik, Bely Bom, Bom Tarlagan rivers					Maly Kamlak, Sema rivers			Anos, Agaira, Karakol rivers					
Hir-nant	List- Kayan	8 g					9 a					9 b			
Katian	Tekhten'	TEKHTEN' FM.  Gray clayey and limy shales, siltstone, sandstone, conglomerate, mudstone including reefal. <i>Plasmoporella crasa</i> Dz., <i>P. convexotabulata</i> Kiaer.  													

Fig. 4. Continued.

		Correlation of local stratigraphic units	
Stage	Horizon	G o r n y   A l t a i	
		Uymen'-Lebed' Zone	
		Outer and inner shelf and delta zone	
Hir- nant	List- vyan- ka	Lebed', Tuloi, Tagaza, Tandoshka, Yurok rivers	
		1 0	
		CHEBOR FM.	
Katian	Tekhten'	Variegated sandstone, siltstone, aleuro-sandstone, shales. <i>Glyptomena subgirvanensis</i> Severg., <i>Austinella</i> sp.	
		> 650 m	
		Upper subformation	
Sandbian	Khankhara	Gray limestone, including algae, biogenic-clastic, stromatolite, oolitic, limy siltstone, mudstone and sandstone.	
		<i>Nyctopora minimalis</i> (Rad.), <i>Calapoecia anticostensis</i> Bill, <i>Karagemia altaica</i> Dz., <i>Wormsipora karasuensis</i> Dz., <i>Sibiriolites lebediensis</i> Dz., <i>Grewinkia lebediensis</i> (Tcherepn.), <i>Glyptorthis praepulchra</i> Severg., <i>Austinella lebediensis</i> Severg., <i>Fardenia scalena</i> Will., <i>Clathrodictyon kirgisicum amzassensis</i> V.Khalif.	
		<i>Phragmodus undatus</i> Br. et M., <i>Panderodus</i> cf. <i>P. gracilis</i> (Br. et M.), <i>Belodina compressa</i> (Br. et M.), <i>Drepanostodus suberectus</i> (Br. et M.), <i>Ballardina altaica</i> Melnik., <i>Eurychilina sennikovi</i> Melnik., <i>Retiprimites formosus</i> Melnik., <i>Laccochilina (Laccochilina) lebediensis</i> Melnik., <i>Bolbina dubia</i> Melnik., <i>Bol. karasuensis</i> Melnik., <i>Soanella petruninae</i> Melnik., <i>Egorovella demissa</i> Melnik., <i>Bullaefurum partitus</i> Melnik., <i>Pseudozygobolbina invisitata</i> Melnik., <i>Allertemmysa (?) incerta</i> Melnik.	
Darrivillian	Bugryshikha	<i>Scandodus</i> sp., <i>Phragmodus undatus</i> Br. et M., <i>Panderodus</i> cf. <i>P. gracilis</i> (Br. et M.), <i>Aphelognathus</i> sp.	
		<i>Boreadorthis togaensis</i> Severg., <i>Triplesia mongolica</i> Tchern., <i>Chaulistomella amzassensis</i> (Severg.), <i>Anoptambonites grayae sibirica</i> Severg., <i>Strophomena lebediensis</i> Severg., <i>Chasmops</i> sp., <i>Ceraurinus icarus</i> (Bill.), <i>Jaboganellus</i> sp.	
		<i>Trematopora propria</i> Jarosh., <i>Rhinidictya lebediensis</i> Jarosh., <i>R. altaica</i> Jarosh.	
Dapingian	Kulib- shevo	<i>Panderodus gracilis</i> (Br. et M.), <i>Belodina compressa</i> (Br. et M.), <i>Phragmodus undatus</i> Br. et M., <i>Eraticodon</i> sp.	
		155-250 m	
		Lower subformation	
Floian	Tuloi (= Lebed')	Alternation of gray-colored limestone, limy siltstone and sandstone, with basal gravelite, rare conglomerate.	
		<i>Fascifera buraensis</i> Severg., <i>Togaella cf. grandis</i> Severg., <i>Eoanastrophia lebediensis</i> (Severg.), <i>Eridorthis subinexpecta</i> Severg., <i>Carolinites</i> sp., <i>Eorobergia lebediensis</i> Petrun.	
		<i>Apatomorpha altaica</i> Severg., <i>Howellites cf. flava</i> (Havl.), <i>Rostricellula stretiniensis</i> Severg., <i>Hemiarges</i> sp., <i>Climacograptus</i> sp., <i>Diplograptus</i> sp. ind.	
Tremadocian	Takoshkin (="Upper Tayanza")	150-250 m	
		KARASA FM.	
		Gray-green aleuro-sandstone, quartz sandstone, siltstone.	
Tremadocian	Takoshkin (="Upper Tayanza")	<i>Hustedograptus teretiusculus</i> (Hisinger)	
		<i>Glyptograptus euglyphus</i> (Lapw.), <i>Diplograptus</i> sp., <i>Amplexograptus coelatus</i> (Lapw.)	
		<i>Amplexograptus confertus</i> (Lapw.), <i>Cyathochitina tuloiensis</i> Obut et Zasl., <i>Conochitina oelandica</i> Eis.	
Tremadocian	Takoshkin (="Upper Tayanza")	<i>Soanella tuloi</i> Melnik., ? <i>Laccoprimitia</i> sp., ? <i>Quadrijugatoriidae</i>	
		<i>Undulograptus dentatus</i> (Brongn.)	
		<i>Parastrophina bilobata</i> Coop., <i>Christiana aff. subquadrata</i> (Hall), <i>Hesperorthis markovae</i> Rozm., <i>Trondorthis sibirica</i> Severg., <i>Isophragma extensum</i> Coop., <i>Glyptomena karasuensis</i> Sev., <i>Ujukites tarykensis</i> Andr., <i>Ceraurina latigenata</i> Petrun., <i>Atryctopyge sibirica</i> Petrun., <i>Cnemidopyge tuloi</i> Petrun., <i>Laccoprimitia</i> sp.	
Tremadocian	Takoshkin (="Upper Tayanza")	<i>Pseudoclimacograptus sharenbergi</i> (Lapw.), <i>Cryptograptus tricornis insectiformis</i> Rued., <i>Eorobergia metopsis</i> Petrun., <i>Carolinites</i> sp., <i>Lonchodomas</i> sp.	
		420-450 m	
		Dark-gray, greenish-gray siltstone, fine-grained sandstone, lilac gravelite and conglomerate.	
Tremadocian	Takoshkin (="Upper Tayanza")	TULOI FM.	
		<i>E. hirundo</i> (Salt.), <i>I. caduceus imitatus</i> Harris	
		<i>Isograptus gibberulus</i> (Nich.), <i>Is. forsipiformis</i> (Rued.), <i>Is. hemicyclus</i> (Harr.), <i>Is. schrenki</i> Obut et Sob., <i>Exp. etensum</i> (Hal.), <i>I. maximo-divergens</i> (Harris), <i>Tetragraptus bigsbyi</i> (Hall.), <i>Eotetragraptus harti</i> (Hall.), <i>Corymbograptus holubi</i> Kraft, <i>Cor. deflexus</i> (El. et W.), <i>Pseudisograptus manubriatus</i> (Hall) <i>Idiostrophia tuloviensis</i> Severg., <i>Archaeorthis altaica</i> Severg., <i>Isophragma ex gr. extensum</i> Coop., <i>Eorobergia integra</i> Petrun., <i>Remopleuridiella altaica</i> Petrun., <i>Vogdesia ? tuloi</i> Petrun., <i>Bodenia aff. aechminiiformis</i> V. Ivan., <i>Bodenia aff. distincta</i> Melnik., ? <i>Tvaerenella</i> sp., <i>Soanella tuloi</i> Melnik., <i>Maraphonia caliginosa</i> Melnik., <i>Pseudophyllograptus angustifolius elongatus</i> Bulm., <i>Phyllograptus anna anna</i> Hall, <i>Expansograptus suecicus</i> (Tullb.), <i>Pendeograptus aff. pendens</i> El., <i>Acrograptus cognatus</i> (Haf. et Thom.) <i>Didymograptus protobifidus</i> Elles <i>Phyllograptus densus</i> Tornq., <i>Phyllograptus ilicifolius glaber</i> Mon. <i>Acr. pussilus</i> (Tul.), <i>Expansograptus extensus</i> (Hall), <i>Exp. taimyrensis</i> Obut et Sob., <i>Nanorthis gloriosus</i> Severg., <i>Hesperonomia paratylyensis</i> Severg., <i>Tersella altaica</i> Petrun., <i>Lapidaria ? ishpensis</i> Petrun., <i>Rhabdochitina turgida</i> lev. <i>Acrograptus balticus</i> (Tullb.)	
Tremadocian	Takoshkin (="Upper Tayanza")	265-600 m	
		<i>Tetragraptus approximatus</i> (Nich.), <i>Eotetragraptus harti</i> (T.S.Hall), <i>Eotet. aff. headi</i> (Salt.), <i>Nanorthis multicostata</i> Ulr. et Coop., <i>Shumardia tagasensis</i> Petrun., <i>Pythine sibirica</i> Petrun., <i>Seleneceme improvisa</i> Petrun.	
		100 m	
Tremadocian	Takoshkin (="Upper Tayanza")	ISHPA FM.	
		Member 3 - variegated siltstone. <i>Apatokephalus ex gr. serratus</i> Sars., <i>Amzasskiella mirabilis</i> Polet., <i>Glaphurus cf. coronatus</i> Z.Max., <i>Nanorthis shoriensis</i> Severg.	
		Member 2 - variegated limestone, sandstone, siltstone. <i>Apatokephalus bijanus</i> Petrun., <i>Symphysurus</i> sp., <i>Adelograptus tenellus</i> (Linn.), <i>Kiaerograptus kiaeri</i> (Mons.), <i>Nanorthis shoriensis</i> Severg.	
Tremadocian	Takoshkin (="Upper Tayanza")	425 m	
		Underlying units	
		€	

Fig. 4. Continued.

		Correlation of local stratigraphic units										Stratigraphic charts for adjacent regions			
Stage	Horizon	G o r n y A l t a i										Siberian platform (Kanygin et al., 2017)  Regional Stratigraphic units			
		Teletskoe Lakeside zone					Ulagan Zone								
		Inner shelf and delta zone					Shallow water part of intra-island acr basin								
		Iogach, Samysh riv., Tozodov, N. Turochackski, Tarlyk brooks, upstream Biya R.					Bolshoi Ulagan R.								
Hir- nant	List- vyan- ka	1 1										1 2		?	
		IOGACH BODY (former "Chebor" Fm.)										SYNTYGAN FM.			
Katian	Tekhten'	At the base red-colored fine-grained sandstone and pebble conglomerate. In lower and middle parts gray, snuffy-gray, green-gray and red-colored sandstone, siltstone, limy-clayey mudstone, ? limestone.										Gray sandstone, siltstone, in the upper part - layers and lenses of limestone and red-colored siltstone, covers of basic volcanics.		BURIAN	
		<i>Schizophorella fallax fallax</i> (Salter), <i>Rostricellula sparsa asiatica</i> Rozman, <i>Rynchotrema</i> sp., <i>Strophomena</i> sp., <i>Cyrtototella</i> sp., <i>Austinella lebediensis</i> Severg., <i>Eridorthis</i> cf. <i>subinexpecta</i> Severg., <i>Glyptorthis balclatchiensis</i> (Dav.), <i>Triplesia mongolica</i> Tschern., <i>Ceraurinus</i> sp., <i>Pterygometopinae</i> , <i>Cyrtophyllum</i> sp., <i>Sibiriolites</i> sp.												NIRUNDIAN	
														DOLBORIAN	
Sandbian	Khankhara	<p>SAMYSH BODY</p> <p>Gray, green-gray and red-colored sandstone, siltstone, limy sandstone, limy siltstone, mudstone, limestone. At the base - red-colored conglomerate and sandstone.</p> <p><i>Sowerbyella sladensis</i> Jones, <i>Cyrtototella semicyrcularis</i> (Eichwald), <i>Lenorthis</i> cf. <i>girardi</i> Andreeva, <i>Rostricellula plena</i> Hall, <i>Hebertella</i> cf. <i>borealis</i> Billings, <i>Syphonotreta</i> sp., <i>Kullervo panderi</i> Opik, <i>Ingria</i> sp., <i>Simphysurus</i> cf. <i>exactus</i> (Tschug.), <i>Asaphus</i> sp., <i>Pliomera</i> cf. <i>insangensis</i> Billings, <i>Sibiriolites</i> sp., <i>Acodus</i> cf. <i>A. combsi</i> Bradshaw, <i>Parapanderodus</i> sp., <i>Drepanoistodus basiovalis</i> (Sergeeva), <i>Paraistodus</i> sp., <i>Periodon</i> sp.</p> <p>TOZODOV BODY</p> <p>Greenish-gray and gray sandstone, siltstone, mudstone, limy mustone, rare lenses of limestone.</p> <p><i>Undulograptus</i> aff. <i>jaroslavi</i> Boucek, <i>Lonchodomas</i> sp., <i>Asaphidae</i>, <i>Egorovella</i> sp., <i>Parapanderodus gracilis</i> (Barnson et Mehl), <i>Yangtzeplacognathus</i> ? sp.</p>										up to 750 m		BAKSANIAN	
Darriwilian	Bugryshikha											KOSBAZHI FM.		KIRENSKIAN-KUDRIAN	
												Red-colored and green clayey gravelite and sandstone, aleuro-sandstone, siltstone, at the base - conglomerate. <i>Angarella lapatini</i> Assat		VOLGIAN	
														MUKTEIAN	
														VIKHOREVIAN	
Dapingian	Kuibyshevo											KYZYL TASH FM.		KIMAIAN	
												Cherry quartz sandstone, siltstone, rare gravelite, conglomerate. In the upper part - cherry clayey shales. <i>Angarella lapatini</i> Assat		~ 600 m	
Floian	Tuloi (= Lebed')											ADYGKHAN FM.		UGORIAN	
												Green and greenish-gray sandstone, conglomerate, purple siltstone.		~ 300 m	
Tremadocian	Takoshkin (= "Upper Tayanza")											ERITAG FM.		NYAIAN	
												Green and purple sandstone, siltstone, conglomerate.		up to 550 m	
Underlying units		Є <sub>1</sub>										?Є		Є <sub>3</sub>	



### 1.1.1. REGIONAL STRATIGRAPHIC UNITS OF THE WESTERN PART OF ASFA

**Takoshkin (=Verkhne-Tayanza) Regional stage (Horizon)** was defined by collective authors (Sennikov et al., 2014, 2018a,b).

The position of the lower boundary of the Ordovician System and, correspondingly, the position of the Tremadocian Stage (Lower Ordovician) were fixed in the Green Point type section on Newfoundland, Canada (Cooper et al., 2001), with the zonal conodont species *Iapetognathus fluctivagus*. The lower boundary of the Tremadocian is now correlated with the lower boundaries of the fossil-bearing Marcheta Formation (the Zasur'ya Group, Gorny Altai). With a low degree of certainty, the lower boundary of the Tremadocian is correlated with the lower boundaries of the nonfossiliferous Suetka and Tekelyu Formations (the Gorny Altai Group, Altai). Hereafter, for younger Ordovician chronostratigraphic intervals, no data are presented on the succession of formations in the Ulagan zone of Gorny Altai: Eritag, Adygkhan, Kyzyltash, Kosbazhi, Pichikhem, and Syntygan (Naumenko and Gutak, 1982). This is explained by the fact that the boundaries of these six formations are tentatively correlated with the lower boundaries of six Ordovician stages of the ISS because of the extremely poor paleontological knowledge of the entire Ordovician section of this part of Altai. According to the old official chart (Decisions..., 1983), the Dobry Horizon was the oldest one in the succession of Ordovician regional stratigraphic units in the western ASFA. This horizon was detected from trilobite assemblages and correlated with the Lower Tremadocian. The Upper Tremadocian included the next horizon - the Tayanza Horizon, substantiated (Decisions..., 1983) by a succession of trilobite assemblages (Petrulina, 1966, 1968, 1973). In recent years, in the Biya-Katun' zone (Gorny Altai), the Middle Kamlak Subformation, correlated on trilobites with the Tayanza Horizon, conodont assemblages (including the Upper Cambrian *Proconodontus* Zone and the lowermost Ordovician *Iapetognathus/Iapetonudus* Zone), which mark the position of the Cambrian/Ordovician boundary in the upper Tayanza Horizon (Sennikov et al., 2014) were found. The Tayanza Horizon on trilobites in the key sections of Gornaya Shoriya was divided into three zones: (1) *Ap. sibiricus/N. oriens*, (2) *Am. mirabilis/Sh. pusillina*, and (3) *Ter. strobilata/Niobe zhulanica* (Petrulina, 1966, 1968). On the other hand, in the Middle and Upper Kamlak subformations of the Biya-Katun' zone (Gorny Altai) and in the Middle and Upper Ishpa subformations of the Uyem'-Lebed' zone, the Tayanza Horizon contains two trilobite assemblages clearly different in stratigraphic position. The first one corresponds to above-mentioned two lower trilobite zones, whereas the upper one corresponds to the third zone. Therefore, the Tayanza regional stratigraphic unit could be accepted in horizon rank and divided into two subhorizons, formally named "Lower-Middle Tayanza" and "Upper Tayanza". The lower unit must correspond to the upper Cambrian (the *Proconodontus* conodont Zone), whereas the "Upper Tayanza" unit must belong to the Tremadocian: the *Iapetognathus/Iapetonudus* conodont Zone and the *B. ramosus/Tr. osloensis/Al. hyperboreus* graptolite Zone (Sennikov et al., 2008, 2014; 2018a,b).

The trilobite species *Harpidoides eximius* Petrun., *Acrocephalina lata* Petrun., *A. contracta* Petrun., *Lusampa cupoides* Petrun., *L. tenuis* Petrun., *Bilacunaspis repentis* Petrun., *B. angusta* Petrun., *Niobella altaiensis* Petrun., *Proapatokephalops altaicus* Petrun., *Plethopeltides (Maximovella) improvisus* Petrun., and *Ishpella repentina* Petrun., which coexist in the Lower Kamlak Subformation with the conodonts of the Upper Cambrian *Proconodontus* Zone, were previously assigned to the Dobry Horizon, correlated with the Lower Tremadocian. According to the presented conodont data, the Dobry Horizon in the western ASFA should be correlated with the Upper Cambrian. Trilobites *Kaufmannella (Butyrinia) robustispina* Petrun. and *Kaltykelina altaica* Petrun., typical of the upper Dobry Horizon and the Lower-Middle Tayanza Subhorizon, are observed in the upper part of the Lower Kamlak Subformation and in the Middle Kamlak Subformation, lower in the section than conodonts *Iapetonudus* sp. These Dobry-Tayanza "transitional" taxa should be assigned to the Late Cambrian. Trilobites *Niobides* cf. *armatus* Harr. et Leanza, *Platypeltoides* cf. *anderssoni* (Troeds.), *Platypeltoides* cf. *wimani* (Troeds.), *Macropyge urceolata* Petrun., *Apatokephalus kamlakensis* Petrun., and *Harpidoides assiensis* Petrun. in the upper part of the Middle Kamlak Subformation were previously assigned to the lower Tayanza Horizon (= "Lower-Middle Tayanza" Subhorizon) and correlated with the lower part of the Upper Tremadocian. As the considered trilobite taxa were found lower in the Kamlak Section than conodonts *Iapetonudus* sp. and *Iapetognathus* sp., the lower part (two-thirds?) of the Tayanza Horizon ("Lower-Middle Tayanza" Subhorizon) should be viewed as the uppermost Upper Cambrian interval in the western ASFA. The trilobite species *Ishpella platycephala* Petrun., observed in all three Kamlak subformations, and the trilobite species *Amzasskiella mirabilis* Polet. in the lower part of the upper Middle Kamlak Subformation and in the Upper Kamlak Subformation within a lower interval than the graptolites of the Upper Tremadocian *ramosus/osloensis/hyperboreus* Zonal Level, previously assigned to the Dobry-Tayanza and Lower- Upper Tayanza "transitional" forms, respectively, should be considered Cambrian-Ordovician "transitional" taxa. In the light of the foregoing, the *Ap. sibiricus/N. oriens* trilobite Zone should be correlated with the Upper Cambrian, whereas the *Am. mirabilis/Sh. pusillina* Zone should occupy a transitional position between Cambrian and Ordovician. The elements of the assemblage of the third trilobite zone of the Tayanza Horizon - the species *Bijaspis katuniana* Petrun., *Parapliomera sibirica* Petrun., *Deltacare sibirica* Petrun., *Euloma shorica* Petrun., *Apatokephalus* ex gr. *serratus* (Sars), *Hysterolesus verus* Petrun., and *Borogothus*

*altaicus* Petrun., observed in the upper part of the Upper Kamlak Subformation, lower than the graptolites of the Upper Tremadocian *B. ramosus*/*Tr. osloensis*/*Al. hyperboreus* Zonal Level, which are assigned to the “Upper Tayanza” Subhorizon and previously correlated with the Upper Tremadocian should be correlated with the chronostratigraphic extent of the entire Tremadocian.

**Tuloi (=Lebed’) Regional stage (Horizon)** was proposed by collective authors (Decisions..., 1983; Sennikov et al., 2014, 2018a,b).

The GSSP of the Lower Ordovician Floian Stage of the ISS was detected in the Diabasbrottet Section, Sweden, and marked by the first appearance of the graptolite marker species *Tetragraptus approximatus* (Bergström et al., 2004). The marker species of the lower boundary of the Floian Stage *T. approximatus* is observed in the Tuloi Formation of the Uymen’–Lebed’ zone of Gorny Altai. Correspondingly, the lower boundary of the Tuloi Formation (Tuloi stratotype section) and the lower boundary of the regional stratigraphic unit based on biostratigraphic data on this formation – the Lebed’ Horizon, are correlated with the base of the Floian Stage of the GSSR. The lower boundary of the Talitsa Formation (Zasur’ya Group) in northwestern Gorny Altai is correlated by conodont assemblages with the lower boundary of the Floian Stage less reliably than the lower boundary of the Tuloi Formation. The *Lagenochitina esthonica chitinozoan* Zone is recognized in the interval corresponding to the Floian Stage, in the GSSP of the younger Dapingian Huanghuachang Stage, China (Wang et al., 2005). The species *Conochitina raymondii* Achab (Achab, 1989) was recognized in this zone on Newfoundland, Canada, and, later, on the Yangtze Plate, China (Wang and Chen, 2003). This species was found in the Tuloi Formation of the Uymen’–Lebed’ zone, Gorny Altai, in the *densus* graptolite Zone in a section near the Tandoshka River (Sennikov and Obut, 2002). It is more complicated to determine the chronostratigraphic position of the upper boundary of the previously used stratigraphic units: the Lebed’ Horizon and Tuloi Formation. Finds of a diverse graptolite assemblage in the new Pridorozhny Section of Altai (Bukolova, 2011), in the uppermost Tuloi Formation of the Uymen’–Lebed’ zone, Gorny Altai, immediately beneath the basal member of the Karasa Formation, require that the upper boundary of the Tuloi Formation (and, correspondingly, the lower boundary of the overlying Karasa Formation) be placed lower, at the boundary between *I. caduceus imitatus* and *U. sinodontatus*/*Cardiograptus* subzones of the *E. hirundo* graptolite Zone. Previously (Decisions..., 1983), the upper Tuloi Formation was correlated with the Kostinsky and lower part of Bugryshikha horizons. This contradicted brachiopod data (Kul’kov and Severgina, 1989) on the presence of the species *Trondorthis sibirica* Severg. in the type section of the Kostinsky beds near former Batun Village, in the lower Karasa Formation, which overlay the Tuloi Formation. With accepting an older age for the upper boundary of the Tuloi Formation (and, correspondingly, the lower boundary of the Karasa Formation), the brachiopod assemblage with *Trondorthis sibirica* Severg. occupies the same stratigraphic position both in the Kostinsky beds of the Voskresenka Formation (the Charysh–Inya zone of Gorny Altai) and the lower part Karasa Formation (the Uymen’–Lebed’ zone of Gorny Altai). When the Lebed’ Horizon was recognized, it encompassed (including the Lebed’ stratotype section near Stretinka Village) the Tuloi Formation of the Uymen’–Lebed’ zone, Gorny Altai, without its uppermost part (Decisions..., 1983). To prevent uncertainty in understanding the phrase “without its uppermost part,” the stratigraphic extent of the Lebed’ stratotype Section should be increased (by less than one-third) to that of the entire Tuloi Formation. Previously (Decisions..., 1983), the Lebed’ Horizon was correlated with the Arenigian (a Stage of the British scale) and did not extend beyond the Lower Ordovician. In the proposed new version, the Lebed’ Horizon in its stratotype coincides with the Tuloi Formation. In the new stage division of the GSSP, the Tuloi (=Lebed’) Horizon will correspond to the Lower Ordovician Floian Stage and three-quarters of the Middle Ordovician Dapingian Stage.

Tuloi (=Lebed’) Horizon is characterised by graptolites zones: *approximatus*, *densus* (*balticus* and *densus* subzones), *protobifidus*, *angustifolius elongatus* – *broggeri*, *gibberulus* (*deflexus* and *maximo-divergens* subzones), *hirundo* (*caduceus imitatus* Subzone).

**Kuibyshevo Regional stage (Horizon)** was proposed by collective authors (Sennikov et al., 2014, 2018a,b). The Dapingian GSSP is the Huanghuachang section, China, which is marked by the first appearance of the conodont species *Baltoniodus triangularis* (Wang et al., 2005). The Huangnitang Section, China, marked by the first appearance of the graptolite species *Undulograptus austrodentatus* (Mitchell et al., 1997), was selected as the Darriwilian GSSP. These two boundaries do not coincide with any of the boundaries of the fossil-bearing Ordovician local stratigraphic units of the western ASFA. With accepting an older age for the upper boundary of the Tuloi Formation, Gorny Altai (see above), a large stratigraphic gap has been detected from graptolite zonal stratigraphy between the upper boundary of the Tuloi (=Lebed’) Horizon and the lower boundary of the overlying Kostinsky Horizon, according to the official Ordovician stratigraphic chart of the ASFA (Decisions..., 1983). The convergence of these two neighboring horizons, required by the stratigraphic chart of 1979 (Decisions..., 1983), was not confirmed in individual sections. In the zonal scale of graptolites from the stratotypes of the considered horizons, the Tuloi (=Lebed’) Horizon has “terminated” (zonal dating of the upper boundary of the horizon), whereas the Kostinsky Horizon has not “begun” (zonal record

of the lower boundary of the horizon). No actual relationships between the horizons (more precisely, neighbouring peculiar faunal assemblages) were observed in the Altai or Salair sections: The stratotype of the Tuloi (=Lebed') Horizon with the Tuloi (=Lebed') assemblage of benthic fauna is localized in the Uymen'-Lebed' zone of Gorny Altai (Lebed' Section), whereas the type sections of the Kostinsky Horizon with a peculiar assemblage of benthic fauna are localized in the Charysh-Inya zone of this region (the Barany Section in the upper Voskresenka Formation near Ust'-Chagyarka Village and the Batun Section near former Batun Village). In the Uymen'-Lebed' Zone of Gorny Altai, the Tuloi Formation (including the stratotype of the Lebed' Horizon near the Lebed' River) is overlain by the basal sandstone member of the Karasa Formation, which does not contain any fossils. Also, they are very few in higher members of the lower Karasa Formation. The Kostinsky beds of the upper Voskresenka Formation (including the formation stratotype near the Barany Creek) in the Charysh-Inya zone of Gorny Altai are underlain by nonfossiliferous sandstone and siltstone members. In connection with these two facts, note that no sections with a continuous succession of benthic faunal assemblages are known from upper Tuloi (=Lebed') Horizon to lower part of Kostinsky Horizon in Gorny Altai or Salair Ridge. The data on the pelagic graptolite assemblages in their synthesized Altai zonal succession, compared with the data from the stratotypes of the Tuloi (=Lebed') and Kostinsky horizons, clearly indicate the presence of a chronostratigraphic interval between the termination of the former and the beginning of the latter. A new horizon is required for a continuous chronostratigraphic succession of regional units; e.g., the Kuibyshevo Horizon, after Kuibyshevo Village, Gorny Altai, with a stratotype in the Maralikha-1 Section, in the Charysh-Inya zone of the region, which contains graptolite assemblages showing a continuous zonal succession (Sennikov et al., 2008). The new horizon includes the *U. sinodentatus*/*Cardiograptus* Subzone of the *E. hirundo* graptolite Zone and the *U. austrodentatus* graptolite Zone. The stratotype section of the Kuibyshevo Horizon contains the graptolite assemblage of the *I. caduceus imitatus* Subzone of the *E. hirundo* graptolite Zone (Sennikov et al., 2008), which corresponds to the uppermost Tuloi (=Lebed') Horizon. Also, the Maralikha-1 Section contains some specimens from benthic groups: trilobites, brachiopods, nautiloids, and crinoids, which might permit determining the possible benthic faunal assemblage for the Kuibyshevo Horizon. In the new stage division of the GSSP, the Kuibyshevo Horizon will correspond to the upper quarter of the Dapingian and the lower quarter of the Darriwilian.

**Kostinsky Regional stage (Horizon)** was defined by E.S. Levitsky (Levitsky, 1963; Severgina, 1973; Sennikov et al., 1982; Decisions..., 1983; Sennikov et al., 2014, 2018a,b). It is aligned with the upper half of the lower Darriwilian (Da1) and middle Darriwilian (Da2).

In connection with new finds of graptolite assemblages in the type locality of the Kostinsky Horizon (the Batun Section near former Batun Village) near the Kostinsky mine in the Charysh-Inya zone of Gorny Altai, the stratigraphic extent of the Kostinsky Horizon should be increased (by less than one-third) to include two graptolite zones: *U. (= ?Eogl.) dentatus* and *E. balhaschensis*/*E. kirgisicus*. Previously, the Kostinsky Horizon included the Lower Llanvirnian and corresponded to the interval of the now-invalid *D. bifidus* graptolite Zone (Decisions..., 1983); later, it was correlated (Sennikov, 1996) with its chronostratigraphic analog – the *E. balhaschensis*/*E. kirgisicus* graptolite Zone. In the new stage division of the GSSP, the Kostinsky Horizon will correspond to the second quarter of the Dapingian. Analysis of available biostratigraphic data permits more precise positioning of the lower boundary of the Zaichikha Formation in the Berd'-Khmelevka zone of the Salair Ridge. Its base was previously correlated with the lower boundary of the *H. teretiusculus* graptolite Zone (Decisions..., 1983) and, later (Petrunkina and Sennikov, 1986), with the upper boundary of the Arenig (top of the *E. hirundo* graptolite Zone). In the Izyrak Section, which is the stratotype for the Izyrak Formation, underlying the Zaichikha Formation, and the parastratotype for the latter, graptolites of the *E. balhaschensis* Zone were detected in the uppermost Izyrak and the lowermost Zaichikha formations. Graptolites of the *E. jakovlevi* Zone were detected 20 m higher in the section than the base of the Zaichikha Formation. The lower boundary of the Zaichikha Formation should be placed in the upper one-third (?upper quarter) of the *E. balhaschensis* Zone, i.e., below the upper boundary of the Kostinsky Horizon in the middle Darriwilian Stage. Similar new data on the distribution of graptolites permit more precise positioning of the base of the Karastun Formation in the Gur'evsk-El'tsovka zone of the Salair Ridge. Its lower boundary was correlated with the lower boundary of the *H. teretiusculus* graptolite Zone (Decisions..., 1983). Graptolites of the *E. balhaschensis*/*E. kirgisicus* Zone were detected in the Korovy Prud Section in this part of the Salair Ridge, in the lowermost Karastun Formation. By analogy with the lower boundary of the Zaichikha Formation (see above), this suggests a correlation between the base of the Karastun Formation and the upper part of the *E. balhaschensis*/*E. kirgisicus* graptolite Zone.

The Kostinsky Horizon is characterized by trilobite fauna *Megalaspides sibirica* Petrun., *Eorobergia compacta* Petrun., *Kolymella* cf. *plana* (Tchug.), *Eccoptochile tchagyrica* Petrun., *Ceraurinella* cf. *frequens* Tchug., and brachiopods *Idiostrophia costata* Ulrich and Cooper, *Chaganella* sp., *Hesperonomia tylyensis* Severg., *Hesperonomiella kuznetskiana* Severg., *Beloviella salairica* Severg., *Trondorthis sibirica* Severg., *Tr. talovkiensis* Severg.

Graptolites of the *dentatus-kirgisicus* Zone are known from the Kostinsky Horizon. Regional *dentatus-kirgisicus* Zone is treated as analogue of the *artus* Zone of the British zonal scale and the *dentatus* Zone of the North-American



zonal scale of the International Stratigraphic Scale, that coincides with lower 4b (=Da2) and transitional 4a/4b (Da1/Da2) of the middle part of the Darriwilian Stage (Webby et al., 2004).

Conodonts of the *E. pseudoplanus* Zone aligned with upper subzone of the variabilis zone from the North-Atlantic zonal scale of the International Stratigraphic Scale, that coincides with the upper part of 4a (=Da1) and lower one third of 4b (=Da2) of the middle part of the Darriwilian Stage (Viira et al., 2001; Webby et al., 2004) are recovered from this horizon.

**Bugryshikha Regional stage (Horizon)** was proposed by (Levitsky, 1963; Severgina, 1965, 1968, 1972, 1973; Gintsinger, V.Sennikov, 1967; Stratigraphic..., 1975; Decisions..., 1983; Sennikov et al., 2018a,b). This horizon is aligned with upper Darriwilian (Da3), lower Sandbian (Sa1) and lower half of upper Sandbian (Sa2).

The Sandbian GSSP, localized in the Fågelsång Section, Sweden, is marked by the first appearance of the graptolite species *Nemagraptus gracilis* (Bergström et al., 2000). This boundary is correlated with the lower boundary of the fossiliferous Gur'yanovka Formation of the Uymen'–Lebed' zone, Gorny Altai. The Sandbian Stage is divided into two time slices (Bergström et al., 2009), proposed as “informal” substages: lower and upper. The lower boundary of the upper one is marked by the base of the *Cl. bicornis* graptolite Zone. Reliable data have been obtained that the lower boundary of the Khankhara Horizon, detected (Sennikov et al., 2008) within the Khankhara Formation of the Charysh–Inya zone of Gorny Altai (and the lower boundary of this formation) corresponds to the middle part of the *Cl. wilsoni* graptolite Zone (Sennikov et al., 2011a). A basal member of oolitic limestones is localized within the Malaya Uskuchevka Section (doublet section of the formation parastratotype near the Bol'shaya Uskuchevka River) and the Khankhara Section (formation stratotype) at the base of the Khankhara Formation. In the Malaya Uskuchevka Section (Sennikov et al., 2008), the boundary between the graptolite assemblages of the *Cl. peltifer/Cl. antiquus lineatus* Zone (which underlies the *Cl. wilsoni* Zone) and the *Cl. wilsoni* Zone is much lower than the base of the oolitic limestones in the middle Bugryshikha Formation. This is to emphasize that the lower boundary of the Khankhara Formation must be localized higher in the section than the lower part of the *Cl. wilsoni* graptolite Zone. In the Ebogon Section, in the Anui–Chuya zone of Altai, the first (lowest) finds of the graptolite assemblage of the *Dicr. clingani* Zone (which overlies the *Cl. wilsoni* Zone) are localized considerably higher than the basal member of oolitic limestones, in the upper Khankhara Formation. This indicates that the lower boundary of the Khankhara Formation must be localized lower than the lower boundary of the *Dicr. clingani* graptolite Zone. The lower boundary of the Khankhara Horizon (and, correspondingly, the base of the Khankhara Formation) should be correlated with the lower boundary of the *Cl. bicornis* graptolite Zone (Teilzone including the upper part of the *Cl. wilsoni* Complexes Zone), which will correspond to the base of the “informal” Upper Sandbian Substage in the new stage subdivision of the GSSP. Conodonts from two interbeds of lumpy and clayey limestones with abundant shell fragments are observed at the lower boundary of the Khankhara Horizon from the section of the middle Gur'yanovka Formation, which is localized on the right bank of the Biya River, near the Chechenek Creek (the Biya Section in the Uymen'–Lebed' zone of Gorny Altai. The conodonts are of medium preservation, and their assemblage includes the species *Panderodus gracilis* (Branson and Mehl), *Phragmodus undatus* Branson and Mehl, and *Erraticodon* sp., typical of the Upper Sandbian *Belodina compressa* and *Phragmodus undatus* zones. Analysis of the biostratigraphic data makes it possible to revise the chronostratigraphic position of the lower boundary of the Veber Formation of the Gur'evsk–El'tsovka zone, Salair Ridge. In the official stratigraphic chart (Decisions..., 1983), the base of the Veber Formation, according to the data from regional correlations of benthic communities, was placed at the base of the *Pl. linearis* graptolite Zone (correlated with the base of the Ashgillian). A brachiopod assemblage with *Boreadorthis togaensis* Severg. was detected in the lower Veber Formation, near El'tsovka Village (Bobrovka limestones). This species is typical of the lower part of the Upper Gur'yanovka Subformation in the Lebed' River Section in the Uymen'–Lebed' zone of Gorny Altai (Kul'kov, Severgina, 1989). The boundary between Lower and Upper Gur'yanovka subformations, based on finds of conodonts (Lebed' and Biya sections) *Scandodus* sp., *Phragmodus undatus* Br. et M., *Panderodus* cf. *P. gracilis* (Br. et M.), *Belodina compressa* (Br. et Mehl), and *Aphelognathus* sp., is now correlated with the lower boundary of the Khankhara Horizon. Thus, with the present state of knowledge, the lower boundary of the Veber Formation of the Gur'evsk–El'tsovka zone of the Salair Ridge should be correlated with the base of the Khankhara Horizon, i.e., with the base of the *Cl. bicornis* graptolite Zone. In the new stage division of the GSSR, the base of the Veber Formation will correspond to the base of the “informal” Upper Sandbian Substage. On the other hand, note that trilobites *Tretaspis* sp. were found in the upper Karastun Formation, which underlies the Veber Formation, in the Karastun stratotype Section, on Mt. Orlinaya, near Gur'evsk, in the Gur'evsk–El'tsovka zone of the Salair Ridge. Trinucleid trilobites, which this species belongs to, are found in Great Britain in the lower Pushgillian Subdivision of the Ashgillian Stage, correlated with the *Pl. linearis* graptolite Zone (Koren', 2002). In the Charysh–Inya zone of Gorny Altai, in the Chineta Section, *Tretaspis* sp. were found in the lowermost Khankhara Formation (Sennikov et al., 2008). This suggests that the base of the Veber Formation (= top of the Karastun Formation) must be localized above the base of the Khankhara Horizon. In the Gur'evsk–El'tsovka zone of the Salair Ridge, the Karastun Formation

is localized below the Veber Formation. It is now the most natural to correlate the upper boundary of the Karastun Formation, which is exclusively terrigenous, with the base of the Khankhara Horizon, i.e., with the base of the *Cl. bicornis* graptolite Zone, by analogy with the upper boundary of the Bugryshikha Formation of terrigenous rocks in the Charysh–Inya zone of Gorny Altai. The Salair Veber Formation, observed (including its stratotype) in the outskirts of Gur'evsk, is underlain by sediments with basal conglomerates (like the previously recognized Gorny Formation, which is now included in the Veber Formation) and, sometimes, unconformity (Gintsinger, 1969). Some researchers (Gintsinger, 1969; Stratigraphic..., 1975; and others) were right in saying that the sedimentation of the Veber Formation might have been preceded by a gap. This gap might have been included the lower Khankhara Horizon – the interval of the *Cl. bicornis* graptolite Zone, i.e., the Upper Sandbian of the GSSP.

The Bugryshikha Horizon is subdivided into two subhorizons. The Lower Bugryshikha Subhorizon is characterized by trilobite fauna *Eorobergia integra* Petrun., *Vogdesia? tuloica* Petrun., *Remopleuridiella altaiensis* Petrun., *Levirobergia oirotica* Petrun., and brachiopods *Archaeorthis altaica* Severg., *Idiostrophia tuloviensis* Severg., and Upper Bugryshikha Subhorizon by trilobites *Pliomerellus latus* Petrun., *Raymondaspis altaicus* Petrun., *Robergiella margofera* Petrun., *Cnemidopyge tuloica* Petrun., *Atractopyge sibirica* Petrun., *Ceraurina latigenata* Petrun., and brachiopods *Glyptorthis primus* Severg., *Parastrophina bilobata* Cooper, *Beloviella bugryshichaensis* Severg., *Christiania* aff. *subquadrata* (Hall), *Glyptomena karasuensis* Severg.

The lower half of the Upper Bugryshikha Subhorizon is dominated by trilobites *Homotelus angustus* Petrun., *Lonchodomas (Foliopyge) levis* Petrun., *Ampyxella (Belaxella) infermicostata* Petrun., *Nileus tengriensis* Web., *Telephina mobergi* (Hadd.), and the upper half of the Upper Bugryshikha Subhorizon – by trilobites *Cybelurus planifrons* Weber, *Raymondella bugryshichiensis* Petrun. On a whole brachiopods *Apatomorpha altaica* Severg., *Leptellina tennesseensis* Cooper, *Hesperorthis markovae* Rozman are distributed in the Upper Bugryshikha Subhorizon.

Graptolites of the *jakovlevi-coelatus*, *teretiusculus*, *gracilis-serratulus*, *multidens* (Subzone of *antiquus lineatus-peltifer* and lower part of Subzone *wilsoni*) zones are typical for the Bugryshikha Horizon.

**Khankhara Regional stage (Horizon)** was proposed by (Sennikov et al., 2008). It could be aligned with upper half of upper Sandbian (Sa2) and lower Katian (Ka1) (Sennikov et al., 2018a,b).

The Katian GSSP, observed in the Black Knob Ridge section, United States, is marked by the first appearance of the graptolite species *Diplacanthograptus caudatus* (Goldman et al., 2007). This boundary does not coincide with any of the boundaries of the fossil-bearing Ordovician local stratigraphic units in the western ASFA. The Katian Stage is divided into four time slices (Bergström et al., 2009), proposed as “informal” substages: lower, first middle, second middle, and upper. The lower boundary of the second substage (first middle) is marked by the base of the *Pl. linearis* graptolite Zone. The Tekhten' Horizon, detected in the Upper Ordovician of the western ASFA (Sennikov et al., 2008), and, correspondingly, the Tekhten' Formation (with a stratotype in the Anui–Chuya zone of Gorny Altai) (Sennikov et al., 2001) were correlated with the Ashgillian Stage of the British scale, whose base corresponded to the lower boundary of the *Pl. linearis* Zone. However, there is no conclusive evidence that the lower boundary of the Tekhten' Horizon corresponds to the base of the *Pl. linearis* Subzone of the *O. quadrimucronatus* graptolite Zone. In the Ebogon Section (the Anui–Chuya zone, Gorny Altai), the latest (uppermost) finds (Sennikov et al., 2001) of graptolites of the *Dicr. clingani* Subzone (the lower of two subzones of the *O. quadrimucronatus* Zone) are observed 300 m lower than the base of the Tekhten' Formation, which overlies the Khankhara Formation. This only shows that the lower boundary of the Tekhten' Formation is localized above the upper part of the *Dicr. clingani* Zone. In the Khankhara Section (the Charysh–Inya zone, Gorny Altai), graptolites of the *Pl. linearis* Zone were found in the lower Tekhten' Formation (previously, the upper Khankhara Formation (Sennikov et al., 1984)). The lower boundary of the Tekhten' Horizon is tentatively correlated with the base of the *Pl. linearis* Subzone, i.e., the base of the second “informal” Katian substage in the GSSP.

The Khankhara Horizon is subdivided into three subhorizons. Trilobites *Chasmopsella unica* Petrun., *Bronteopsis gregaria* Raum., *Jaboganellus gornoaltaicus* Petrun., *Otarionelliana koksoriana* Korol., *Eorobergia lebediensis* Petrun., and brachiopods *Onniella chancharica* (Severg.), *Plectocamara uscuchiensis* Severg., *Fascifera buraensis* Severg., *Bimuria bugryshichaensis* Severg., *Chaulistomella inaquistriata* (Cooper), *Eoanastrophia lebediensis* (Severg.) are typical for the Lower Khankhara Subhorizon.

Trilobites *Ceraurinus icarus* (Bill.), *Calyptaulax bellatulus* Petrun., *Paracybeloides loveni* (Linrs.), and brachiopods *Boreadorthis togaensis* Severg., *Chaulistomella amzassensis* (Severg.), *Strophomena lebediensis* Severg., *Rostricellula ainsliei amzassica* Severg., *Togaella grandis* Severg. are typical for the Middle Khankhara Subhorizon.

Trilobites *Holotrachelus punctiliosus* Tornø., *Iliaenus oviformis* Warb., *Iliaenus* cf. *septentrionalis* Tchug., *Amphilichas sniatkovi* Weber, *Brontocephalina nuda* (Ang.), *Isocolus sjogreni* Ang., *Chasmops saliricus* Petrun., *Eucrinuroides bobroviensis* Petrun., and brachiopods *Eospirigerina sublevis* Severg., *Austinella lebediensis* Severg., *Salopina uxunaica* (Severg.), *Glyptorthis praepulchra* Severg., *Gl. balclatchiensis* (Dav.), *Hesperorthis lebediensis* Severg., *Dulankarella magna* Ruk., *Catazyga salairica* (Severg.) are typical for the Upper Khankhara Subhorizon.

The following graptolite zones are distinguished for the Khankhara Horizon: upper part of *multidens-wilsoni*, *bicornis*, *clingani-caudatus* and *linearis*.

**Tekhten' Regional stage (Horizon)** was proposed by N.V. Sennikov, Z.E. Petrunina and L.A. Gladkikh (Sennikov et al., 2001). It is aligned with middle (Ka2, Ka3) and upper (Ka4) Katian, as well as with lower Hirnantian (Hi1).

The Tekhten' Horizon is subdivided into three subhorizons. The Lower Tekhten' Subhorizon is characterized by brachiopods *Eospirigerina orloviensis* (Severg.), *Oxoplectia platystrophoides* Schubert et Cooper, *Catazyga inensis* (Severg.), *Catazyga anuensis* Severg., *Eridorthis subinexpecta digna* Severg., *Schizophorella fallax* Salter.

Trilobites *Stenoblepharum warburgae* (Prib.) and brachiopods *Giraldibella bella* (Bergst.), *Thebesia thebesensis* Amsden, *Leangella septata* (Cooper) are typical for the Middle Tekhten' Subhorizon. The Upper Tekhten' Subhorizon is characterized by trilobites *Mucronaspis mucronata* (Brongniart) and brachiopods *Dalmanella testudinaria* (Dalm.), *Zygospiraella indistincta* Kulk. et Severg., *Streptis altosinuata* (Holt.), *Hirnantia* aff. *noixella* Amsden, *Brevilamnulella gromotuchaensis* Severg.

The Hirnantian GSSP, observed in the Wangjiawan North section, China, is marked by the first appearance of the graptolite species *Normalograptus extraordinarius* (Chen et al., 2006). The lower boundary of the Hirnantian Stage is correlated with the lower boundary of the Chebor Formation of the Uymen'-Lebed' zone, Gorny Altai. The Hirnantian Stage is divided into two time slices (Bergström et al., 2009). Note that the lower boundary of the upper "informal" substage is marked by the end of the Hirnantian Isotopic Carbon Excursion (HICE), observed worldwide (Bergström et al., 2006, 2009; Chen et al., 2006; Kaljo and Martma, 2011; Mitchell et al., 2011; Underwood et al., 1997), rather than by the marker species of a graptolite or conodont zone. For the Ordovician sediments of the western ASFA, a new horizon – the Listvyanka Horizon (Sennikov et al., 2008) was proposed within the Upper Hirnantian. It includes the mudstones of the lower Vtorye Utyosy Formation (with a stratotype in the Charysh-Inya Zone of Gorny Altai) with graptolites of the *Nor. persculptus* Zone. It is now proposed that the stratigraphic extent of the Listvyanka Horizon should be increased (by less than one-third), so that it includes the dalmanitinae carbonate member of the upper Tekhten' Formation with trilobites *Mucronaspis mucronata* (Brongniart), which underlies the Vtorye Utyosy Formation. According to the latest data, this member (Sennikov and Ainsaar, 2012), correlated with the lowermost *Nor. persculptus* graptolite Zone, has a considerably increased content of heavy carbon isotopes, which corresponds to the HICE. Thus, the Ordovician Listvyanka Horizon of the ASFA will include the uppermost Tekhten' Formation (its peculiar dalmanitinae terminal member) and the lower Vtorye Utyosy Formation. The Listvyanka Horizon will be correlated with the *Nor. persculptus* graptolite Zone in its full extent. In the ISS, it will correspond to the "informal" Upper Hirnantian Substage. Previously, the dalmanitinae carbonate member belonged to the Tekhten' Horizon. The Tekhten' Horizon will correspond to the Tekhten' Formation without its uppermost part – the dalmanitinae member. Note that the Tekhten' Formation in the Tekhten' Section of the Anui-Chuya zone of Altai does not contain this characteristic dalmanitinae member in the upper part of its stratotype (Sennikov et al., 2001, 2008).

Tekhten' Horizon is characterised by graptolites of the *linearis*, *supernus*, *ornatus*, *pacificus*, *ojsuensis/mimyensis* zones and by conodonts of *ordovicianus* Zone.

**Listvyanka Regional stage (Horizon)** was proposed by (Sennikov et al., 2008, 2014, 2018a,b). It is aligned with uppermost upper Hirnantian (Hi1).

Graptolites of the *persculptus* Zone are common for the Listvyanka Horizon. Along with graptolites chitinozoans *Conochitina microcantha* Eisenack have been recovered. The other fauna at this stratigraphic level on the Gorny Altai are represented trilobites *Dalmanitina mucronata* (Brong.) and brachiopods.

Fig. 4 presents intra-regional correlation charts of local Ordovician stratigraphic units of the Gorny Altai based mainly on synthetic modern paleontological data.

## 1.2. ORDOVICIAN GRAPTOLITE ZONES OF THE GORNY ALTAI

The Ordovician sediments of the Gorny Altai contain the following succession of graptolite biostratigraphic units (Fig. 5) (Obut, Sennikov, 1984, Bukolova, 2011; Sennikov et al., 2015a, 2018a,b): Tremadocian, the *kiaeri/tenellus* Zone (the upper half of the zone is regarded as the *ramosus/osloensis* Subzone) (Takoshkin = Upper Tayanza Horizon, Kamlak and Ishpa formations); Floian, the *approximatus* (Tuloi Horizon, Tuloi Formation), *densus* (the lower half is regarded as the *balticus* Subzone) (Tuloi Horizon, Tuloi Formation), and *angustifolius elongatus/broggeri* (without the uppermost part) (Tuloi Horizon, Tuloi Formation) zones; Dapingian, the *angustifolius elongatus/broggeri* (uppermost part) (Tuloi Horizon, Tuloi Formation), *gibberulus* (with the *deflexus* lower Subzone and the *maximo-divergens* upper Subzone) (Tuloi Horizon, Tuloi Formation), and *hirundo* (with the *caduceus imitatus*



ISC, 2008			Time Slises (TS) (Bergstrom et al., 2009)	Correlation of the ISC/GSS stages, horizons, and zonal subdivisions	Altai-Sayan Folded Area					Global correlation levels of lower boundaries of zones and beds with fauna
System	Series	Stage			Horizon (Sennikov et al., 2008)	Salair	Gorny Altai			
Ordovician	Upper	Hirnantian	Index		(Sennikov, 1992, 1996; Sennikov and Bukolova, 2010, Sennikov et al., 2015)	Kuznetsk Alatau, specified after (Obut and Sennikov, 1984; Sennikov, 1996)	Specified after (Bukolova, 2011; Obut et Sennikov, 1984; Sennikov, 1996, 2012; Sennikov et Ainsaar, 2012; Sennikov and Bukolova, 2010; Sennikov et al., 2008, 2011a,b)			
			Hi <sub>2</sub> End of the Hirnantian Isotopic Carbon Excursion (HICE)	11	Listvyanka	persculptus	persculptus / End of the HICE		persculptus / End of the HICE	
			Hi <sub>1</sub> Normalograptus extraordinarius zone (Gr)	10	Tekhten'	Слой с <i>Normalograptus mirnyensis</i> ?	ojsuensis / mirnyensis			
		Ka <sub>4</sub> Dicellograptus complanatus zone (Gr)		Not detected		supernus	pacificus	pacificus		
		Ka <sub>3</sub> <i>Amorph. ordovicicus</i> zone (C)		?			ornatus			
		Ka <sub>2</sub> Pleurogr. linearis zone (Gr)	9	Слой с <i>Orthograptus ex gr. quadrimucronatus</i>		quadrimucronatus	linearis	linearis		
		Ka <sub>1</sub> Diplocanthograptus caudatus zone (Gr)	8	?			clingani	caudatus		
		Sandbian	Sa <sub>2</sub> Climacograptus bicornis zone (Gr)	7	Khan-khara	?		wilsoni	bicornis	bicornis
			Sa <sub>1</sub> Nemagraptus gracilis zone (Gr)			Not detected	foliaceus	peltifer / antiquus lineatus		
				6		?	gracilis / serratus / bekkeri	gracilis / bekkeri		
						teretiusculus	teretiusculus	teretiusculus		
		Middle	Dartwiliian	Da <sub>3</sub> Pygodus serra zone (C)		Kostinsky	jakovlevi / geminus	jakovlevi / coelatus		
	Da <sub>2</sub> Didymograptus artus zone (Gr)				balhaschensis / kirgisicus		balhaschensis / kirgisicus			
	Da <sub>1</sub> Undulograptus austrodentatus zone (Gr)			5			dentatus	dentatus		
	Dapingian		Dp <sub>3</sub> Oncograptus zone (Gr)	4	Kuibyshevo	sparsus	hirundo	sinodentatus / Cardiograptus		
			Dp <sub>2</sub> Isograptus victoriae maximus zone (Gr)					caduceus imitatus		
			Dp <sub>1</sub> Baltoniodus triangularis zone (C)			gibberulus	maximo-divergens	maximo-divergens	maximo-divergens	
								deflexus	gibberulus	
	Floian		Fi <sub>3</sub> Didymograptus protobifidus zone (Gr)	3	Tuloi (=Lebed')	broggeri	angustifolius elongatus / broggeri	protobifidus	protobifidus	
			Fi <sub>2</sub> Oepikodus evae zone (C)			densus	densus	balticus	densus	
			Fi <sub>1</sub> Tetragraptus approximatus zone (Gr)	2		?	approximatus		approximatus	
		Tr <sub>3</sub> Paroistodus proteus zone (C)				Слой с <i>Dictyonema kravtsovi</i> ?	?	ramosus/osloensis / hyperboreus		
Tr <sub>2</sub> Paltodus deltifer zone (C)				Слой с <i>Anis. richardsoni</i> ?		kiaeri / tenellus		Слой с <i>Anis. richardsoni</i>		
Lower	Tremadocian	Tr <sub>1</sub> Iapetognathus fluctivagus zone (C)	1	Takoshtin (=Upper Tayanza)	Грaptолиты отсутствуют	?		Iapetognathus zone (C)		

Fig. 5. Graptolite zonation and correlation levels for the Ordovician of the Gorny Altai and Salair.

lower Subzone and the *sinodentatus/Cardiograptus* upper Subzone) (upper Tuloi–lower Kuibyshevo horizons, Tuloi and Karasa formations) zones; Dartwiliian, the *austrodentatus* (Kuibyshevo Horizon, Voskresenka Formation), *dentatus* (Kostinsky Horizon, Voskresenka and Karasa Formations), *balhaschensis/kirgisicus* (Kostinsky Horizon, Voskresenka Formation), *jakovlevi/coelatus* (Bugryshikha Horizon, Bugryshikha and Karasa formations), and *teretiusculus* (Bugryshikha Horizon, Bugryshikha and Karasa formations) zones; Sandbian, the *gracilis/serratus/bekkeri* (Bugryshikha Horizon, Bugryshikha Formation), and *foliaceus* (= former *multidens* Zone) (with the *peltifer/antiquus lineatus* lower Subzone and the *wilsoni* upper Subzone) (upper Bugryshikha–lower Khankhara horizons, upper Bugryshikha and lower Khankhara formations) zones; Katian, the *quadrimucronatus* (with the *clingani* lower Subzone and the *linearis* upper Subzone) (upper Khankhara–lower Tekhten' horizons, upper Khankhara and lower Tekhten' formations), *supernus*, *ornatus*, and *pacificus* (Tekhten' Horizon, Tekhten' Formation) zones; Hirnantian, the *ojsuensis/mirnyensis* (Tekhten' Horizon, Tekhten' Formation), *persculptus* (Listvyanka Horizon, lower Vtorye Utyosy Formation) (Bukolova, 2011; Obut and Sennikov, 1984; Sennikov, 1976, 1996, 2012, 2013; Sennikov et al., 2008, 2011a, b). Besides, the *protobifidus* Teilzone (Sennikov, 2013; Sennikov et al., 2008), which marks the Upper Floian of the ISC (Section Salair), is distinguished in the Ordovician of Gorny Altai, in the upper part of the *densus* Zone and the lower part of the *angustifolius elongatus/broggeri* Zone. The *bicornis* Teilzone, which marks the Upper Sandbian of the ISC (Section The Taimyr Peninsula), is recognized in the Ordovician of Altai, in the upper part of the *wilsoni* Subzone of the *multidens* Zone. The *caudatus* Teilzone is distinguished in Ordovician sections of Gorny Altai, in the lower part of the *clingani* Subzone of the *quadrimucronatus* Zone. The index species of this zone, *Diplocanthograptus caudatus* (Lap.), marks the Katian Stage of the ISC in the GSSP (Black Knob Ridge Section, United States) (Goldman

et al., 2007). The *caudatus* Zone is distinguished in the Ordovician of Scotland (as a subzone of the *clingani* Zone) (Zalasiewicz et al., 2009) and North America (Loydell, 2012). In northeastern Russia (Kolyma River), finds of the species *Dip. caudatus* (Lap.) (Obut and Sobolevskaya, 1968) in the lower part of the *ingens wellingtonensis* Zone (lower zone of the Katian Stage) allow to define beds with *Dip. caudatus*. The overwhelming majority of the graptolite zones (their assemblages and index species) and beds defined in the Ordovician of Gorny Altai are used worldwide (Cooper et al., 2004; Gradstein et al., 2012; Loydell, 2012; Webby et al., 2004; Zalasiewicz et al., 2009). The exceptions are the *balhaschensis/kirgisicus* Zone, which is described above (Section Salair), and the *jakovlevi/coelatus* Zone. Data on the species *Expansograptus jakovlevi* (Kel.), found in Salair together with *Didymograptus geminus* (His.), are considered above (Section Salair). The species *Amplexograptus coelatus* (Lap.) occurs in the Ordovician sediments of Great Britain (Elles and Wood, 1901–1918; Strachan, 1996, 1997; Zalasiewicz et al., 2009), in the *murchisoni* Zone (probably, also in the uppermost part of the *underlain artus* Zone (former *bifidus* Zone), and spreads to the overlain zones. In Altai the species *Am. coelatus* (Lap.) and the corresponding assemblage are observed in numerous sections below the assemblage of the *teretiusculus* Zone (Sennikov et al., 2008).

### 1.3. ORDOVICIAN CONODONT ZONES OF THE GORNY ALTAI

The first data on conodonts from the Ordovician sediments of Gorny Altai were obtained not long ago (Moskalenko, 1977), but the present state of knowledge of this group (Iwata et al., 1997; Izokh et al., 2003, 2005) is similar to that of conodonts in folded regions such as the Urals and Kazakhstan. As there are not enough monographic studies of the Ordovician conodonts of Altai (Sennikov et al. 2015) and the rare and low-abundance species are not clearly identified, the biostratigraphic units are based on species with a wide geographic range and cosmopolitan taxa, yet allow direct correlations between the proposed zonation and the zones of the ISC/GSS (Fig. 6). Gorny Altai is the only Russian region with detected elements of the zonal taxon *Iapetognathus* and the associated genus *Iapetonodus* (Sennikov et al., 2014) characterize the base of Ordovician in the type sections of North America (Cooper et al., 2001).

The appearance of elements of *Cordylodus angulatus* in the Upper Kamlak Subformation marks the lower boundary of the overlain zone. Beds with *Paroistodus proteus* were defined in the cherts of the upper Talitsa Formation (Zasur'ya Group) of the Takoshkin Horizon and in the cherts of the Marcheta Formation (Zasur'ya Group) of the Tuloi Horizon in its stratotype. The assemblage of the beds includes *Paroistodus* cf. *proteus* (Lind.), *Paracordylodus gracilis*, *Cornuodus longibasis*, *Drepanodus reclinatus*, *Oneotodus* sp., and some other species defined in open nomenclature. The conodont assemblage allows dating this interval as Late Tremadocian – Early Floian. Beds with *Oepikodus evae* were established in the cherts of the upper Marcheta Formation (Zasur'ya Group), Tuloi Horizon. The conodont assemblage in the beds includes *O. evae*, *Periodon* cf. *flabellum*, *Prioniodus* cf. *P. elegans*, *Baltoniodus* sp., and *Drepanoistodus* sp. The beds are correlated with the same-named zone in the upper Floian. The upper Voskresenka Formation (Kuibyshevo Horizon) in its type section contains beds with *Periodon flabellum* – *Parapanderodus striatus*, whose assemblage includes *Semiacontiodus?* *mufushanensis*, *Acodus elatus*, *Juanognathus jaanussoni*, *Protoprioniodus* sp., *Cooperignathus* sp., *Tangshanodus tangshanensis*, *Dr. suberectus* s.l., *Triangulodus larapintinensis*, *Anodontus longus*, *Naimanodus degtiarevi*, *Panderodus?* *nogami*, and others. These taxa permit assigning the considered beds and the Kuibyshevo Horizon to the Upper Dapingian – lowermost Darriwilian. The upper Voskresenka Formation (Kostinsky Horizon) contains beds with *Eoplacognathus pseudoplanus*, whose assemblage includes *Periodon aculeatus*, *Scolopodus* sp., *Drepanodus arcuatus*, *Ansella* sp., and *Paroistodus* sp. (Izokh et al., 2005). The beds are

ISC, 2008			Altai-Sayan Folded Area	
System	Series	Stage	Correlation of the ISC stages, horizons, and zonal subdivisions	Horizon
Ordovician	Upper	Hirnantian		Listvyanka
		Katian		Tekhten'
				Beds with <i>Protopanderodus liripipus</i>
		Sandbian	Base of Katian ↓	Khankhara
	Middle			Bugsy-shikha
		Darriwilian		Kostinsky
		Dapingian		Kuibyshevo
	Lower		Base of the upper part of Floian ↓	Tuloi (=Lebed')
		Floian	Base of Floian ↓	
		Tremadocian	Base of Tremadocian ↓	Takoshkin (=Upper Tayanza)
Cambrian	Upper	Batyr-baian		Lower Tayanza

Fig. 6. Conodont zones and correlation levels for the Ordovician of the Gorny Altai and Salair.

correlated with the same-named zone in the lower Darriwilian (Middle Ordovician) of the northwestern East European Platform. Beds with *Ph. undatus* and *Belodina compressa* were defined in the lower and middle Gur'yanovka Formation (Khankhara Horizon). Along with the nominal species, the assemblage includes *Panderodus* cf. *P. gracilis*, *Aphelognathus* sp., *B. compressa* and *Drepanoistodus suberectus*. The coexistence of *Ph. undatus* and *B. compressa* is typical of the upper Sandbian–lower Katian on many continents, including North America and Eurasia (East European Platform); it permits a reliable correlation of the beds defined in the Gur'yanovka Formation with this stratigraphic interval. Beds with *Protopanderodus liripipus* were established in a cherty terrigenous series (Tekhten' Horizon) in the western part of the region. Thin limestone layers contain a conodont assemblage, in which along with the nominal species *Periodon grandis*, *Panderodus* sp., *Decoriconus* sp., *Paroistodus ?mutatus* and *B. compressa* were identified; it allows a correlation of the distinguished beds with the upper Katian.

#### 1.4. LOCAL ORDOVICIAN STRATIGRAPHIC UNITS OF THE GORNY ALTAI

Ordovician strata in the Gorny Altai occur in two genetically different types of sections that record oceanic and shelf deposition.

##### 1.4.1. OCEANIC DEPOSITION

Oceanic Ordovician sections are known in the western and northwestern Gorny Altai (Anui-Chuya and Charysh-Inya facies zones) and consist of the following units.

##### Zasur'ya Group

The Zasur'ya stratigraphic unit was first distinguished as a formation by O.P. Goryainova in 1956 (unpublished evidence). Tikhonov (1956) in his publication referred to Goryainova's definition but applied the name Zasur'ya to the lower subformation of a variegated formation in the western Gorny Altai. The Zasur'ya unit has no stratotype section and its typical locality is in the area northwest of Krasnoshchekovo Village on the right side of the Charysh River, in the catchments of the Zasur'ya, Molchanikha, and Berezovy brooks. Sennikov et al. (2001, 2003) suggested to distinguish the Zasur'ya unit as a group divided into three formations.

According to the present knowledge, the lithology of the Zasur'ya Group consists of clayey-siliceous schists, jade, chert, mudstone, siltstone, sandstone, tuffaceous sandstone, tuff breccias, and gabbro, gabbro-diorite and diorite dikes, and mafic volcanics. According to the chemistry of basalts, the group was deposited in an ocean (Iwata et al., 1997; Buslov et al., 1999, 2000; Sennikov et al., 2003; Safonova et al., 2011). Variolite pillow lavas, aphyre or less often plagioclase-pyroxene and pyroxene-plagioclase porphyry basalts and scarce andesites are of low-K oceanic tholeiite affinity and can be classified as mid-ocean ridge (MORB) or ocean-island (OIB) basalts (Figs 7-11). Basalts are quite rare in the sections of the Zasur'ya Group and exist as 1 to 10 m thick or thicker layers.

Conodont assemblages in the Zasur'ya Group are localized at two intervals: of (1) Late Cambrian (Aksai Stage – Early Batyrbai Substage) and (2) Early Ordovician (Late Tremadocian Substage – Floian Stage). No transitional Batyrbai-Tremadocian conodont zones in the group have been found so far. All conodonts were extracted by hydrofluoric acid dissolution from chert, siliceous mudstone, and other siliceous rocks. The absence of conodont finds from the Cambrian-Ordovician boundary strata in the Zasur'ya Group is implicit evidence that a large part of the sections lack siliceous rocks. This hypothesis is supported by the stacking pattern of the Zasur'ya sections along the Talitsa River where the group shows distinct division into three units, with a purely terrigenous middle unit (the Talitsa Formation) that is almost void of siliceous sediments.


##### Listvenny Formation

The Listvenny Formation, 200 to 300 m thick, is the lowermost unit in the Zasur'ya Group (Sennikov et al., 2001, 2003). Its stratotype section occurs in the Listvenny Brook – Talitsa River divide in the western Gorny Altai. The Listvenny Formation contains, in all its sections, red or more rarely gray siliceous rocks, often with volcanic intercalations, among red and gray terrigenous members. The formation bears conodonts, radiolarians, and siliceous sponge spicules. The conodont age of the Listvenny Formation is defined at five stratigraphic intervals corresponding to the conodont zones: (1) Aksai age, *W. matsushitai* Zone, (2) Aksai age, *M. erectus* Zone, (3) earliest Batyrbai age, *P. muelleri* Zone, (4) Batyrbai age, *E. notchpeakensis* Zone, (5) middle Batyrbai age, *C. minutus* Zone. Relationships of the Listvenny Formation with the underlying rocks remain unknown. The formation is conformably overlain by the Talitsa Formation.






**Fig. 7.** Original/sedimentary field contacts of oceanic light gray to brown chert and basalt in the Lower Ordovician of the Gorny Altai (Marcheta Formation, Charysh River).





**Fig. 8.** The typical aphyric basalt (thin-section) in the Lower Ordovician of the Gorny Altai (from Safonova et al., 2011).



**Fig. 9.** Chemical composition of basaltoids of the Zasur'ya Group (Gorny Altai). a – classification diagram; b, c – discrimination diagrams. Tholeiitic series: TR – rhyolite; TD – dacite; TA – tholeiite andesites; HFT – high-ferrous tholeiite. Calc-alkalic series: CR – rhyolite; CD – dacite; CA – calc-alkali andesites; HMT – high-magnesium tholeiite; BK – komatiite basalt; PK – komatiite; IAT – island-arc tholeiites; MORB – mid-ocean ridge basalts; OIT – ocean-island tholeiites; OIA – ocean-island alkalic basalts; CAB – calc-alkalic basalts; OIB – ocean-island basalts (from Sennikov et al., 2003).

**Fig. 10.** Rare-earth element distribution curves of basaltoids of the Zasur'ya Group. a – MORB-normed, b – chondrite-normed (from Sennikov et al., 2003).





**Fig. 11.** Scheme for the 500 Ma transitional (A) and 470 Ma enriched (B) hot-spot oceanic basalts (Groups 2 and 3) of the Zasukh Group in respect to the thickness of the oceanic lithosphere. At 500 Ma the lavas probably erupted closer to MOR, i.e., under a thinner/younger oceanic lithosphere. This resulted in the relatively high degrees of melting and therefore in a larger portion of incompatible element depleted refractory material in the melt. At 470 Ma, when the plume is located under a thicker but older oceanic lithosphere, the melting column was shorter, the average degree of melting was lower and the melt was more contributed by incompatible element enriched less refractory material of mantle heterogeneities.

C – in the Ordovician the crust of the Paleo-Asian Ocean subducted beneath the SW active margin of the Siberian continent (from Safonova et al., 2011).

### Talitsa Formation

The 400–450 m thick Talitsa Formation, the middle unit of the Zasukh Group, consists of gray (or less often variegated) terrigenous rocks. Its stratotype is located in the middle reaches of the Talitsa River (right tributary of the Charysh River) in the western Gornyy Altai (Sennikov et al., 2001, 2003). The faunal groups found in the formation are radiolarians and siliceous sponge spicules. The Talitsa Formation conformably overlies the Listvennyy Formation and is conformably (?) overlain by the Marcheta Formation.


### Marcheta Formation

The Marcheta Formation, the upper unit of the Zasukh Group, varying in thickness from 600 to 950 m, is composed of alternating red and gray terrigenous and siliceous rocks (Figs 12, 13). The stratotype of the formation was distinguished as two composite sections along the Marcheta River, the left tributary of the Muta River in the northwestern Gornyy Altai (Sennikov et al., 2001, 2003). No reliably proven basalts have been found so far in the paleontologically constrained sections of the Marcheta Formation. The Marcheta rocks include scarce layers of tuff and tuffaceous sandstone. The formation contains conodonts, radiolarians, and siliceous sponge spicules. The conodont age of the Marcheta Formation of the Zasukh Group is defined at three stratigraphic intervals corresponding to the conodont zones: (1) Tremadocian-Floian boundary strata, *P. proteus* Zone, (2) upper part of the Floian, *P. elegans* Zone, and (3) middle Floian, *O. evae* Zone. The Marcheta Formation conformably (?) overlies the Talitsa Formation. The relationship of the formation with the overlying strata remains uncertain.



**Fig. 12.** Massive red chert yielded radiolarians in the Lower Ordovician of the Gornyy Altai (Marcheta Formation, Charysh Section).





**Fig. 13.** Lithology and ranges of fossil taxa from Charysh Section.

The legend (for this and other logs) : 1 – limestone, 2 – massive limestone, 3 – lenses, 4 – algal limestone, 5 – clayey limestone, 6 – sandy limestone, 7 – mudstone, 8 – siltstone, 9 – aleuro-sandstone, 10 – sandstone, 11 – conglomerate, 12 – sodded interval, 13 – breccia, 14 – cherts, 15 – limy sandstone, 16 – limy siltstone, 17 – limy mudstone, 18 – “twisting” structures.



## 1.4.2. SHELF DEPOSITION

Ordovician shelf deposition in the Gorny Altai is known in the western, northwestern, central, northern, northeastern, and eastern parts of the area. There are three main local Ordovician successions that were deposited in different environments of a single basin.

### 1.4.2.1. THE FIRST FACIES TYPE OF SHELF DEPOSITION

**SUCCESSION 1** (Uymen'-Lebed' facies zone) of Ordovician shelf facies comprises the Ishpa, Tuloi, Karasa, Gur'yanovka, and Chebor formations which occur in the northeastern and eastern Gorny Altai and were deposited on a relatively shallow inner shelf proximal to the shore.

#### Ishpa Formation

The Ishpa Formation in the northeastern Gorny Altai (Krivchikov et al., 1976; Decisions..., 1983; Stratigraphic..., 1991) reaches a thickness of 1000 m. Its stratotype section is located along the Ishpa River (left tributary of the Biya River) near Verkh-Biysk Village (Petrulina et al., 1984). The basal member of the formation is conglomerate of pebbles and cobbles of sedimentary, volcanic, and intrusive rocks. The Ishpa Formation is generally composed of mudstone, siltstone, and sandstone, with limestone and conglomerate interbeds. The faunas include algae, trilobites, brachiopods, and graptolites. Graptolite assemblages correspond to the *tenellus* – *kiaeri* Zone. The formation rests upon the Middle-Late Cambrian Tandoshka Formation and under the Tuloi Formation with unconformities along both contacts. Its lower section correlates with the upper Upper Cambrian, the middle part is the Lower Tremadocian, and the upper section is the Upper Tremadocian.

#### Tuloi Formation

The Tuloi Formation, up to 2600 m thick, occurs in the northeastern Gorny Altai. It was originally distinguished (Krivchikov et al., 1976; Decisions..., 1983; Petrulina et al., 1984; Stratigraphic..., 1991) as the lower unit of the Stretinka Formation (Group). The formation is represented by its stratotype section on the right side of the Biya River 2.4 km downstream of the Tuloi inflow and then on the right side of the lower Tuloi River. The formation consists of interbedded mudstone, siltstone, silty sandstone, and sandstone. Its basal member is composed of coarse conglomerate and sandstone with floating lenses of intrusive, volcanic, and terrigenous pebbles. The Tuloi strata contain trilobites, brachiopods, graptolites, and chitinozoans. According to graptolite zonation, the formation spans the *approximatus*, *densus*, *angustifolius elongatus*, *gibberulus* and *hirundo* zones. The formation conformably oversteps the Tremadocian Ishpa Formation and is conformably overlain by the Middle Ordovician Karasa Formation. The Tuloi Formation correlates with the Floian and the Dapingian (without terminal part).

#### Karasa Formation

The Karasa Formation, 450 m thick, is known in the northeastern Gorny Altai. It was originally recognized (Krivchikov et al., 1976; Decisions..., 1983; Stratigraphic..., 1991) as the upper unit of the Stretinka Formation (Group). Its stratotype section crops out on the right side of the Tuloi River downstream of the Karasa River mouth. The formation basal member consists of light gray and yellowish-gray quartz sandstone and the full section comprises interbedded mudstone, siltstone, silty sandstone, and sandstone. It contains trilobites, brachiopods, ostracods, crinoids, hyoliths, orthoceratites, gastropods, graptolites, and chitinozoans. There are two graptolite zones: *dentatus* and *teretiusculus*. The formation lies concordantly over the Tuloi Formation and is erosively overlain by the Gur'yanovka Formation. The Karasa Formation correlates with upper part Dapingian and the Darriwilian.

#### Gur'yanovka Formation

The Gur'yanovka Formation reaching a thickness of 1000 m occurs in the northeastern Gorny Altai. The formation was recognized by V.M. Sennikov (Sennikov, 1962; Stratigraphic..., 1975; Decisions..., 1983) and is represented by a stratotype section on the right side of the Lebed' River upstream of the Bura River mouth, near former Gur'yanovka Village. The formation is composed of siltstone, sandstone, limy-clayey mudstone, and limestone, with the basal member of gravelstone and coarse to medium sandstone, or less often of fine conglomerate. The faunas include tabulate and rugose corals, stromatoporoids, trilobites, brachiopods, ostracods, bryozoans, pelecypods, gastropods, orthoceratites, conodonts, and graptolites. The Gur'yanovka Formation rests on the weakly eroded surface of the Middle Ordovician Karasa Formation and is conformably overlain by the Chebor Formation. The formation conventionally correlates with the Sandbian and the Katian.

#### Chebor Formation

The Chebor Formation, up to 1100 m thick, is known in the northeastern Gorny Altai in the Lebed', Baigol, Biya, and Samysh catchments (northeastern and central Uymen'-Lebed' Synclinorium). The formation was distinguished by V.M. Sennikov (Sennikov, 1962; Stratigraphic..., 1975; Decisions..., 1983) and has its stratotype section on the right side of the Lebed' River upstream of the Ayugozha mouth, the left Lebed' tributary, near Chebor Mountain.

The Chebor section consists of mudstone and siltstone, less abundant sandstone, with thin layers and lenses of gray limestone. The basal member is composed of fine conglomerate and gravelstone with quartz pebble. The fauna finds are brachiopods and tabulate corals. The formation concordantly overlies the Middle Ordovician Gur'yanovka Formation and is discordantly overlain by the Silurian Tochilny Formation (?). The Chebor Formation conventionally correlates with the Hirnantian.

#### 1.4.2.2. THE SECOND FACIES TYPE OF SHELF DEPOSITION

**SUCCESSION 2** (Biya-Katun' facies zone) of shelf facies includes the Kamlak, Khankhara, and Bulukhta formations found in the northern Gorny Altai. They were deposited at the outer-to-inner shelf transition. Some sections of the local stratigraphic units in succession 2 formed in estuaries of large rivers that most likely flew from mountains.

##### Kamlak Formation

The Kamlak Formation up to 2000 m thick is cropped out in the northern Gorny Altai. It was recognized as a separate stratigraphic unit by (Ermikov et al., 1979; Decisions..., 1983; Petrunina et al., 1984; Petrunina, 1990; Stratigraphic ..., 1991; Contributions..., 2001; Sennikov et al., 2008). The formation's composite stratotype section occurs on the left bank of the Maly Kamlak River and along the Tokoshkin Brook, its left tributary. The Kamlak Formation consists of interbedded sandstone, siltstone, limestone, conglomerate, gravelstone, and mudstone and contains trilobites, brachiopods, graptolites, and conodonts. The conodonts correspond to *Cordylodus lindströmi* – *Iapetognathus fluctivagus* zones and graptolites – to the *osloensis* – *ramosus* zone. Contact with underlain and overlain formations is unconformable. The Kamlak Formation is subdivided into three subformations. The Lower Kamlak Subformation (about 120 m thick) is aligned with the upper Cambrian, the most part of the Middle Kamlak Subformation (~440 m) with the uppermost Cambrian; terminal part of the Middle Kamlak Subformation (~15 m) with the lowermost part Tremadocian, and the Upper Kamlak Subformation (up to 1400 m) with the Tremadocian.

##### Khankhara Formation

For data on the Khankhara Formation see below (Succession 3).

##### Bulukhta Formation

The Bulukhta Formation, about 500 m thick, is known on the northeastern periphery of the Anui-Chuya area of the Gorny Altai (Bulukhta and Sarasa rivers). A.B. Gintsinger (Gintsinger, 1958, 1964; Vinkman, Gintsinger, 1967; Stratigraphic ..., 1975; Decisions..., 1983; Sennikov et al., 1995) distinguished it as a separate stratigraphic unit. The formation's stratotype section occurs in the middle reaches of the Bulukhta River (left tributary of the Ulus-Cherga River). It is composed of sandstone, siltstone, mudstone, limestone, and conglomerate and contains tabulate and rugose corals and brachiopods. The formation concordantly oversteps the Khankhara Formation and is discordantly (?) overlain by the Vtorye Utyosy Formation. The Bulukhta Formation correlates with the second half of the Katian.

#### 1.4.2.3. THE THIRD FACIES TYPE OF SHELF DEPOSITION

**SUCCESSION 3** (Anui-Chuya and Charysh-Inya facies zones) includes the Voskresenka, Bugryshikha, Khankhara, and Tekhten' formations, and the basal layers of the lower Vtorye Utyosy Formation found in the western, northwestern, and central Gorny Altai. The deposition occurred far offshore, on a relatively deep outer shelf and on a shallow-marine carbonate platform along the shelf edge at the foot of the continental slope.

##### Voskresenka Formation

The Voskresenka Formation occupies the western and central parts of the Gorny Altai (Sennikov et al., 1979, 1982; Ermikov et al., 1979; Decisions..., 1983; Stratigraphic..., 1991, Contributions..., 2001). Its stratotype section, about 300 m thick, has been documented in the western Gorny Altai at the Barany – Voskresenka brooks divide (left tributaries of the Charysh River) near Ust'-Chagyrka Village. The Voskresenka Formation, 285 m to 900 m thick, consists of sandstone, siltstone, mudstone, less often conglomerate and limestone. The rocks contain trilobites, brachiopods, crinoids, gastropods, graptolites, and conodonts. Graptolites include zonal assemblages of the *approximatus*, *densus*, *gibberulus*, *austrodentatus-hirundo*, and *dentatus-kirgisicus* zones (Petrunina et al., 1984; Sennikov, 1996). The *E. pseudoplanus* conodont zone in the upper part of the Voskresenka Formation (Kostinsky Beds) correlates with the upper subzone of the *variabilis* Zone in the North Atlantic standard of the International Stratigraphic scale: upper part 4a (=Da1) and lower one third 4b (=Da2) of the lower and middle Darriwilian (Webby et al., 2004). The formation lies discordantly over different Cambrian units and is conformably overlain by the Bugryshikha Formation. The Voskresenka Formation correlates with the Floian, Dapingian, and lower and middle Darriwilian of the international scale.

### **Bugryshikha Formation**

The Bugryshikha Formation varies in thickness from 140 m to 1600 m and occurs in the Charysh-Inya and Anui-Chuya areas of the Gorny Altai near Bugryshikha Village. The formation was first distinguished as an unnamed unit by A.A. Nikonov (Nikonov, 1931; Usov, 1936; Stratigraphic..., 1956, 1975; Perfil'ev, 1959; Sennikov et al., 1959; Decisions..., 1959, 1983; Petrunina, Severgina, 1962; Contributions..., 2001); V.A. Kuznetsov (1948) coined the name Bugryshikha Formation. The stratotype of the formation is a composite section near Bugryshikha Village along the Bolshaya and Malaya Uskuchevka rivers, the right tributaries of the Belaya River. The lower part of the formation crops out along the Bugryshikha River (the Belaya left tributary) where it is exposed lying discordantly upon variegated terrigenous rocks of the Suetka Formation of the Gorny Altai Group. The Bugryshikha Formation is composed of sandstone, siltstone, or more rarely mudstone and conglomerate. The Bugryshikha strata bear trilobites, brachiopods, and graptolites. The section spans the graptolite zones: *jakovlevi-coelatus*, *teretiusculus*, *gracilis-serratus*, and *multidens* (*antiquus lineatus-peltifer* and lower *wilsoni* subzones). The Bugryshikha Formation conformably lies over the Voskresenka Formation and under the Khankhara Formation. It correlates with the upper half of the Darriwilian and the lower half of the Sandbian.

### **Khankhara Formation**

The Khankhara Formation, with a variable thickness of 60 to 800 m, is found in the western, northwestern, northern and central Gorny Altai. It was distinguished in 1929 by A.A. Nikonov (Usov, 1936; Stratigraphic..., 1956, 1975; Tikhonov, 1956; Sennikov et al., 1959; Decisions..., 1959, 1983; Contributions..., 2001). Its stratotype section has been designated on the left side of the Malaya Khankhara River near Chineta Village, and a section on the right side of the Malaya Uskuchevka River near Bugryshikha Village is its hypostratotype. The Khankhara Formation consists of sandstone, calcareous-argillaceous siltstone, and mudstone with limestone and marl interbeds. The base of the formation is a prominent lithological marker being composed of gray sandy and clayey limestone or often oolitic limestone occasionally grading into limestone conglomerate. The fauna found in the Khankhara section includes tabulate corals, gastropods, crinoids, trilobites, brachiopods, and graptolites.

The graptolite zonation includes the upper *multidens-wilsoni* Zone and *bicornis*, *clingani-caudatus* and *linearis* zones.

The formation rests conformably upon the Middle Ordovician Bugryshikha Formation and is gradually overlain by the upper Upper Ordovician Tekhten' Formation. The Khankhara Formation correlates with the upper half of the Sandbian and the lower half of the Katian.

### **Tekhten' Formation**

The Tekhten' Formation, from 115 m to 700 m thick, occurs in the western, northwestern, and central Gorny Altai. It was distinguished (Sennikov et al., 2001; Contributions..., 2001) in upper Ordovician strata which make up a single complex of carbonate-terrigenous rocks standing out against the strata above and below. The carbonates are most often of reef origin. Terrigenous rocks are found on the periphery of reefal carbonates, where they replace the flanks of the latter along the strike, as well as inside the carbonate bodies. Reef frameworks are present in different strata but most often are found in the lower and upper parts of the section. Limestone locally predominate throughout the formation thickness (invalid Orlov Formation) or are restricted mainly to the lower layers (invalid Chakyr Formation) while the upper section consists of terrigenous sediments with thin limestone layers (invalid Dietken Formation). In many sections the carbonate-terrigenous proportions are intermediate between the two extremes. The stratotype section of the Tekhten' Formation is located on the right side of the Tekhten' River (right tributary of the Muta River). The formation is composed of limestone with algal bioherms alternating with calcareous siltstone, sandstone, or less often mudstone. The Tekhten' strata contain tabulate and rugose corals, stromatoporoids, trilobites, brachiopods, ostracods, gastropods, crinoids, graptolites, conodonts, scolecodonts, radiolarians, siliceous sponge spicules, and chitinozoans. The graptolite assemblages in the formation correspond to the *supernus* Zone with the *supernus* and *ornatus* subzones. The Tekhten' Formation concordantly lies over the Khankhara Formation and under the Vtorye Utyosy Formation. It correlates with the upper half of the Katian and with the lower half of the Hirnantian.

### **Vtorye Utyosy Formation**

The greatest part of the Vtorye Utyosy Formation consist of black and dark gray mudstone and is aligned with the Llandovery Series of the Silurian. The basal beds Vtorye Utyosy Formation marked by the *persculptus* graptolite Zone of the terminal Ordovician (upper half).



## 2. TRACES OF GLOBAL SEDIMENTARY EVENTS IN THE ORDOVICIAN SECTIONS OF THE GORNY ALTAI

Many global sedimentary events are related to the Cambrian / Ordovician boundary and the lowermost Ordovician. The uniqueness of the Cambrian-Ordovician boundary interval has been inferred from the following evidence:

a) The concentration of numerous global abiotic (sedimentary) events, whereas biotic extinction events were extremely rare. The Late Devonian – Early Carboniferous proves to be another "geo-critical" interval in the Phanerozoic where global sedimentation events sided with global extinction events (Waliser, 1996).

b) The considered boundary is characterized by a revolutionary restructuring of marine communities, which resulted in the formation of complicated pelagic biota structure.

c) The Paleo-Tethys and large-scale displacements of the continental blocks (Dobretsov, 2003, 2011) gave rise to the Caledonian Salair orogeny represented by the Salair and Gorny Altai terranes. The Early Caledonian endo — and exogenous events were manifest as large-scale collision-accretional events, i.e. accretion of the Gondwanian back-arc and island-arc structures and terranes (Altai-Mongolian, Chulyshman etc.) onto the margin of Siberia, which was accompanied by the basins deformation, orogeny, formation of a coarse-clastic (conglomerate, brecciated, olistostromes, produced by submarine landslide, etc.) bodies.

A new active volcanism event (following the large-scale Early-Middle Cambrian) began in the western part of the region (Kuznetsky Alatau, Salair, Altai), with subsequent "closure" of the Altai segment of the Paleoasian ocean and the onset of the Altai-Salair margin of the Siberian craton transition from active to amagmatic passive margin. The onset of typical shelf sedimentation was characteristic of most of the Gorny Altai area (western, northwestern, central and northern parts) (Sennikov et al., 2008; Sennikov et al., 2018a,b).

d) The existence of the Khadar magnetochronostratigraphic superchron in the studied interval (Dobretsov, 2011). Superchron is a polarity interval when the orientation of a planet's geomagnetic field does not change which suggests the absence of geomagnetic reversals (changes in the Earth's magnetic polarity). The three superchrons identified in the Earth's history correspond to the periods of: 490–460, 300–260 and 124–86 Ma.

### 1. Terminal Late Cambrian sedimentation regressive event Lange Ranch Eustatic Event (LREE).

The LREE was designated on the basis on the basis of material from North America and China (Miller, 1984) and is currently understood as coinciding with the base of *Cordylous proavus* conodont Zone.

In the Gorny Altai, terminal Cambrian conodonts of *C. proavus* Zone (end of Batyrbaian) were recovered at the base of the Lower Kamlak Subformation of shelf genesis. However, a lithological marker for the LREE has not been found yet.

### 2. Global Early Ordovician Acerocare regressive event (ARE).

The ARE event was defined in carbonate shelf sequences of North American, Siberian and Chinese platforms and is aligned with the base of the *iapetognathus fluctivagus* conodont Zone and the base of the *Rhabdinopora flabelliformis parabola* graptolite Zone. Previously, the ARE event was believed to correspond to LREE eustatic event (Erdtmann, Miller, 1981).

The ARE event is designated within the Kamlak Formation (lower part of the upper subformation), in the northern part of the Altai shelf basin (Petrulina et al., 1984; Sennikov et al., 2008). A 35 m-thick member of conglomerate identified within the Kamlak Section grades into sandstone along strike. Limestone from the uppermost part of Middle Kamlak Subformation in the Kamlak Section yielded conodonts of the *Iapetognathus fluctivagus* Zone. The ARE event is designated within the Kamlak Formation (Anui-Chuya facies zone) in the northern part of Altai shelf basin, where it coincides with the Lower/Middle Kamlak subformation boundary. At the base of the Middle Kamlak Subformation there are three consecutive 30 m thick conglomerate beds. These conglomerate is represented by middle-sized, poorly sorted, well-rounded pebbles, which occupy 50 % of the rock and include diorite, quartzite, gneiss, and crystalline schist.

### 3. Black Mountain transgressive event (BME).

The event is linked to the base of the *Cordylodus angulatus* conodont Zone and defined in Australia (Miller, 1984). In the sections at Salair region (near Gorny Altai), the BME event was not manifest in the shelf basin. The conodonts found there indicate that the event should have proceeded inside an unified member of reefal limestone of the Tolstochikha Formation.

The transgressive event, BME, in the Gorny Altai was not proved by paleontological data, since conodonts of the *C. angulatus* Zone are absent. However, noteworthy is that within the relatively shallow-water Upper Kamlak



**Fig. 14.** Tremadocian black mudstone in the Kamlak Formation (Kamlak Section).



**Fig. 15.** Terrigenous rock overlain limestone (Ishpa Formation, Ishpa Section).

the *T. osloensis*/*Al. hyperboreus* Zone, stratigraphically fully aligned with the *Kiaerograptus* Zone, were recovered from the sandstone that overlies the Kamlak conglomerate. The Baltoscandian standard *Kiaerograptus* Zone is below the base of the *Par. proteus* conodont Zone, which will make it possible to identify the KCE when graptolites of this zone are found.

#### 6. Global sedimentary regressive «Ceratomyge» event (CRE).

Ceratomyge Regressive event (Erdtmann, 1986) was established in North America and positioned in the upper Tremadocian, middle parts of the *Araneograptus murrayi* graptolite and *Paroistodus proteus* conodont zones, respectively. The CRE regressive event in the Gorny Altai is identified in the marine sediments (fragments of the Paleasian ocean) at the base of the Marcheta Formation (Anui-Chuya facies zone) by the appearing red-color terrigenous rocks and red chert (Sennikov et al., 2001, 2003; Sennikov et al., 2008). Terrigenous rocks from the underlying Talitsa Formation are mainly grey-colored, rarely lilac, and chert are mainly violet and red-brown. In the Marcheta-2 Section, conodonts of the middle part of *Par. proteus* Zone were recorded.

#### 7. Global sedimentary regressive «Basal Arenig BioEvent» (BAgB) (= «Late Tremadoc - Early Arenig Lowstand»).

A global biotic Basal Arenig Bio-Event (BAgB) is defined at the Tremadocian/Floian (Arenigian) boundary (Walliser, 1986), with the biota responding by a dramatic change in the structure of benthic-pelagic communities. The event was assumingly triggered by a global transgression. The BAgB biotic event is aligned with the base of the *T. approximatus* graptolite Zone. In the Altai shelf basin, the biotic BAgB event and, thus, the global transgressive event, is identified in the lower part of the Tuloi Formation (Uimen'-Lebed' facies zone) (Krivchikov et al., 1976; Petrunina et al., 1984; Sennikov et al., 2008; Noskov, 2007). This formation overlies, with dip and azimuthal unconformity and thick (up to 130 m) basal conglomerate, different Cambrian horizons: the Lower Cambrian in the Lebed' Section,

Subformation (Anui-Chuya facies zone) of the Altai paleobasin a "sudden" appearance of deep-water black shales is observed, that could be regarded as BME traces (Fig. 14).

#### 4. Global sedimentary regressive «Peltocare» event (PRE).

Peltocare Regressive event (Erdman, 1986) was established in the Baltic paleobasin and correlated with the lower boundary of the *Adelograptus tenellus* graptolite Zone. The regressive event PRE is defined in the Ishpa Formation (Uimen'-Lebed' facies zone), the Altai shelf basin. It could be recognized by the appearance of sandstone and mudstone in the upper part of formation overlying mainly organogenic limestone from the middle part of formation (Fig. 15) (Krivchikov et al., 1976; Petrunina et al., 1984; Sennikov et al., 2008). Graptolites of the *Ad. tenellus* Zone were identified at the Pereval Section, in the basal layers of the upper Ishpa Formation.

#### 5. Global sedimentary regressive «Kelly Creek» event (KCE).

Kelly Creek Regressive event (Nicoll et al., 1992) was designated in Australia below the base of the *Par. proteus* conodont Zone. In the Gorny Altai, the KCE regressive event was defined in the upper Kamlak Subformation based on the presence of three members of basal conglomerate up to 170 m thick (Petrunina et al., 1984; Sennikov et al., 2008; Noskov, 2007). Conglomerate have large- and medium-size, poorly sorted, well-rounded, pebbles include granite, granodiorite, syenite, diorite, felsite, basalt, quartzite, chert, sandstone, siltstone, limestone. Graptolites of



the Middle Cambrian in the Tagaza section, and the Upper Cambrian in the Tandoshka Section. It also overlies the Lower Ordovician (Tremadocian in the Ishpa and Tuloi sections). Basal conglomerate of the Ishpa Formation have large-, occasionally medium-size, poorly sorted, well-rounded pebbles, which include granite, granodiorite, syenite, diorite, gabbro, tuffs, quartzite, sandstone, hornfel (Figs 16, 17). Graptolites of the *T. approximatus* Zone were recovered at the lower parts of the formation overlying the conglomerates in the Tuloi, Lebed' and Tagaza sections.

#### 8. Global sedimentary transgressive «Mid Arenig Highstand» event (= «Helskjer Drowning Event»).

The identification of the considered transgression event in the Gorny Altai has thus far been disputable. The end of sedimentary hiatus in the Suetka-Kuibyshev, Charysh-Inya and Anui-Chuya facies zones (Gorny Altai Group) and the onset of Ordovician-Silurian deposition of a coarse clastic basal unit of the Voskresenka Formation lying with angular and azimuthal unconformity over Cambrian-Tremadocian sediments could be in a way interpreted as its traces (Sennikov et al., 2008; Sennikov et al., 2018a,b). In some areas, in western parts (in modern coordinates) of the Altai basin (Charysh-Inya facies zone), the middle Floian Stage was marked by inception of sedimentary strata exhibiting characteristic features of shelf deposition (the Voskresenka Formation). Their formation was substantially contributed by the degrading subducted oceanic pelagic sediments (Zasur'ya Group). The oceanic deposition (Marcheta Formation) also continued. Middle Floian oceanic sediments (upper part of the Marcheta Formation) are found in two regions, to the north and east (in modern coordinates) relative to Floian shelf sediments, specifically, the continental slope position is reconstructed from the shelf Floian deposits (northerly direction), suggesting that younger oceanic fragments were consistently attaching to the edge of accretionary complex. A global regression with the sea level dropped by 80–100 m was established within the latest Dapingian – earliest Darriwilian stratigraphic interval.

#### 9. Global sedimentary regressive «Stein Lowstand Event» or «Late Arenig – Early Llanvirn Lowstand».

Large-scale global regression occurred at the end of Dapingian- beginning of Darriwilian, marked by 80–100 m sea level fall (Nielsen, 2003, 2004, 2011; Munnecke et al., 2010; Gradstein et al., 2012). This event is broadly registered within the Middle Ordovician sections across the Siberian Platform (Kanygin et al., 2010).

The “Stein Lowstand Event” global regression is proposed as the major cause of “unexpected” occurrence of the offshore bar quartz sandstone within the studied medium-to-great depth sections of Gorny Altai. Yellow-gray sandstone well sorted and rounded of the Karasa Formation overlay deep-water black mudstone of the Tuloi Formation (Fig. 18).

Traces of regressive Stein Lowstand Event were recorded in the Uimen'-Lebed' paleobasin (Pridorozhny, Tuloi sections) with stratigraphic position within the boundary between graptolite subzones: *caduceus imitatus* and *sinodontatus/Cardiograptus* subzones of the *E. hirundo* graptolite Zone. Well-sorted and well-rounded yellow-grey sandstone of bar origin (Karasa Formation) overlie black deep-water mudstone of the Tuloi Formation.

The lowermost Karasa Formation within the Lebed' (25 m), Pridorozhny (25 m) and Tuloi (80 m) sections is comprised by the basal bed of the medium-well sorted, well-rounded medium-to-coarse-grained quartz sandstone (90 %) and quartzite (10 %) (Noskov, 2007). It is interpreted either as an offshore bar (Sennikov, 1962; Bukolova,



Fig. 16. Conglomerate of basal member of the Tuloi Formation (Tuloi Section) (General view).



Fig. 17. The pebbles in the conglomerate of basal member of the Tuloi Formation (Tuloi Section).





**Fig. 18.** Massive beds of the bar sandstone of the basal member of the Karasa Formation (Tuloi Section).

There are no offshore-bar quartz sandstone documented at the base of the Karasa Formation in the Yurok Section. Hence, it carries no records of the “Late Arenig–Early Llanvirn Lowstand” global event, fixed at the base of the Karasa Formation in the other studied section of northeastern Gorny Altai, possibly due to a comparatively greater depth and distal remoteness of the Yurok Section area from the shore in the Late Arenig–Early Llanvirn time. At the maximum of the sea-level fall, the upper part of the “Yurok uplift” was proposedly still situated below the fair-weather wave base and was not affected by erosional processes. Considering the total sea-level fall estimation at 80–100 m, the top of the uplift was located deeper than 25–50 m even during the maximum of the regression. Consequently, prior to initiation of the “Stein L.E.” (upper third of the Dapingian, the uplift had been localized as deep as 100–150 m. The maximum of the “Furudal Highstand” (upper third of the Darriwilian) placed the top of the uplift 150–200 m below the sea base. A comparatively low thickness (at least 45 m) and specific lithological composition (fine, homogenous, well-sorted terrigenous material) of the lower Karasa Formation within the Yurok Section reveal its offshore remoteness, and depth at 150–200 m, same as for the uppermost Tuloi Formation.

It is of special notification, the Dapingian–Darriwilian stage in the evolution of graptolites, juxtaposed between the “Basal Llanvirn” event (extinction of isograptids and anisograptids in the *gibberulus* graptolite Zone) and “Basal Caradoc” event (didymograptids went extinct at the base of the *peltifer* graptolite Zone), displays no ecological crises (Barnes et al., 1996; Cooper et al., 2004; Webby et al., 2004; and others). Such periods of ecological stability are usually marked by equalization of specimens within the genera both, in the certain locality and within the whole paleobasin. It is the pattern identified in the Yurok Section, where the quantity dominating taxa can not be established within the paleocommunity during the Dapingian–Darriwilian interval. However, the pre-crisis period (*teretiusculus* graptolite Zone) in the Yurok and Lebed’ sections reveals minimum of the equalization, with *Hustedograptus teretiusculus* (His.) being drastically dominant (several dozen of the specimens, where the other taxa being rare and scarce).

The Yurok Section demonstrates, that after the global regression and “Basal Llanvirn” biotic event, accompanied by rapid depletion of taxonomical abundance and decrease of the population densities of the graptolites, the Upper Dapingian – Lower Darriwilian stage was characterized by a comparatively rapid revival of the population densities and taxonomical abundance of the graptolites inhabiting the offshore-remote zones with depths at 150–200 m. At the same time, the comparatively proximal and less-deep (< 100–150 m) zones of the paleobasin (Lebed’, Pridorozhny, and Tuloi sections) demonstrate a considerably decreased rate of the revival.

The overlaying siltstone and mudstone of the Karasa Formation in the Lebed’ (400 m) and Tuloi (370 m) sections, corresponding the *austrodentatus*, *dentatus*, *balhaschensis*, *jakovlevi*, and *teretiusculus* graptolite zones, mirror a gradual transition to the medium-depth zone (100–150 m). The “Late Arenig–Early Llanvirn Lowstand” was followed by a rapid sea level rise, culminated in the global “Furudal Highstand” (Nielsen, 2003, 2004, 2011), also known as the “Late Llanvirn–Caradoc Highstand” (Gradstein et al., 2012), within the *teretiusculus* graptolite Zone. Total sea level rise during the transgression is estimated at 100–150 m (Gradstein et al., 2012).

### 11. Global sedimentary regressive «Vollen Lowstand» event.

The high stand climax (Late Llanvirn–Caradoc Highstand transgressive sedimentation event) was followed by a regression event with a large-scale sea level drop by 100–130 m (Nielsen, 2004, 2011; Munnecke et al., 2010; Gradstein et al., 2012). The basal coarse-grained (conglomerate, gravelite, sandstone with floating gravel and coarse sand) member that belongs to the Gur’yanovka Formation (Uimen’-Lebed’ facies zone), dated to the earliest Sandbian,

2011), or beach deposits (Noskov, 2007). Therefore, the mid upper Dapingian strata of these sections were accumulated within the lower shoreface (< 10 m) and upper shoreface (0–10 m).

### 10. Global sedimentary transgressive «Furudal Highstand» event or «Late Llanvirn – Caradoc Highstand».

The estimated amplitude of water level rise between transgressive Furudal Highstand (Nielsen, 2003, 2004, 2011) and regressive Stein Lowstand Event events is 100–150 m (Gradstein et al., 2012). A specific basal member of “bar” quartz sandstone at the base of the Karasa Formation is absent from the Yurok Section, which implies that the latter bears no evidence of Late Arenig – Early Llanvirn Lowstand global regression Event recorded in other sections of this part of Altai.

bears evidence of the global early Sandbian regression Vollen Lowstand in the Late Ordovician Altai sections, rather than of local break in sedimentation. The basal member of the Gur'yanovka Formation is composed of pebbles and gravel which include jasperoid, granite, porphyrite, diabase, sandstone, and siltstone (Noskov, 2007). While talus composed of a variety of Ordovician, as well as older sediments could be supplied from the areas, which became exposed during the previous regression.

## 12. Global sedimentary regressive «Frognerkilen Lowstand Event» or «Solvang Lowstand Event».


The described below phenomena manifest in the Gur'yanovka Formation sections (Uimen'-Lebed' facies zone) could be regarded as signatures left by the two global early Katian regressions in the Gorny Altai: (i) the appearance of carbonate gravelite layer (calcirudite with unsorted medium-rounded fragments) in the Bura Section (8th member); (ii) a layer of conglomerate with pebbles of limestone in the Tuloi Section (8th member); and (iii) solitary grains of coarse-grained sand (medium-rounded and -sorted) in fine- and medium-grained sandstones of the Biya section (9th member). The early Katian regressions are most likely to have been caused by the first multi-stage Late Ordovician episodes of global glaciation. The chronostratigraphic position of the two events is recorded at the lowest portion of the *Dicranograptus clingani* graptolite Zone for the former and at the topmost part of this zone for the latter. Estimations of the extent of such regressions yielded a 100 m drop in sea level (Nielsen, 2004, 2011; Munnecke et al., 2010; Gradstein et al., 2012). The Ebogon Section (Anui-Chuya facies zone) is composed of black to dark gray shale with graptolites (*Dicranograptus clingani* Zone) and trilobites. These beds formed at depths of 100 m and below are abruptly overstepped by limestone containing oolitic intercalations, which is interpreted as the result of the wave action, being the signatures left by the "Solvang Lowstand Event".

## 13. Global sedimentary maximum regressive event connected with the Late Ordovician Glacial – «Ashgill Lowstand Interval» (= «Hirnantian Lowstand Event»).

A global sea level drop from 80 to 150 m was reported for the latest Ordovician (Late Katian and Hirnantian stages), i.e. during the period of most extensive Late Ordovician Glacial there was, which prompted the appearance of shallow-water sedimentary formations and breaks in sedimentation. In the Gorny Altai, this is evidenced by the gray-colored carbonate and thin clastic sediments of the Gur'yanovka Formation (Uimen'-Lebed' facies zone) sharply replaced by the overlying red-colored, clastic (sandstone and gravelite), extremely shallow-water sediments of the Chebor Formation. The Lebed' Section (Chebor Formation) is featured by: gravel-pebble diamictites; tempestites; oblique and cross-lamination; multidirectional and multi-scale wave and current ripples; traces of high-amplitude wave action (Fig. 19, a-e). The desiccation cracks, ichnofossils and other signatures of tidal zone and coastal plain are characteristic of the Gur'yanovka Glade Section (Chebor Formation) (Fig. 19,f).

## 14. Global Hirnantian sedimentary event (HICE).

A positive carbon isotope excursion pronouncedly distinguished at the latest Ordovician (Hirnantian Stage) with a maximum recorded in the middle part of a specific member with the so-called Hirnantian-Dalmanitian assemblages of benthic fauna which corresponded to either transitional layers between *Nor. extraordinarius* and *Nor. persculptus* graptolite zones, or the lower half of the latter. This excursion reflecting a sharp increase in carbon isotope content developing to



**Fig. 19.** Lithologic manifestation of the Hirnantian Lowstand Event (Ashgill Lowstand Interval) in the Gorny Altai (Lebed' River).






**Fig. 20.** The 3 m thick Dalmanitina Limestone Member in the Burovlyanka Section with the light gray lower unit and the dark gray (beige) upper unit. The ruler is 2.8 m long.

peaks and indicating maximum absolute values for carbon isotopes throughout the Ordovician, was termed “Hirnantian isotope carbon excursion” (HICE).

The HICE abiotic event has thus far been recorded in almost all Late Ordovician sections, including: a) the Global boundary Stratotype Section and Point (GSSP) for the base of the Hirnantian Stage (the Ordovician System), (Wangjiawan North Section in China) (Chen et al., 2006); b) GSSP for the Rhuddanian (the Silurian System) (Dob’s Linn Section in Scotland) (Underwood et al., 1997); c) in the Mirny Section in the Kolyma region in NE Siberia (Russian regional standard of the Hirnantian) (Kaljo, Martma, 2011) and in the key sections of the Ordovician-Silurian boundary stratigraphic interval of numerous geological regions across the continents (Bergström et al., 2006; Kaljo, Martma, 2011; Meidla et al., 2011; Mitchell et al., 2011; Schonlaub et al., 2011 et al.). Continuous Ordovician-Silurian sections with finds of both pelagic assemblages, including graptolites, conodonts and radiolarians, and benthic Dalmanitina fauna were recorded in the Gorny Altai (Charysh-Inya facies zone) (Sennikov, 1998, 2012; Sennikov et al., 2008; Obut, Semenova, 2011; Sennikov et al., 2011b). In the Gorny Altai sections, these complexes allowed a distinct differentiation of the Hirnantian Stage, which corresponds to *Nor. ojsuensis* and *Nor. persculptus* graptolite zones. While studying the Burovlyanka Section (Charysh-Inya facies zone), a key section for the Gorny Altai, carbon isotopes enabled the records



**Fig. 21.** Stratigraphy, selected fossils (after Sennikov et al., 2014) and carbon isotope data of the Burovlyanka section. Note that thicknesses of the stratigraphic units are not to scale, except for the sampled intervals (members 1 and 5). D – Dalmanitina Member (Member 5) (from Sennikov et al., 2015b).

of the HICE sedimentation event in the lowermost part of the Dalmanitina member, corresponding to the lowermost part of *Nor. persculptus* Zone (Sennikov, Ainsaar, 2012; Sennikov et al., 2015b) (Figs 20, 21).



### 3. PALEO GEOGRAPHY OF THE WESTERN ALTAI-SAYAN FOLDED AREA (ORDOVICIAN ALTAI-SALAIR BASIN)

#### 3.1. SEDIMENTARY TYPES AND BIOTAS OF THE ALTAI-SALAIR ORDOVICIAN BASIN

The Altai-Salair Ordovician basin represented shelf continental-margin basin of the Siberian Craton. The so-called Paleo-Asian Ocean was directly linked with the Altai-Salair shelf basin. Its separate fragments could be observed as tectonic blocks in the Gorny Altai area.

Local lithostratigraphic subdivisions (formations) in the Altai-Salair Ordovician basin characterize wide range of sedimentary facies: from shelf to oceanic genesis. Formations could be subdivided into two groups. First is represented by only one rocks association, for example, sandstone and siltstone, confined to a single specific facies. The second group includes formations represented by the diverse rock associations, for example, limestone, mudstone, siltstone and sandstone. Such rocks in the various sections characterize relatively same facies that on the other hand could be assigned to the different paleogeographic environments.

##### 3.1.1. OCEANIC GENESIS

**1. Volcanic-siliceous-terrigenous sedimentary type** (massive, coarse-laminated tuff, tuff sandstone, chert, siliceous mudstone, siltstone, sandstone yielded numerous siliceous sponge spicules, radiolarians and rare conodonts). As an example: Marcheta Formation (Marcheta-2 and Talitsa sections) – Late Tremadocian – Floian.

**2. Siliceous-terrigenous sedimentary type** (massive, coarse-laminated chert, siliceous mudstone, siltstone, sandstone with numerous siliceous sponge spicules, radiolarians and rare conodonts). As an example: Marcheta Formation (Kamyshenka and Charysh sections) – Late Tremadocian – Floian.

##### 3.1.2. CONTINENTAL SLOPE GENESIS

**1. Terrigenous flysch sedimentary type** (coarse-laminated rhythmic sandstone, siltstone, mudstone, with rare but taxonomically diverse trilobites, brachiopods, single poor graptolites). As an example: Bugryshikha Formation (Malaya Uskuchevka and Pichuzhikha-2 sections) – Late Darriwilian – Early Sandbian.

**2. Terrigenous underwater-sliding (gravitation-mixtite) sedimentary type** (non-bedded, often lense with landslide traces and small isolated sphere jointing in terrigenous rocks (“twisting”), chert, siliceous mudstone, siltstone, with rare unvaried radiolarians and rare siliceous sponge spicules). As an example: siliceous-terrigenous sequence (Suetka Section) – Early Hirnantian.

**3. Siliceous-terrigenous (gravitational-mixtite) sedimentary type** (fine-laminated, often lenses with traces of sliding and isolated sphere jointing in terrigenous rocks (“twisting”), chert, siliceous mudstone, siltstone with few and taxonomically monotonous radiolarians and rare siliceous sponge spicules). As an example: siliceous-terrigenous sequence (Suetka Section) – Early Hirnantian.

##### 3.1.3. SHELF GENESIS

The outer shelf, which is quite far from the shore, must have had depths of 150–250 m, that is, greater than those in the inner shelf. It supported a carbonate platform, which occupied at least 70 % of the studied strip. The platform, in its turn, developed reefs, whose tops reached the sea surface (so-called level reefs). About 20–25 % of the outer shelf was occupied by the inner edge of the carbonate platform, with terrigenous-carbonate and carbonate-terrigenous sedimentation; 5–10 %, by the outer edge and bottom of the carbonate platform, with carbonate-terrigenous or, less often, siliceous-terrigenous sedimentation.

**1. Volcanic-terrigenous sedimentary type** (volcanic islands and arcs).

A. Volcanoes slope facies (porphyrite, tuff, sandstone lenses and rare limestone lenses with single trilobites and brachiopods). As an example: Salair, Krasnoe Formation (Section Krasnoe), El'tsovka Formation (El'tsovka Section) – Tremadocian.

B. Facies distant from the volcanic arc (porphyrite, tuff, beds and lenses of sandstone, siltstone and limestone with few taxonomically diverse trilobites and brachiopods). As an example: Gorny Altai, Agayra Formation (Anos Section) – Tremadocian.

**2. Siliceous-terrigenous sedimentary type.**

A. Facies of the distant from shore deep shelf (chert, siliceous mudstone, siltstone with taxonomically diverse radiolarians and rare taxonomically diverse conodonts siliceous sponge spicules). As an example: Tekhten' Formation (sections Tachalov and Barany), siliceous-terrigenous sequence (Suetka Section) – Early Hirnantian.

B. Facies of the shelf foreland, near continental slope edge (sandstone, siltstone, siliceous mudstone with few poor graptolites). As an example: Tekhten' Formation (Rudovozy Section) – Early Hirnantian.

### 3. Terrigenous sedimentary type.

A. Avandelta front of the mountain river (varicolored conglomerate and sandstone). As an example: Upper Subformation of the Kamlak Formation (Section Takoshkin) – Late Tremadocian; Bulukhta Formation (Boriskin Log Section) – Early Hirnantian.

B. Riverside facies:

a) location close to cliffy (iron-bound) desiccated river bank (beds and lenses of the well-rounded and well-sorted conglomerate, gravelstone, sandstone). As an example: lower basal bed of the Tuloi Formation (Mingalevsky, Ishpa sections) – Early Floian;

b) location close to relatively plane bank (massive, well-sorted and well-rounded sandstone). As an example: lower basal bed of the Karasa Formation (Tuloi Section) – Middle Darriwilian.

C. Facies distant from the bank (siltstone, mudstone, rarely sandstone with taxonomically diverse graptolites, trilobites and brachiopods). As an example: middle and upper parts of the Tuloi Formation (Stretenka, Tagaza, Tandoshka sections) – Floian; Bugryshikha Formation (Maralikha Section) – Late Darriwilian – Early Sandbian; Khankhara Formation (Ebogon Section) – Late Sandbian – Early Katian; Ilovaty Formation (Ilovaty, Cheremshanka sections) (Salair) – Floian – Early Darriwilian; Karastun Formation (Korovy Section) (Salair) – Middle Darriwilian – Sandbian; Izyrak Formation (Izyrak Section) (Salair) – Floian – Early Darriwilian.

D. Facies of the underwater highs tops (fine-platy and cross-laminating mudstone with traces of maceration and yielded abundant taxonomically diverse graptolites). As an example: Voskresenka Formation (Pichuzhikha Section) – Floian.

E. Facies of the underwater highs bottoms (gravitation-mixtite) (coarse-laminated sandstone and siltstones with the middle size isolated sphere jointing in terrigenous rocks (“twisting”), up to 0.2–0.3 m in diameter, with rare taxonomically monotonous graptolites). As an example: Voskresenka Formation (Maralikha Section) – Lower Darriwilian.

### 4. Carbonate-terrigenous sedimentary type.

A. Facies slightly distant from the shore (carbonate mudstone, marl, clayey limestone with rare taxonomically poor corals and brachiopods). As an example: Diskovaya Formation (Algain Section) – Early Katian (Gornaya Shoriya).

B. Facies distant from the shore (intercalation of prevailed carbonate mudstone and siltstone, with the secondary in amount middle-laminated clayey limestone yielded few taxonomically diverse brachiopods and trilobites, rare taxonomically poor corals and graptolites). As an example: Khankhara Formation (Verkhnyaya Karasu, Nizhnyaya Karasu, Marcheta-4 sections) – Early Katian.

### 5. Terrigenous-carbonate sedimentary type.

A. Facies distant from the shore (intercalation of prevailed fine-laminated carbonate clayey limestone with the secondary in amount of carbonate mudstone and siltstone yielded abundant taxonomically diverse brachiopods and trilobites, rare taxonomically monotonous corals). As an example: Khankhara Formation (Marinikha, Kholmogorikha sections) – Early Katian.

B. Facies adjacent to distant parts of the reefs (middle and fine-laminated clayey limestone with intercalates of carbonaceous mudstone with abundant diverse corals, trilobites, brachiopods). As an example: Tekhten' Formation (Elanda Section) – Late Katian.

C. Facies of bays on the reef edges (lenses of the algae bioherms small in size, to 1–2 m in diameter, in the siltstone-mudstone matrix with rare taxonomically monotonous brachiopods and graptolites). As an example: Tekhten' Formation (Burovlyanka Section) – Early Hirnantian.

### 6. Carbonate (reef) sedimentary type

A. Facies of the groups of large reefs (up to 2–3 km in diameter) on the carbonate platform at shelf edge:

a) the central parts of separate reefs (massive, un-laminated limestone, with large algae bioherms up to 20–30 m, with rare taxonomically monotonous corals). As an example: Tekhten' Formation (Orlov, Burovlyanka, Marcheta-4, Tekhten', Chakyr, Muta, Bely Bom sections) – Late Katian – Early Hirnantian;

b) the marginal part of the separate reefs (coarse- and middle laminated limestone, with large taxonomically diverse corals). As an example: Tekhten' Formation (Tekhten', Muta sections) – Late Katian;

B. Facies of middle-sized reefs (0.5–1 km in diameter) on the slopes of fade volcanoes (massive, un-laminated limestone, with abundant taxonomically diverse trilobites, as well as rare taxonomically poor brachiopods and conodonts). As an example: Tolstochikha Formation (Orlinaya Section) – Tremadocian (Salair).

C. Facies of separate, small (0.05–0.1 km in diameter), isolated reefs (patch-reef). As an example: Tekhten' Formation (sections Tachalov, Burovlyanka) – Early Hirnantian; Veber Formation (Spornaya Sopka Section) – Early Hirnantian (Salair).

D. Shallow-water distant from shore facies (middle-laminated oolitic limestone, with rare brachiopod and trilobite fragments). As an example: basal bed of the Khankhara Formation (Malaya Uskuchevka, Ebogon, Belaya sections) – Late Sandbian.

### 3.2. SEDIMENTARY MODEL PROFILES FOR THE ORDOVICIAN OF THE ALTAI BASIN

Regularities in lithofacies succession in the different facies area of the Gorny Altai was found out during alignment of the Ordovician local lithostratigraphic subdivisions based on graptolite and conodont zonation (Figs 22–24).

Analysis of fauna paleo-assemblages obtained from the local stratigraphic subdivisions (formations) turned out variations in composition and population density within different fauna groups (Fig. 25).




Fig. 22. Correlation of the Ordovician key sections of the Uymen'-Lebed' facies zone.






Fig. 23. Correlation of the Ordovician key sections of the Charysh-Inya facies zone.

ISC, 2008/ GSC, 2012	Gorny Altai		Revised graptolite zonal successions																																																																																																																																																																																																																																																																																																																																																																										
	Stage	Series	Formation	"Lebed"	"Pridorozhny"	"Tula"	"Yurok"																																																																																																																																																																																																																																																																																																																																																																						
Ordovician	Middle	Dapingian	Kubyshevo	Tula (=Lebed)	gibberulus angustifolius elongatus / broggeri proto-bifidus	gibberulus maximo-divergens deflexus	gibberulus maximo-divergens deflexus	gibberulus																																																																																																																																																																																																																																																																																																																																																																					
									Fragment of graptolite zonal succession (Sannikov et al., 2014, 2015a)	teretiusculus	No graptolites documented	No graptolites documented	No graptolites documented																																																																																																																																																																																																																																																																																																																																																																
														teretiusculus	No graptolites documented	No graptolites documented																																																																																																																																																																																																																																																																																																																																																													
																	jakovlevi / coelatus	No graptolites documented	No graptolites documented																																																																																																																																																																																																																																																																																																																																																										
																				balhaschensis / kirgisicus	No graptolites documented	No graptolites documented																																																																																																																																																																																																																																																																																																																																																							
	Koshtanovskiy	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo				Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo	Kubyshevo

Sections	Lebed'	Bura	Gyr'yanovka Meadow	Tuloi-Karasa	Chechenek	Biya
Distribution of lithological types of rocks						
Sandstone (including gravelstone and conglomerate)	~20%	~3%	15%	~40%	47%	~40%
Mudstone and siltstone	~30%	~9%	30%	~40%	43%	~42%
Limestone (including clayey and sandy)	~50%	~88%	55%	~20%	10%	~18%
Frequency of faunal groups occurrence in separate layers and beds						
High (> 100 specimen)	Brachiopods, tabulate corals /heliolithids	Tabulate corals /heliolithids, brachiopods		Brachiopods	Brachiopods	Brachiopods
Middle (100-10 specimen)	Ostracods, nautiloids	Ostracods, trilobites	Brachiopods, ostracods, tabulate corals	Ostracods		Conodonts, ostracods
Low (< 10 specimen)	Conodonts, trilobites	Conodonts, rugose corals, stromatoporoids		Trilobites, nautiloids		
Rare findings	Rugose corals		Rugose corals, bryozoans, trilobites, ichnofossils	Tabulate corals	Trilobites, crinoids	Corals, bryozoans, trilobites, nautiloids
Taxonomic variety of faunal groups entirely throughout the section						
High (> 10 specimen)		Brachiopods		Brachiopods, ostracods		
Middle (10-5 specimen)	Brachiopods, conodonts	Tabulate corals/heliolithids, trilobites	Brachiopods	Trilobites		Brachiopods
Low (< 5 specimen) and weakly studied	Tabulate/heliolithids, ostracods, trilobites, rugose corals, nautiloids	Bryozoans, rugose corals, conodonts, stromatoporoids	Rugose corals, bryozoans, tabulate corals, ostracods, ichnofossils	Tabulate corals, nautiloids	Brachiopods, trilobites, crinoids	Conodonts, trilobites, bryozoans, corals, ostracods, nautiloids

**Fig. 25.** Correlation of identified zonal successions in the studied sections of the Uymen'-Lebed' structure-facies zone, and regional graptolite zonation for the Lower (excluding Tremadocian) and Middle Ordovician of the Gorny Altai.



In recent years, the event stratigraphy has been increasingly corroborated by numerous data on the world ocean level fall (during regressions) and rise (during transgressions) in absolute ages (Nielsen, 2003, 2004, 2011; Munnecke et al., 2010; Gradstein et al., 2012). The granulometric composition of terrigenous sediments, their sedimentation behavior and fauna characteristics enable first-approximation estimates of absolute paleodepths of the sedimentation (Fig. 26). A comparative study of abrupt changes in the lithological composition within a series of Ordovician sections of the Gornyy Altai allowed marking some of the abrupt large-scale changes in the Ordovician Altai basin depths (Fig. 27).


**Fig. 26.** Standard profile for depths of marine basin.

**Fig. 27.** Reconstructed depth variations within the studied areas of the Gorny Altai paleobasin in the Early–Middle Ordovician.


Integrated analysis of sedimentary features and composition of fauna groups (paleo-communities) revealed the patterns of their relationships based on : a) the distance from the provenance area of terrigenous material supplied to the paleobasin; b) the paleobasin's depth; c) the paleo-seafloor topography (micro-lows, large slope angles, etc.), d) sediment composition, e) grain-size of clastic sediments, f) thickness of forming beds (Fig. 28). The patterns variability is most prominent for the succeeding stratigraphic intervals.

Reconstruction of spatial facies substitution of lithological types of Ordovician sedimentary rocks in the Gorny Altai area throughout the sections revealed different patterns of facies relationships between clastic and carbonate rock associations and, accordingly, changes of fauna groups in paleocommunities (Figs 29, 30).


While analysis and generalization of the revealed patterns allowed to construct the so-called model profiles of Ordovician sedimentation in the paleobasins of the Gorny Altai (Figs 31–33). Each of these profiles has its own prototype (see above Figs 29, 30), represented by a group of real sections of a certain chronostratigraphical interval (straddling usually 1–2, rarely 3 stages) for specific areas corresponding to any offshore part of the Ordovician Altai paleobasin. Comparison of the model profiles allows to reveal specific evolution of individual parts of the paleo-offshore areas and to identify the trends in sedimentation processes within the paleobasin, which serves as the basis for a comparative analysis of the Ordovician Altai basin development and evolutionary stages of other known Ordovician




**Fig. 28.** Model profile of the Altai paleobasin (Uymen'-Lebed' facies zone / Tuloi and Karasa formations / Dapingian Stage – Darriwilian Stage).




**Fig. 29.** Lithological profile of the Altai paleobasin (Anui-Chuya facies zone / Khankhara Formation / Upper part Sandbian Stage – Lower part Katian Stage).



**Fig. 30.** Lithological profile of the Altai paleobasin (Anui-Chuya facies zone / Tekhten' Formation / upper part Katian Stage – Hirnantian Stage).




**Fig. 31.** Model profile of the Altai paleobasin (Charysh-Inya facies zone / Voskresenka Formation / upper part Floian Stage – lower part Dariivilian Stage).




**Fig. 32.** Model profile of the Altai paleobasin (Anui-Chuya facies zone / Khankhara Formation / upper part Sandbian Stage – lower part Katian Stage).





**Fig. 33.** Model profile of the Altai paleobasin (Anui-Chuya facies zone / Tekhten' Formation / Upper part Katian Stage – Hirnantian Stage).



**Fig. 34.** General structural scheme of the Late Ordovician Altai basin.

basins in Siberia, the Urals, Taimyr, the Baltic region and across the world.

Generalization of the series of facies profiles across the outcrops of Ordovician shelf sediments in the Gorny Altai area allowed an inference about the predominant sedimentation types in the Late Ordovician basin described as: (i) reefogenic type in the carbonate-platform zone and (ii) terrigenous-carbonate and terrigenous type in the inner-shelf zone (Yolkin et al., 1994; Sennikov et al., 2008) (Fig. 34).

### 3.3. PALEOGEOGRAPHIC RECONSTRUCTION OF THE ORDOVICIAN ALTAI BASIN


This study is the systematic synthesis of successive changes in paleogeography and contours of the Ordovician Altai basin. The synthesis is based on a series of stage- (and substage) scale paleogeographic charts for the whole Ordovician-Silurian basin history which have been compiled and partly published by today (Yolkin et al., 1994; Sennikov, 2006a, b; Sennikov et al., 2008).

#### 4. ORDOVICIAN KEY SECTIONS IN THE GORNY ALTAI

The described sections are grouped geographically into areas of the western, northwestern, central, northern and northeastern Gorny Altai. The member subdivision and bed-by-bed description of the sections have been the responsibility of N.V. Sennikov, Z.E. Petrunina, L.A. Gladkikh, A.V. Krivchikov, O.T. Obut, E.V. Bukolova (Lykova). The section symbols are keyed as follows. The letters stand for author's name (e.g., Yo for Yolkin, S for Sennikov, P for Puzyrev, R for Russkikh, B for Bukolova in Yo-7039, S-78115, P-78032-1/12, R-7812 and B-097) or for location (LSS for location from succession of sections, F for faunal location, H for shallow hole). Two first numerals after hyphen denote the year when the description was made (70 for 1970, 78 for 1978, etc.) and the following digits before hyphen are section or locality numbers (39, 115, 032); numerals after second hyphen are member numbers (-1 for first member), and numerals after slash mark the sampling depth in meters above the member base (/12 means that the sample was collected at 12 m above the base of the member).

#### 4.1. NORTH-EASTERN GORNY ALTAI (Uymen'–Lebed' facies Zone)

Ordovician sedimentary sequences cropped out in the north-east of the Gorny Altai form a large-scale anticline, accompanied by synclines and tectonic dislocations and belong to the Uymen'-Lebed' facies zone (Fig. 35).



**Fig. 35.** Sketch map of the lower and middle stream of the Biya River (Sennikov et al., 2008).

#### 4.1.1. AREA OF VERKH-BIYSK VILLAGE

##### Pereval Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Tremadocian.

**Regional stratigraphic subdivisions:** Takoshkin Regional stage (Horizon).

**Local lithostratigraphic subdivisions:** Ishpa Formation.

**Zones:** *tenellus* – *kiaeri* graptolite zones.

**Fauna:** trilobites, graptolites.

A small fragment of the Ishpa Formation (S-0726) is exposed in a quarry on a pass along the left side of the roadway from Gorno-Altai to Verkh-Biysk Village (distance mark 105/9) (Figs 36, 37).



Fig. 36. General view of the Pereval Section (upper part of Ishpa Formation).

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Graptolites	Trilobites
Ordovician	Lower	Tremadocian	Ishpa	<i>tenellus</i> - <i>kiaeri</i>	3	>25	Sandstone: greenish-gray, fine to medium, well sorted, and tuffaceous sandstone, with scarce thin (to 10 cm) layers of greenish-gray clayey siltstone; sandstone shows 1.5 m cleavage and 20-30 cm flagginess; thin (1-3 cm) layers of fine sandstone are of cross bedding; fine sandstone encloses lens-shaped (10-15 cm long and 0.5-1 cm thick) medium to coarse sandstone and scarce floating siltstone pebble of 3-10 cm in diameter; there are large (to 15 cm) "tongues" of slumping soft sediment.	<ul style="list-style-type: none"> <li>Adelograptus aff. <i>tenellus</i> (Linnarsson)</li> <li>Paradelograptus sp.</li> <li>Schizograptus sp.</li> <li>Anisograptus sp.</li> </ul>	<ul style="list-style-type: none"> <li>Symphysurus sp.</li> <li>Geragnostus sp.</li> <li>Raphiophoridae</li> </ul>
					2	60	Dark olive-gray siltstone.		
					1	15	Sandstone, dark olive, rather quartz, fine to coarse, polymictic, alternating with dark olive clayey siltstone, often with conchoidal cleavage; both sandstone and siltstone layers are 1 m thick; sandstone is of high and medium roundness and shows good 1-2 cm sorting from fine to medium and on to coarse grain sizes; locally there are syndepositional lenses (from 1-3 cm to 5-7 cm long and 1-3 cm thick) and patches of siltstone in sandstone.		

Fig. 37. Lithology and ranges of fossil taxa from the Pereval Section.



Dark olive-gray siltstone in bedrock exposures along the right side of the roadway between the two quarries (S-0726 and S-0727) contains graptolites (loc. S-0726) *Dictyonema* sp., *Kiaerograptus kiaeri* (Monsen), (loc. S-78146) *Adelograptus* aff. *tenellus* (Linnarsson), *Kiaerograptus kiaeri* (Monsen), and sandstone (loc 205) contains trilobites *Symphysurus* sp., *Geragnostus* sp., Raphiophoridae; graptolites *Anisograptus* sp. are found at loc. 4062a/1668 in the same area; (loc. B-097-099) graptolites *Adelograptus* aff. *tenellus* (Linnarsson), *Adelograptus* sp., *Paradelograptus* sp., *Kiaerograptus kiaeri* (Monsen), *Schizograptus* sp. According to rock lithology and faunas (trilobite and graptolite assemblages), the section fragment belongs to the Ishpa Formation. Graptolites correspond to the Tremadocian *Adelograptus tenellus* and *Kiaerograptus kiaeri* zones.

The total thickness of the Ishpa composite section (S-0726, S-0727, S-78146) may reach no less than 100 m.

#### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

The siltstone topping the studied section comprise layers with traces of washouts formed in the extremely shallow intertidal zone (Fig. 38). Alternatively, rare graptolites were found in the lower and middle parts of the Pereval Section, which may indicate average depths (50–100 m) of the enclosing rocks deposition.



**Fig. 38.** Lithology of the Pereval Section (upper part of the Ishpa Formation).

#### **Ishpa Section**

##### ***Chronostratigraphic subdivisions of the International Stratigraphic Scale:***

Cambrian (Furongian), Ordovician (Tremadocian).

***Regional stratigraphic subdivisions:*** Takoshkin Regional stage (Horizon).

***Local lithostratigraphic subdivisions:*** Ishpa Formation.


***Fauna:*** trilobites, brachiopods, algae, siliceous sponges.

Lower parts of the Ishpa Formation cropped out on the left bank of Biya River, near Village Danilkino: conglomerate greenish-gray polymictic, more than 200 m thick. In the carbonate pebbles trilobites known from the uppermost Cambrian were found.

The two sections accepted as composite stratotype of the Ishpa Formation are located: (1) 350 m upstream of the river's mouth on the left bank of the Ishpa Rv. (Figs 39, 40); (2) one km upstream of the former, on the right bank of the Ishpa river. The latter section reveals the upper horizons of the Ishpa Formation with the concordantly overlying Tuloi Formation (Floian, Dapingian), with conglomerate and siltstone. The total thickness of the Ishpa Formation including the 200 m member of conglomerate in the vicinity of Danilkino Village measures 1085 m. The left-bank part of the stratotype defined from trilobites is dated as Late Cambrian, while the right-bank part is dated as Tremadocian.

#### ***Facies and depositional settings.***

In the studied section, one of the conglomerate members of the Ishpa



**Fig. 39.** Sketch map of the Ishpa area.






















Ordovician					Lithology	Trilobites	Brachio- pods	Algae	Siliceous sponge spicules													
System	Series	Stage	Formation	Member No.						Thickness, m												
Cambrian	Furongian	Stage 10 (Batyrbain)	Tremadocian	Ishpa	13	>80		Siltstone: greenish-gray, snuffy-gray, intercalated with limestone: gray, sandy.														
					12	75		Sandstone: greenish-gray and gray-lilac polymictic, medium-grained, calcareous. Rare siltstone.														
					11	25		Sandstone: gray, pink-gray, arkosic, calcareous.														
					10	75		Sandstone: from fine to coarse-grained, calcareous, and gravelstone. Rock color gray, rare gray-lilac.														
					9	20		Large -pebble conglomerate. In pebbles: sedimentary, volcanic and intrusive rocks. Rock color is greenish-gray.														
					8	5		Limestone: gray, massive, algal-organic, with layers of sandy limestone and siltstone.														
					7	30		Conglomerate: greenish-gray, large-pebble. In pebbles: sedimentary, volcanic and intrusive rocks. Sandstone polymictic, conglomeratic. Contact with the underlying member - tectonic.														
					6	>60		Limestone: gray, yellow-gray, dark gray, massive, fine-grained. Contact with the underlying member - tectonic.														
					5	20		Silty sandstone and calcareous siltstone with gray sandy limestone nodules. Rock color is pinky- and yellow-gray, greenish-gray.														
					4	10		Conglomerate: brownish-gray, in pebbles: granit, porphyrite, limestone, sandstone, siltstone, quartz.														
					3	>60		Silty sandstone and siltstone: bedded, calcareous, with gray limestone nodules, sometimes sandy. Rock color is yellow- and greenish-gray. Lenses and layers of gray limestone and small-pebble conglomerate.														
					2	400		Sodded interval.														
					1	>10		Conglomerate: greenish-gray, polymictic. In pebbles: medium- , coarse-grained biotitic and leucogranite, plagiogranite, quartz-feldspar rocks, quartz, microgranite, gabbro-syenite, gabbro, plagioclase porphyrite, quartzite, limestone, siltstone.														
											<i>Promegalaspides</i> sp. <i>Dolgeuloma</i> sp. <i>Eocheirurus</i> sp.		<i>Acrocephallina</i> sp.		<i>Emsurina</i> sp. <i>Niobella</i> sp.		<i>Pseudoacrocephalites ishpensis</i> Petrun. <i>Bijaspis krivtchikovi</i> Petrun. <i>Mansiella</i> sp. <i>Parakoldinia</i> (?) sp. <i>Lopeuloma</i> sp.		<i>Proapatokcephalops cf. altaicus</i> Petrun. <i>Pseudagnostus</i> sp. <i>Bilacunaspis</i> (?) sp.		<i>Linguella</i> sp. <i>Obolus</i> (?) sp. Acrotretacea Eoorthidae	

Fig. 40. Lithology and ranges of fossil taxa from the Ishpa Section.



Formation consists of the group of conglomerate-breccia layers with the characteristically almost unrounded fragments of tabular sandstone stratification-oriented layers. The deposition of such rocks is largely influenced by coastal erosion (persistent wave action) affecting a steep cliff composed by horizontally lying, weakly cemented sedimentary rocks, with rapid burial of such buildups in wave-cut notches. In some cases, even sandy cement is absent from large fragments (Fig. 41e, f).



**Fig. 41.** Conglomerate in the Ishpa Section (lower part of the Ishpa Formation).

a and b – mixed unrounded blocks and rounded pebbles; c – packages of the fallen down layers near coastal cliff; d – poorly rounded pebble layers: wave-cut notches; e and f – pebbles almost without matrix: regular wave activity.



#### 4.1.2. AREA OF TUROCHAK VILLAGE (LEBED' RIVER)

##### Lebed' Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Floian, Dapingian, Darriwilian, Sandbian, Katian, Hirnantian.

**Regional stratigraphic subdivisions:** Tuloi, Kuibyshevo, Kostinsky, Bugryshikha, Khankhara, Tekhten' and Lystvyanka regional stages (horizons).

**Local lithostratigraphic subdivisions:** Tuloi, Karasa, Gur'yanovka and Chebor formations.

**Zones:** "approximatus", *densus*, *protobifidus*, *angustifolius elongatus*, *gibberulus*, "hirundo – jakovlevi/coelatus", "teretiusculus" graptolite zones.

**Fauna:** graptolites, brachiopods, trilobites, ostracods, orthoceratids, crinoids, gastropods, bryozoans, tabulates, heliolitids, chitinozoans.

The Lebed' Section – is one of the most well studied Ordovician section in Altai (Figs 42–46).

The Gur'yanovka Formation parastratotype is exposed in the right side of the Lebed' River, upstream of former Stretinka Village, with visible contacts with the underlying Karasa Formation and the overlying Chebor Formation. A large part of the Gur'yanovka Formation in the middle of the Bura stratotype section in the Lebed' River right side near former Gur'yanovka Village is hidden under vegetation and encloses dike layers. The Lebed' and Bura sections, with fossils therein, were extensively studied in previous years (Dzyubo, 1960; Sennikov, 1962; Krivchikov et al., 1976; Severgina, 1978, 1984; Petrunina et al., 1984; Kul'kov and Severgina, 1989; Sennikov et al., 2008, 2018a,b; Melnikova, 2010). In the Lebed' Section, the Gur'yanovka Formation basal member composed of greenish and dark gray medium to coarse (up to gravel) polymictic sandstone lies over greenish-gray siltstone of the Karasa Formation top containing graptolites of the terminal Darriwilian *teretiusculus* Zone, in the right side of the Lebed' River, 300 m upstream of the third river shallow from former Stretinka Village (Fig. 42). The basal member is overlain by organic limestone intercalated with limy siltstone and mudstone.

The Gur'yanovka Formation is overlain by purple clayey mudstone of the Chebor Formation with gradual 1 m thick transition. The boundary between the two formations in this section may follow the base of the purple and lilac mudstone or rather the top of uppermost limestone or limy mudstone. The latter interpretation is consistent with red coloration appearing in the Gur'yanovka Formation limestone at the top of the Gur'yanovka Glade and Biya sections (see below). Therefore, the Gur'yanovka Formation – Chebor Formation boundary may correspond to the member 17 base instead of the member 18 top. The thickness of the Gur'yanovka Formation in the Lebed' Section is within 200 m, though it was previously overestimated to >500 m (Krivchikov et al., 1976), possibly, because some riverside outcrops became hardly accessible and the strike of some members aligns with the river stream direction.

**Peculiarities in facies, faunal assemblages and sedimentary environments** (notably for the Gur'yanovka Formation).

According to previous data (Dzyubo, 1960; Krivchikov et al., 1976; Kul'kov and Severgina, 1989; Sennikov, 1962; Melnikova, 2010), the Gur'yanovka Formation (stratotype) in the Lebed'-Stretinka Section contains the fauna assemblages of *Severginella altaica* (Sev.), *Salopia uxunaica* (Sev.), *Glyptomena subgirvanensis* Sev., *Schizophorella altaica* (Sev.), *Severginella shorica* (Sev.), *Chaulistomella lebanonensis* (Cooper), *Austinella lebediensis* Sev., *Boreadorthis togaensis* Sev. brachiopods; *Ceraurinus* cf. *icarus* (Bill.) trilobites; *Grammolomatella* sp., *Eurychilina*? sp. ostracods; *Nyctopora elandensis* Dz. tabulates; and *Cyrtophyllum* ex gr. *jaconurensis* Dz., *Sibiriolites lebediensis* Dz. heliolitids. We have additionally identified *Chaetetes tchakerensis* Dz. heliolitids (member 6); *Nyctopora denticulate* Sok. et Tes., *Nyct. nicholsoni* (Rad.), *Calapoecia baragashiensis* Dz. tabulates (members 7, 9, and 13); and *Scandodus* sp., *Phragmodus undatus* Br. et M., *Panderodus* cf. *P. gracilis* (Br. et M.), *Aphelognathus* sp., *Belodina compressa* (Br. et M.) and *Drepanoistodus suberectus* (Br. et M.) conodonts (member 13).

Limestone occupies a half of Gur'yanovka Formation in the total Lebed' Section volume. It is commonly dark gray, with low clay contents, and enclose 0.5–1.0 cm pure algal nodules, as well as larger and purer nodules of 1–3 cm.

Clay material often produces nodular textures on eroded surfaces or forms separate 5–7 cm layers. The sediments contain locally numerous round or oval algal nodules, 0.5 to 1.0–1.5 cm in diameter, and abundant (70–80 % of the rock volume) 2–3 cm algal calyptrae with knobby surfaces. There appear large (15–20 cm) and small (3–5 cm) colonies of tabulates turned upside down relative to their lifetime position. Large colonies are coated with clayey limestone, which indicates their transport on the sea bottom. Sandstone occupies about 20 % of the section volume (Fig. 47).

The sand material is well sorted and rounded, like siltstone (20 % of the section). Clay most often occurs as a minor component of limestone or less often as separate layers (no more than 10 % of the section volume).

The Gur'yanovka Formation oryctocenoses in the Lebed' Section comprise abundant brachiopods, tabulates, and heliolitids, less abundant ostracods and nautiloids, few trilobites and conodonts, and sporadic rugoses. The taxonomic diversity is the greatest among brachiopods, medium in conodonts, and low in tabulates, heliolitids,

ostracods and trilobites. Lithological features, such as: clayey limestone with nodular textures, limestone with lumpy beds, nodular mudstone, fine-grained well sorted and rounded sand in limestone, algal nodules and calyptrae, colonies of tabulates turned upside down relative to their lifetime position, including those coated with clayey limestone (Fig. 47), etc., indicate shallow marine deposition environments, within 10 m above fair-weather wave base (FWB) for the most of the Gur'yankovka Formation units and subunits in the Lebed' Section. On the other hand, 1–2 mm lamination in mudstone produced by clay laminae indicates deposition at >10 m below FWB. The sea apparently had a rough bottom topography, with rises and depressions of different magnitudes (the laminated mudstones are indicators of shallow depressions). Some carbonate layers in the section also may have deposited >10 m below FWB, judging by brachiopod shelly banks with preserved intact double-valved shells. Note, however, the presence of algal calyptrae that were produced by limy blue-green algae (Cyanobacteria) which bloom only within the euphotic zone, within depths of 30–80 m (Chuvashov et al., 1987; Luchinina, 1973).

In the Lebed' Section, the Chebor Formation is represented by variegated, mostly red sandstone, siltstone, and mudstone. The rocks exhibit different-scale ripple structures, traces of multidirectional ripples, cross-bedding, traces of storm activity (tempestites), desiccation cracks on the mudstone bed surface filled with sandy material of the overlying layer, traces of bioturbation, chaotic (mainly quartz) unrounded to poorly rounded fragments of pebble and gravel particle size in a poorly sorted sandy matrix (Figs 48, 49).





Fig. 42. Sketch map of the Lebed' River area.

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	
Ordovician	Middle	Darrivilian	Karasa	"teretiusculus"	28	22	Alternating siltstone greenish-gray and yellowish-gray and clayey mudstones.	
					27	52	Silty sandstone, rare siltstone calcareous, greenish-gray.	
					26	23	Siltstone greenish-gray.	
					25	102	Silty sandstone, rare sandstone polymictic, calcareous, fine-grained, green-gray, green.	
				"hirundo-jakovlevi" / "coelatus"	24	18	Silty sandstone thin-bedded, gray with lilac color.	
					23	130	Silty sandstone bedded, greenish-gray, green.	
					22	100	Sodded interval.	
					21	>25	Alternating siltstone clayey, yellowish-gray and sandstone polymictic, fine-grained, gray-yellow.	
					20	~20	Siltstone clayey, blue-green.	
				Dapingian	19	<10	Siltstone clayey, bluish-gray.	
					18	25	Sandstone quartz, fine and medium grained, yellowish-gray, massive.	
					gibberulus	17	260	Siltstone, mudstone and silty sandstone calcareous-clayey, thin-bedded, greenish-yellowish-gray, partly dark gray. At the top of the member - rare layers of sandstone polymictic, fine-grained, 3-5 m thick. Rocks are to different grade calcareous, often thin-laminated due to grading by size and color.
						16	335	Alternating mudstone clayey, high foliated and calcareous-clayey siltstone with subordinate sandstone layers, solid, fine-grained, polymictic. Rock color changes from greenish-gray to dark-gray, almost black.
					Lower	Floian	Tuloi	angustifolius elongatus
	14	190	Siltstone clayey, thin-bedded, dark-gray and siltstone calcareous-clayey, greenish-gray, with rare layers of silty sandstone.					
	13	205	Alternating shale silty-clayey, highly cleavaged, siltstone and silty sandstone slightly calcareous, thin-bedded. At the bottom of the member - subordinate layers (up to 1,5 m) of sandstone fine-grained, polymictic. Rock color changes from greenish-gray at the bottom to dark gray at the top of the member.					
	12	95	Sandstone medium-coarse-grained, slightly calcareous, polymictic, medium and thin-bedded, with layers of conglomerate with medium-size pebbles: granite, siltstone, sandstone and quartz.					
	11	140	Sandstone medium-grained, slightly calcareous, medium-bedded. At the bottom of the member - rare conglomerate layers with small-pebbles, gray and greenish-gray, in pebbles: siltstone, sandstone and quartz.					
	10	50	Siltstone clayey, dark-gray, almost balck.					
	densus	9	10	Siltstone clayey, light-gray, snuffy.				
		8	38	Siltstone clayey, black.				
		7	30	Sandstone polymictic, fine-medium-grained, gray yellow.				
		6	15	Siltstone clayey, dark gray, almost balck.				
		5	20	Alternating siltstone clayey and silty sandstone gray-yellow.				
		4	100	Mudstone clayey, high cleavaged, dark-gray to black.				
	"aporoxi-matus"	3	25	Siltstone clayey, gray, iron-gray to black.				
		2	32	Siltstone calcareous-clayey, thin-bedded, with rare layers of sandstone fine-grained, slightly calcareous, gray and yellowish-gray.				
		1	~35	Sandstone fine-grained, quartz, yellowish-gray and gray, with rare layers of quartz gravelstone of same color (several cm thick). Rare quartz pebbles up to 0,5 cm in diameter occur from time to time.				

Fig. 43. Lithology and ranges of fossil taxa from the Lebed' Section (Tuloi and Karasa formations).







**Fig. 44.** General view of the Lebed' Section (lower part, Tuloi Formation).




**Fig. 45.** General view of the Lebed' Section (middle part, Karasa Formation).










**Fig. 47.** Lithological peculiarities of limestone in the Gu'yanovka Formation sections.

The Gur'yanovka Formation – Lebed' Section (b, d, f-i), the Bura Section (a, e, j, k), the Biya Section (c): a – river terrace outcrop ( $2 \times 2$  m) that exposes a sequence of medium-bedded nodular clayey limestone; b – upside down and clay-coated isolated colonies of tabulate corals; c – filled erosion channels in submarine sand bars; d – algal calyptrae in lifetime position; e – nodular mudstone (upper layer on the left) upon clayey limestone surface with wave ripple marks; f – whole brachiopod shells in a coquina bank; g – outcrop ( $3 \times 5$  m) of knobby-lumpy-laminated clayey limestone; h – separated dorsal shells of brachiopods, convex upward, on bed surface; i – lumpy clayey limestone; j – low-angle crossbeds of sandy limestone; k – algal nodules (rolls).





**Fig. 48.** General view of the Lebed' Section (upper part).



**Fig. 49.** Lithological peculiarities of sandstone and siltstone in the Chebor Formation – Lebed’ Section.

a and g – ditches and washouts in the near-shore zone; b and f – tempestites: storm activity; c – sigmoidal micro-layers: tidal markers; d – traces of intensive bioturbation; e – doubled layers: tidal traces.



### Pridorozhny Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Floian, Dapingian, Darriwilian.

**Regional stratigraphic subdivisions:** Tuloi and Kuibyshevo regional stages (horizons).

**Local lithostratigraphic subdivisions:** Tuloi and Karasa formations.

**Zones:** *gibberulus* (*deflexus* and *maximo-divergens* subzones), *hirundo* (*caduceus imitatus* Subzone) graptolite zones.

**Fauna:** graptolites, brachiopods.

The Pridorozhny Section is composed of the Tuloi and Karasa formations (Figs 50, 51).

### *Peculiarities in facies, faunal assemblages and sedimentary environments.*

The mentioned above specific aspects of graptolite paleocommunities and indicating taxa allow the paleobasin reconstruction as follows. Since the Floian–Dapingian–Darriwilian graptolites are documented in more than 100 localities in the studied sections, the embedding strata within the Uymen’–Lebed’ structure-facies zone (northeastern Gorny Altai) were proposedly accumulated in relatively distal shelf environments of the paleobasin. An average amount of the graptolite colonies within each the locality counts 5–10, rarely 20–30 rhabdosomes. The occurrences containing hundreds of the colonies are extremely rare. The graptolite assemblages are taxonomically dominated by species and subspecies of *Isograptus* Moberg, *Pseudisograptus* Beavis, *Parisograptus* Chen et Zhang, *Corymbograptus* Obut et Sob., and *Acrograptus* Tzaj genera, which are also specific for distal parts of paleobasins.

The studied graptolite assemblages contain specific depth indicating taxa. The shallow-water and medium-depth bioindicators are represented by numerous *Hustedograptus teretiusculus* (Hus.) (>40 specimens from 7 localities within the Lebed’ and Yurok sections), *Pseudotrigonograptus ensiformis* (Hall) (~20 specimens from 5 localities within the Lebed’, Pridorozhny and Tuloi sections), species of *Pseudophyllograptus* Cooper et Fortey (>10 specimens from 3 localities within the Lebed’, Tuloi, and Yurok sections), species of *Phyllograptus* Hall (>30 specimens from more than 10 localities within the Lebed’ and Yurok sections), and numerous didymograptins species of *Corymbograptus* Obut et Sob. and *Expansograptus* (Boucek et Pribyl) (>200 specimens from several dozen localities within the Lebed’, Pridorozhny and Tuloi sections).

Besides that, the Lebed’, Pridorozhny and Tuloi sections reveal numerous bioindicators of deep-water environments – species of *Pseudisograptus* Beavis, *Parisograptus* Chen et Zhang, and *Isograptus* Moberg. Rhabdosoms of *Pseudisograptus* Beavis are abundant in the Pridorozhny Section, whereas the Tuloi Section is characterized only by scarce occurrences. Various species of *Parisograptus* Chen et Zhang and *Isograptus* Moberg demonstrate considerable population densities in the Lebed’, Pridorozhny and Tuloi sections. There are only occasional occurrences of deep-water bioindicators (*Pseudisograptus* Beavis and *Parisograptus* Chen et Zhang) documented in the Yurok section.




Fig. 50. General view of the Pridorozhny Section.

Ordovician										System
Middle										Series
Dapingian										Stage
Tuloi										Formation
gibberulus				hirundo		Karasa		Zone		
deflexus		maximo-divergens		caduceus imitatus				Subzone		
								Member No.		
								Thickness, m		
Lithology										Graptolites
				</						

Fig. 51. Lithology and ranges of fossil taxa from the Pridorozhny Section.



All these parameters provide a basis for the bioindicator estimate of the studied sections of the Altai Ordovician paeobasin. Accordingly the dominating graptolite bioindicators, the Pridorozhny Section is reconstructed within the deepened outer shelf zone (>200 m), the Tuloi and Lebed' sections correspond the lower outer shelf environments (150–200 m), whereas the Yurok Section is situated in the upper outer shelf zone (100–150 m) of the paleobasin.

Upstream from the Lebed' Section on the right bank of Lebed' River near mouth of its right tributary Bura River, in the vicinity of the former Gur'yanovka Village, the upper part of the Gur'yanovka Formation and stratotype of the Cheborka Formation are situated (Bura and Gur'yanovka Glade sections) (Fig. 52).




Fig. 52. Sketch map of the Gur'yanovka area.





Fig. 53. General view of the Bura Section (lower part).

### Bura Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Sandbian.

**Regional stratigraphic subdivisions:** Khankhara and Tekhten' regional stages (horizons).

**Local lithostratigraphic subdivisions:** Gur'yanovka Formation.

**Fauna:** conodonts, trilobites, ostracods, brachiopods, stramatoporoidea, tabulates, heliolitids, rugoses, bryozoans.

The Bura Section crops out in the right side of the Lebed' River, upstream of the Bura mouth (Figs 52–54). It was first documented by V.M. Sennikov (1962) and then by other researchers (Krivchikov et al., 1976; Kul'kov and Severgina, 1989). The river has changed its course for the past fifty years, the formerly exposed terraces became overgrown with trees and shrubs, and most of the Gur'yanovka Formation stratotype section became almost inaccessible. According to our observations upstream of the Bura mouth, limy gray mudstone lies over the 5 m thick top member of the Karasa Formation composed of dirty yellow or snuffy fine- to medium-grained massive sandstone with low lime contents and under limestone layers with different clay contents intercalated with scarce sandstone and mudstone layers and sets of layers. The total currently exposed thickness of the lower Gur'yanovka Formation in the Bura Section is at least 170 m, which is much less than 500 m reported by Sennikov (1962), without the 150 m thick vegetated interval. Kul'kov and Severgina (1989) estimated the thickness of the middle Gur'yanovka Formation as 400 m.

**Peculiarities in facies, faunal assemblages and sedimentary environments.**

Earlier studies (Cherepnina, 1960; Dzyubo, 1960; Krivchikov et al., 1976; Kul'kov and Severgina, 1989; Melnikova, 2010; Yaroshinskaya, 1960) revealed the following fauna assemblages: (1) *Onniella* cf. *flava* (Havl.), *Apatamorphia altaica* Sev., *Fascifera buraensis* Sev., *Eoanostrophia lebediensis* Sev., *Eridorthis subinexpecta* Sev. brachiopods and *Homotelus* sp., *Eorobergia* sp., *Bumastus* sp. trilobites in the lower part of Gur'yanovka Section; (2) *Eridorthis subinexpecta* Sev., *Catazyga salairica* (Sev.), *Boreadorthis togaensis* Sev., *Strophomena lebediensis* Sev. in Rozm., *Dactylogonia subgeniculata* Sev., *Chaulistomella amzassensis* (Sev.), *Severginella altaica* (Sev.), *Parastrophinella salairica* Sev. brachiopods; *Ceraurinus icarus* (Bill.), *Iliaenus* sp. trilobites; *Stallipora vesiculosa* Modz., *Phenopora multifera* Nekh. bryozoans; *Grewinkia altaica* Tcherepn. rugoses; *Eofletcheria mironovae* Dz., *Nyctopora elandensis* Dz. tabulates; and *Cyrtophyllum baragashensis* Dz., *Sibiriolites lebediensis* Dz. heliolitids in the middle part of section part; (3) *Glyptorthis praerulchra* Sev., *Hesperorthis lebediensis* Sev., *Austinella lebediensis* Sev., *Salopina uxunaica* (Sev.), *Strophomena lebediensis* Sev., *Glyptomena subgirvanensis* Sev., *Fardenia* cf. *scalena* Will., *Parastrophinella salairica* Sev., *Anoptambonites grayae sibirica* Sev., *Catazyga salairica* (Sev.), *Severginella schorica* (Sev.), *Triplexia ainca* Sev., *Rhynchotretoidea aincus* Sev., *Spirigerina sublevis* Rozm.





brachiopods in the upper section part, along with *Egorovella demissa* Melnik ostracods; *Nictopora minimalis* (Rad.), *Nic. tschakerenensis* Dz., *Calapoecia baragashensis* Dz., *Cal. anticostensis* (Bill.), *Paleofavosites inkensis* Tscherep. tabulates; *Sibiriolites lebediensis* Dz. heliolitids; *Clathrodictyon kirgizicum amzassensis* V. Khalf. stromatoporoids; *Constellaria floridaformis* Jarosh. bryozoans; *Anclotichia commutatis* Jarosh., *Stellopora vesiculosa* Modz. We have found also *Scandodus* sp. and *Panderodus* sp. conodonts in member 10 of the Bura Section.

Most of the Bura Section (88 %) is occupied by limestone of massive algal, detrital, nodular clayey (clay nodules), gravelly carbonate (calcirudite), and oolitic varieties and brachiopod coquina, with frequent algal calyptrae and sporadic graded bedding (according to the sizes of benthic clasts). Sandstone in the Bura Section makes 3 % of the section volume. It is well rounded and sorted.

Siltstone and mudstone constitute 9 % of the total thickness and are often poorly consolidated, with signatures of transport and rolling by storms. Low-angle crossbeds delineate small depressions of the sea bottom.

The Gur'yanovka Formation oryctocenoses in the Bura Section comprise predominant brachiopods, tabulates, and heliolitids, less frequent ostracods and trilobites, and few conodonts, stromatoporoids and rugoses. The taxonomic diversity is the greatest in brachiopods, intermediate in trilobites, tabulates and heliolitids, and low in bryozoans, rugoses, and conodonts. Lithology features, such as: 1) limestone with knobby lumpy beds; 2) nodular clayey limestone; 3) twisted lumpy mudstone layers; 4) carbonate gravelstone (calcirudites); 5) algal nodules and calyptrae; 6) sorting into layers according to sizes of benthic fauna; and 7) brachiopod coquina, suggest shallow-water deposition (3–5 to 10 m above fair-weather wave base) of most units and subunits in the Bura Section. The presence of calyptrae and hermatypic corals records deposition in the euphotic zone, to depths of 30–80 m. It is possible to make up a composite section (of three segments) of the Gur'yanovka Formation/Chebor Formation transition in the immediate vicinity of the Bura Section, near former Gur'yanovka Village; for the Chebor Formation this corresponds to the lower one third of its stratotype. The lower part of the Gur'yanovka Glade Section correlates with the second half of the upper Gur'yanovka Formation stratotype (Krivchikov et al., 1976; Kul'kov and Severgina, 1989; Sennikov, 1962) and continues the Bura Section, judging by the strike of rocks (Krivchikov et al., 1976; Sennikov, 1962). At present, the section part corresponding to the Gur'yanovka Formation top is hidden under thick vegetation while that correlated with the overlying Chebor Formation became better exposed as a result of the Lebed' River course change. Thus, we observed (Figs 55, 56) two terminal units of the composite Gur'yanovka Formation stratotype: (i) intercalated gray mudstone, siltstone and fine sandstone with sporadic lenses of reddish limestone and (ii) intercalated gray and greenish-gray clayey limestone and limy sandstone.

### **Gur'yanovka Glade Section**

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Katian, Hirnantian.

**Regional stratigraphic subdivisions:** Khankhara and Tekhten' regional stages (horizons).

**Local lithostratigraphic subdivisions:** Gur'yanovka Formation.

**Fauna:** trilobites, ostracods, tabulate and rugose corals, bryozoans, ichnofossils.



**Fig. 55.** General view of the Gur'yanovka Glade Section (middle part).



System	Series	Stage	Formation	Member No.	Thickness, m	Lithology	Brachiopods	Trilobites	Ostracods	Tabulate corals	Rugose corals	Bryozoans	Ichnofossils
Ordovician			Upper Gur'yanovka Subformation										
Ordovician			Upper Katian										
Ordovician			Upper Chebor										
Ordovician			Himantian										

Fig. 56. Lithology and ranges of fossil taxa from the Gur'yanovka Glade Section.

#### 4.1.3. AREA OF TULOI VILLAGE

##### Tuloi Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Floian, Dapingian, Darriwilian, Sandbian.

**Regional stratigraphic subdivisions:** Tuloi (Lebed'), Kuibyshevo, Kostinsky and Bugryshikha regional stages (horizons).

**Local lithostratigraphic subdivisions:** Tuloi, Karasa and Gur'yanovka formations.

**Zones:** *aproximatus*, *densus*, *angustifolius elongatus*, *gibberulus*, *hirundo*, *dentatus* graptolite zones; *Cyathochitina parvicolla*, *Cyath. calix* chitinozoan zones.

**Fauna:** graptolites, brachiopods, trilobites, ostracods, orthoceratids, crinoids, gastropods, bryozoans, hyolithes, chitinozoans.

The stratotype section of the Tuloi Formation extends along the right side of the Biya River downstream of the Tuloi inflow and then upstream the Tuloi on its right side (Figs 57–62). According to graptolites, trilobites, and brachiopods, the Tuloi Formation spans the Floian, Dapingian, and lower Darriwilian. The formation overlaid conformably over the Ishpa Formation and is underlain by the Karasa Formation.





Fig. 57. Sketch map of the Tuloi Village area.



Fig. 58. Upper part of the Tuloi Formation in the Tuloi Section.





**Fig. 59.** Upper part of the Tuloi Formation in the Tuloi Section.

Members 1 through 19 belong to the Tuloi Formation, members 20 through 23 belong to the Karasa Formation, and members 24 and 25 belong to the Gur'yanovka Formation. The total thickness of the Tuloi Formation in the section exceeds 2500 m, the Karasa Formation is 450 m thick, and the Gur'yanovka Formation is more than 150 m thick.

The graptolite zonation of the section is as follows: members 1 through 5 belong to the *aproximatus* Zone (loc. LSS-401a, S-75141); member 6 may conventionally correspond to the *densus* and *angustifolius elongatus* zones (loc. H-81), and members 7 and 8 are conventionally aligned with the *gibberulus* Zone (loc. 282, H-79); the base of member 9 correlates with the *gibberulus* – *hirundo* boundary, and members 9 through 16 belong to the *hirundo* Zone (loc. H-78, H-75, H-73, LSS-422a, LSS-423, H-91, H-84, H-85, H-87, LSS-418, H-88); members 17 through 22 contain no graptolites; members 18 through 22 correlate with the Darriwilian according to trilobites and brachiopods (loc. H-95, LSS-415, H-92, H-102, H-103); graptolites from member 23 (loc. H-104, LSS-409) correspond to the *dentatus* Zone.



System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Chitinozoans
Ordovician	Upper	Sandbian	Gur'yankova		25	56	Siltstone, greenish-gray, calcareous, with silty fine sandstone and sandy limestone.	<i>Conochitina oelandica</i> Eisenack <i>Conochitina simplex</i> Eisenack <i>Conochitina bacillum</i> Obut et Zaslavskaya <i>Cyatochitina calyx</i> (Eisenack) <i>Cyatochitina tuloyensis</i> Obut et Zaslavskaya <i>Desmochitina minor</i> coeca Eisenack <i>Rhabdochitina regula</i> Obut et Zaslavskaya  <i>Desmochitina minor typica</i> Eisenack <i>Hercochitina</i> sp. <i>Rhabdochitina</i> sp. <i>Desmochitina minor amphorea</i> Eisenack <i>Desmochitina cf. minor elongata</i> Eisenack <i>Lagenochitina aff. deunffi</i> Paris <i>Conochitina infraspinoza</i> Wilson et Dolly <i>Conochitina</i> sp. <i>Conochitina parvicolla</i> Tougeourdeau
					24	90	Conglomerate: greenish-gray or purple-gray, fine, with pebbles of granite, sandstone, siltstone, siliceous rocks, quartz, etc., conglomerated sandstone, and medium to coarse sandstone.	
	Middle	Darrivillian	Karasa	dentatus	23	125	Siltstone: greenish-gray, more rarely silty sandstone; a 10-15 cm thick siltstone layer in member upper part abounds in diverse well preserved fossils; the rock can be classified as trilobite-brachiopod coquina.	
					22	95	Silty sandstone: greenish-gray, less often fine polymictic sandstone and siltstone.	
					21	150	Sandstone: gray, greenish-gray to green, polymictic, fine, occasionally siltstone and silty sandstone, locally limy sandstone.	
				hirundo	20	80	Sandstone: gray or light gray, quartz, medium to coarse.	
					19	165	Interbedded greenish-gray siltstone, mudstone, and silty sandstone.	
					18	10	Sandstone: greenish-gray and brownish-gray polymictic, rarely siltstone.	
					17	120	Interbedded greenish-gray silty sandstone and siltstone, with scarce layers of fine sandstone.	
					16	40	Interbedded greenish-gray fine sandstone and silty sandstone, with scarce thin (to 1 cm) layers of gray and dark gray siltstone.	
					15	90	Silty sandstone: greenish-gray, less often greenish-gray fine polymictic sandstone, and dark gray siltstone.	
					14	50	Silty sandstone: greenish-gray, less often greenish-gray fine polymictic sandstone, and dark gray siltstone.	
					13	45	Silty sandstone: greenish-gray, more rarely fine polymictic sandstone, and greenish-gray, gray to dark gray siltstone.	
					12	55	Interbedded fine polymictic sandstone and silty sandstone, greenish-gray and gray, with scarce thin (to 2 cm) layers of dark gray calcareous siltstone.	
					11	65	Interbedded siltstone, silty sandstone, and rarely fine sandstone, greenish-gray, occasionally with thin layers of gray to dark gray siltstone.	
					10	230	Sandstone: greenish-gray or dark olive-gray, fine, polymictic, alternating with silty sandstone, or less often siltstone.	
					9	20	Siltstone: gray or dark gray, occasionally silty sandstone.	
					8	155	Siltstone: greenish-gray, less often silty sandstone (to fine polymictic sandstone).	
					7	150	Sandstone: greenish-gray, polymictic, fine to medium.	
					6	230	Interbedded greenish-gray siltstone and silty sandstone, less often fine sandstone.	
					5	700	Interbedded siltstone, silty sandstone, and fine polymictic sandstone, greenish-gray, dark olive-gray, locally dark gray.	
	Lower	Floian	Tuloi	angustifolius elongatus	4	175	Sandstone: greenish-gray, polymictic, fine to medium.	
					3	150	Siltstone: greenish-gray, more rarely gray to dark gray, rarely purple.	
					2	70	Sandstone: variegated, polymictic, fine to medium.	
					1	130	Conglomerate, variegated, coarse, or less often conglomerated sandstone, with pebbles of granite, microgranite, porphyry granite, pegmatite granite, intermediate and basic porphyry, tuff, quartz, siltstone, sandstone, clayey and siliceous shale.	
				densus				
				approximatus				

Fig. 60. Lithology and ranges of fossil taxa from the Tuloi Section.





Member No.	Thickness, m	Lithology	Graptolites	Other groups
16c	>8	Interbedded (at 0.3-0.5 m in lower layers and at 0.1-0.3 m in upper layers), silver-gray well rounded and well sorted fine polymictic sandstone and dark gray clayey siltstone.	<ul style="list-style-type: none"> <li>■ <i>Isograptus gibberulus</i> (Nicholson)</li> <li>■ <i>Isograptus caduceus nanus</i> (Ruedemann)</li> <li>■ <i>Isograptus imitata</i> (Harris)</li> <li>■ <i>Isograptus maximo-divergens</i> (Harris)</li> <li>■ <i>Isograptus</i> sp.</li> <li>■ <i>Pseudoisograptus manubriatus</i> (T.S. Hall)</li> <li>■ <i>Tetragraptus harti</i> (Hall)</li> <li>■ <i>Expansograptus</i> sp.</li> </ul>	<ul style="list-style-type: none"> <li>■ Brachiopods</li> </ul>
16b	1,5	Siltstone: gray.		
16a	0,7	Sandstone: olive, slightly calcareous, fine to medium, well rounded and well sorted.		
15c	10	Thinly (1-3 cm) interbedded siltstone and fine sandstone; sandstone layers are slightly thicker than those of siltstone; sandstone shows cross bedding; the section is cut by a fault apparent in the middle of the quarry.		
15b	5	Mudstone and siltstone, dirty olive, of thin 1-3 cm banding produced by more silty layers in mudstone.		
15a	3	Mudstone and clayey siltstone, dark silver-gray or black.		

**Fig. 61.** Lithology and ranges of fossil taxa from the middle part of the Tuloi Section.



**Fig. 62.** Upper part of the Karasa Formation in the Tuloi Section.



## Yurok Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Dapingian, Darriwilian.

**Regional stratigraphic subdivisions:** Tuloi, Kuibyshevo, Kostinsky and Bugryshikha regional stages (horizons).

**Local lithostratigraphic subdivisions:** Tuloi and Karasa formations.

**Zone:** “*angustifolius elongatus*”, “*gibberulus*”, “*hirundo*”, “*sinodontatus*”, “*austrodentatus*”, “*dentatus*”, “*balhaschensis/kirgiscus*”, “*jakovlevi/coelatus*”, “*teretiusculus*” graptolite zones.

**Fauna:** graptolites, trilobites, brachiopods, gastropods, echinoderms, nautiloides, bryozoans, bivalves, phyllocarids.

The Yurok Section was studied during 2011–2015 (Sennikov et al., 2018a,b). It is localized on a side of the road, connecting Artybash and Turochak villages, 240 m northward from a bridge across the Yurok Creek (right tributary of the Biya River). The section comprises subvertical beds of the Upper Tuloi and Karasa formations, which are stratigraphically arranged towards the Teletskoe Lake, with a total thickness along the road exceeding 50 m. The section is paleontologically characterized by graptolites, trilobites, phyllocarids, gastropods, brachiopods, ostracods, bryozoans, bivalves, echinoderms, and nautiloids. The Yurok Section is logged as follows (Figs 63–65):

### *Peculiarities in facies, faunal assemblages and sedimentary environments.*

In the Yurok Section, the upper Tuloi Formation (20-m thick interval corresponding the Upper Floian and Lower–Middle Dapingian *angustifolius elongatus*, *gibberulus* and *hirundo* graptolite zones) and lower Karasa Formation (up to 45 m of the Darriwilian *austrodentatus* – *teretiusculus* graptolite zones) are composed of monotonous clayey mudstone with scarce interbeds of siltstone and yellow-brownish-gray (snuffy-gray) fine-grained sandstone. The basal quartz sandstone member of the Karasa Formation is not represented in the Yurok Section. It could be caused by a hidden tectonic fault, since there is a low-amplitude fault zone documented within the lower Karasa Formation, between the first and second members of the section. Specific lithological composition (very scarce sandstone interbeds, dramatically reduced thickness) of the Upper Floian – Lower–Middle Dapingian upper Tuloi Formation interpret the sedimentary record comprised in the Pridorozhny and Yurok sections as the distal parts of the shelf paleobasin. Black and dark color of the rocks and lack of the benthic fauna in the Pridorozhny Section substantiate an interpretation of this section as the most distal one. However, the light (snuffy) color of the strata, as well as both abundant and diverse fossils of the benthic fauna (trilobites, ostracods, phyllocarids, gastropods, brachiopods, bryozoans, echinoderms) in the Yurok Section contradict its interpretation within the deepened outer shelf, or even within the comparatively deep-water environments. At the same time, there are numerous specimens of full



Fig. 63. General view of the Yurok Section.

System	Series	Stage	Formation	Zone, subzone, "zonal level"	Member No.	Thickness, m	Lithology	Graptolites	Trilobites	Other groups																																																																																									
Ordovician																																																																																																			
Middle																																																																																																			
Darriwilian																																																																																																			
Karasa																																																																																																			
Dapingian	Tuloi	1	"gibberulus"	"hirundo" → "sino-dentatus"	2	6,5	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Unidentifiable graptolites Pseudophyllograptus angustifolius angustifolius (Hall) Callograptus sp. Pterograptus sp. (? "Didymograptus" sp.) "Undulograptus" sinodontatus (Mu et Lee) Undulograptus sp. "Glyptograptus" (s.l.) sp. "Glyptograptus" eosicatus Tzai Phyllograptus sp. Phyllograptus ilicifolius Hall Phyllograptus anna longus Rued. Phyllograptus anna Hall "Pseudoclimacograptus" sp. Hustedograptus teretiusculus (Hisinger)	Raymondaspis sp. Lonchodomas rostratus (Sars) Robergia sparsa Nikol. Cybelurus altaicus Levit. Paracybeloides (?) sp. Unidentifiable trilobites	Phyllocarids Gastropods Brachiopods Ostracods Bryozoans Nautiloidea																																																																																									
											3	3	4	8,5	5	> 30	Mudstones: yellow-brownish-gray ("tobacco-gray"), massive, fissile clayey. Individual convoluted pillows, 40-50 cm in diameter, are documented in the lower part of the interval.	Megistaspis (M.) polyphemus Brogger Amplexograptus confertus (Lapw.) Amplexograptus sp. "Glyptograptus" euglyphus (Lapw.) Glossograptus acanthus Elles et Wood	Carolinites sp.	Echinoderms																																																																															
																					Alternating siltstones: yellow-brownish-gray, clayey mudstones: brown-yellowish-gray and sandstones; light yellowish-brown, fine-grained, well-sorted. The siltstone beds are 30-50 cm-thick, thickness of the mudstones varies at 10-100 cm, and the sandstones beds are at 5-10 cm.	Fault	Siltstones: dark yellow-brownish-gray ("tobacco-gray") clayey, with columnar fissility.	Clayey mudstones: yellow-brownish-gray ("tobacco-gray") with columnar fissility. Pillows of convoluted lamination, 40-50 cm in diameter, are scarcely developed throughout the interval.																																																																											
																									"balhaschensis" / "jakovlevi" / "kirgisicus" / "coelatus"	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco"), clayey.	Mudstones: yellow-brownish-gray ("tobacco-gray") and grayish-yellowish-brown ("grayish-tobacco

Fig. 64. Lithology and ranges of fossil taxa from the Yurok Section.





**Fig. 65.** Subaqueous slumping “twistings and rolls” in mudstone found in the Yurok Section (Karasa Formation).

trilobite skeletons found in the Yurok Section, which preservation conditions postulate a lack of wave influence on the benthic communities. Therefore, the upper Floian and lower-to-middle Dapingian strata of the Yurok Section must be accumulated not only below the fair-weather wave base, but even below the base of storm wave agitation – within the upper outer shelf and medium-depth inner shelf (50–100–150 m). Considering both remoteness of the Yurok Section from the shoreline and relatively shallow depth, we interpret the upper Floian and lower–middle Dapingian strata within the section to be accumulated on a top or slope of a local submarine uplift. This assumption is further supported by the gravity slump structures (convoluted pillows) found in the Yurok Section.

Amongst the trilobite taxa from the Yurok Section above enlisted, *Cybelurus altaicus* Levit. has the biggest potential for appliance as an age-constraint and correlative marker. This taxon was originally defined on a material obtained from the Bugryshikha and Gora Altai sections (stratotypes of Bugryshikha Formation and Bugryshikha Horizon), localized near the Bugryshikha Village in the northwestern Gorny Altai (Charysh–Inya facies zone). The type material originates from the stratigraphic level, characterized by the upper Darriwilian graptolites of *teretiusculus* Zone (Levitsky, 1963; Sennikov et al., 2008). Within the Loktevka–Batun structure-facies zone of the Gorny Altai, this trilobite species was found in the Batun Section (named after the Batun Village, nearby located), within the Voskresenka Formation which comprises there a stratotype of the Kostinsky Horizon (Levitsky, 1962, 1963; Perfil'ev and Levitsky, 1963; Sennikov, 1977; Sennikov et al., 1982, 2014, 2015a). Trilobites *Cybelurus altaicus* Levit. are found there in association with middle–upper Darriwilian graptolites of *dentatus* and *balhaschensis* zones. Composite trilobite complex of the Bugryshikha and Batun areas, associated with *Cybelurus altaicus* Levit., comprise *Cybelurus planus* Levit., *Cybelurus batunensis* Levit., *Aksanatella altaica* Petrun., *Aksanatella bugryshichica* Petrun., *Eorobergia ojrotica* Levit., *Raymondaspis communis* Levit., *Ceraurinus abnormalis* Levit., *Ceraurina* cf. *frequens* Tschug., *Bijacybele* cf. *strigosa* Petrun., *Lonchodomas communis* Levit., *Bathyurellus nonnulus* Tschug., *Pliomera fischeri asiatica* Tschug., *Pliomerellus amplissimus* Petrun., *Pliomerellus* cf. *jacuticus* Tschug., *Kolymella* aff. *plana* (Tschug.), *Glaphurus altaicus* Weber, *Homotelus* sp., *Carrickia* sp., *Carolinites* sp., *Nileus* sp. Within the northeastern Gorny Altai (Uymen'–Lebed' structure-facies zone), trilobites *Cybelurus altaicus* Levit. were identified in the Tuloi Section, in the Karasa Formation (Andreeva, 1985). This stratigraphic level was lately constrained as the upper Darriwilian *dentatus* graptolite Zone. *Cybelurus altaicus* Levit. is also known from Tuva Region. These trilobites were revealed in the stratotype section of the Tarlyk Formation of the Malinovka Group, localized near the Tarlyk Village. Trilobite complex, including *Cybelurus altaicus* Levit., was found in the lower Tarlyk Member, which





Fig. 66. General view of the Biya Section.

section represents a stratotype of the Lower Tarlyk Horizon (Andreeva, 1985; Levitsky, 1962; Sennikov et al., 2006a, b, 2015a), in association with the Darriwilian graptolites and lower Darriwilian conodonts (*Eoplacognathus variabilis* – *E. suecicus* Zone) (Sennikov et al., 2000, 2015a). This trilobite complex contains *Cybelurus altaicus* Levit., *Cybelurus planifrons* (Weber), *Carolinites spinosus* And., *Carolinites marophthalma* (Harr. et Leanz), *Carolinites* aff. *genacinaca* Ross, *Bulbaspis* cf. *ovulum* (Weber), *Plesiomegalaspis* aff. *estonica* Tjernv., *Plesiomegalaspis* sp., *Cybele* cf. *bellatula* Dalm., *Symphysurus* cf. *exactus* Tschug., *Symphysurus* cf. *kujandensis* Tschug., *Ampyx* aff. *politus* Raymond, *Ampixella clavata* And., *Ogigites* aff. *almatyensis* Tschug., *Robergia deckeri* Coop., *Lonchodomas eximius* And., *Malinaspis tuvaensis* And., etc. Until recently, the upper member of the Tuloi Section has been characterized by trilobite taxa, identified by Z.E. Petrunina (Sennikov et al., 2008), and along with new species including *Encrinuroides* sp., *Ceraurinella* sp., *Atractopyge* sp., *Pliocybele* sp., *Raymondaspis* sp., *Robergia* sp., *Robergiella* (?) sp., *Remopleurella* sp., *Sphaerexochus* sp., *Carolinites* sp., *Hemiarges* sp., *Trinodus* sp., *Calyptaulax* sp., *Stegnoipsis* (?) sp., *Iliaenus* sp., *Dimeropyge* (?) sp., *Otarion* sp., *Lonchodomas* sp., *Ampyx* sp., *Nilieus* sp. Our recent discoveries demonstrate the trilobite complex of the upper member of the Karasa Formation within the Tuloi Section sits inside the middle Darriwilian *dentatus* graptolite Zone, and contains certain trilobites taxa, not registered within the Yurok Section. The Tuloi trilobite complex is comprised by *Megistaspis* (*M.*) *polyphemus* Brogger, *Lonchodomas rostratus* (Sars), *Robergia sparsa* Nikol., *Ampyx* sp., *Paracybeloides* (?) sp., *Carolinites* sp., *Raymondaspis* sp., *Agerina* sp., *Niellus* sp. In Baltoscandia, trilobites *Megistapis* Jaanusson gave a name for “*Megistapis* limestones” (type area Bornholm Island), localized in the upper Volkhov Horizon and allegedly accumulated on a relatively distal shelf (Bergström et al., 2013; Pärnaste and Bergström, 2013), adjacent to deep-water facies of graptolite mud (Männil, 1966).

### Biya Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** upper Sandbian, Katian.

**Regional stratigraphic subdivisions:** Khankhara and Tekhten’ regional stages (horizons).

**Local lithostratigraphic subdivisions:** Gur’yanovka and Chebor formations.

**Fauna:** crinoids, trilobites, ostracods, bryozoans, nautiloids, corals, conodonts.

The Biya section, which was mentioned in geological survey reports and in some early publications (Kul’kov and Severgina, 1989; Sennikov et al., 1959; etc.) and documented in detail by our team for the first time, is located in the right side of the Biya River upstream of the Chechenek Creek mouth. The base of the Gur’yanovka Formation does not crop out in this section (see above). River terrace outcrops expose alternating sandstone, mudstone, and clayey limestone



System	Series	Stage	Formation	Member No.	Thickness, m	Lithology	Brachiopods	Crinoids	Ctenoids	Trilobites	Brachiopods	Nautiloids	Corals	Conodonts
Ordovician	Upper	Katian	Chebor Formation	24	4.8	Moderately sorted rounded gray fine to medium polymictic rather quartz sandstone, with poorly sorted subrounded and subangular floating gravel (3–4% of total rock volume); up to 2 cm cracks (possibly produced by desiccation) filled with sand; medium to coarse sandstone in member upper part.								
				23	2.9	Gray or light gray massive clayey and sandy limestone, with algal rolls (1–2 cm in diameter); clay material often forms random spots.								
				22	1.4	Well sorted and rounded gray rather quartz fine to medium polymictic sandstone, with signatures of submarine sand bar erosion by currents (up to 20 cm outcut).								
				21	1.7	Lilac-red or gray siltstone and sandstone similar to rocks of member 19; limy siltstone in member middle; coarser material (to fine sandstone) in upper part.								
				20	1.2	Lumpy limestone with honeycomb and nodular textures; algal rolls (up to 1 cm in diameter).								
				19	6	Siltstone and sandstone similar to those in member 17; isolated 20 cm beds of well sorted rounded lilac-red massive fine sandstone with low lime contents; mostly yellowish-dirty gray rocks in upper part.								
				18	0.6	Gray massive limestone, reddish-lilac in upper part; detrital organic limestone in upper part.								
				17	12	Thin-bedded (1–5 cm) siltstone, purple with gray spots, with low lime contents; 15 cm thick bed of gray limestone at member base; 5 cm thick interbed of red limestone with brachiopods at 2 m above member base; 3–5 cm interbeds of gray massive siltstone and fine sandstone with high lime contents upsection; gray siltstone grading into fine sandstone in member upper part.								
				16	1.7	Beds of olive green or greenish-gray thin-platy limy siltstone alternating with thick beds (3 to 5 m) of purple siltstone, grading to one another along strike.								
				15	12	Well sorted rounded light greenish-gray flaggy massive fine polymictic sandstone; 40 cm thick beds in member lower part and 5–10 cm beds in member middle and top; thin-bedded sandstone grading into purple clayey siltstone in upper one third.								
				14	10	Thin-bedded (3–5 cm) platy light snuffy-gray slightly lumpy clayey siltstone with low lime contents.								
				13	7	Alternating lilac and greenish-gray fine to medium sandstone; 25–30 cm flaggy bedding in 0.5–1.0 m thick lilac beds; 15 cm thick bed of gray-pale reddish limestone at 2 m above member base; medium bedding (10–15 cm) with perfectly smooth plane surfaces at 3 m above member base.								
				12	3	Sandstone similar to massive sandstone of member 9.								
				11	7	Moderately sorted rounded lilac and pale lilac fine to medium sandstone with lime cement; lighter-colored greenish rocks in member upper part.								
				10	10	Sandstone similar to sandstone of member 9.								
				9	4	Poorly sorted subrounded dirty green-gray fine to medium sandstone, with moderately sorted subrounded floating coarse sand.								
				8	6	Thin-bedded (5–10 cm) flaggy sandstone similar to that in member 4; interbeds with lime cement.								
				7	12	Gray and dark gray massive limestone with low clay contents; clay material forms laminae and lenses in member lower part; such laminae and lenses are fewer in member middle but are quite frequent in upper part.								
				6	4	Gray and dark gray massive limestone with low clay contents.								
				5	4	Snuffy-gray clayey mudstone.								
				4	0	Well sorted rounded snuffy-dirty gray fine to medium polymictic sandstone.								
				3	9	Dark gray massive limestone alternating with nodular (3–5 cm) limestone and snuffy-gray clayey mudstone; clayey limestone, massive limestone, and mudstone occupy, respectively, gray massive 70–75%, 10%, and 20% of member thickness; bed thicknesses: 0.2–0.3 m for mudstone and up to 0.5 m for limestone.								
				2	5	Dark gray-snuffish nodular (2–5 cm) clayey limestone; clay material often forms laminae and 15–30 cm long lenses; gray massive limestone with lower clay content in upper part.								
				1	15	Moderately sorted and rounded snuffy fine polymictic sandstone; 15–20 cm or less often 3–5 cm flaggy bedding; lime cement in some interbeds (5–15 cm).								
Ordovician	Lower	Sambian	Gur'yankovka Subformation											

Fig. 67. Lithology and ranges of fossil taxa from the Biya Section.

(Figs 66, 67). The Gur'yankovka Formation has a total thickness of 170 m in the section (without the buried basal part). Its boundary with the overlying Chebor Formation corresponds to the top of the terminal limestone bed, and the uppermost member (N 24) should belong to the Chebor Formation rather than to the Gur'yankovka Formation. Note that the presence of floating pebbles and gravel among sandstone and siltstone was mentioned earlier (Krivchikov et al., 1976; Sennikov, 1962) in the terminal Biya section member, as well as in the basal members of different Chebor Formation sections. Note also that we observed a fragment of the Chebor Formation composed of alternating lilac and gray sandstone, siltstone, and mudstone 100–150 m south of the Biya section end, in an isolate fault-bounded block on the right side of the Biya River.

### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

The previously reported fauna found in the section on the right side of Biya River, near the Chechenek Creek mouth, and assigned to the Karasa Formation (Kul'kov and Severgina, 1989) was restricted to *Glyptorthis primus* Sev. brachiopods. L. Severgina also found *Strophomena lebediensis* Sev. in Rozm., *Chaustimella amzassensis* (Sev.), *Rostricellula ainsliei amzassica* Sev., *Schizophorella altaica* (Sev.), *Triplesia ainca* Sev., *Boreadorthis togaensis* Sev., and *Eridorthis subinexpecta* Sev. brachiopods typical of the Gur'yanovka Formation fauna during geological surveys, between 510 and 615 m downstream of the Yurok River inlet into the Biya. We discovered *Belodina compressa* (Br. et M.), *Panderodus gracilis* (Br. et M.), *Phragmodus undatus* Br. et M., and *Erraticodon* sp. conodonts in members 3 and 20.

Both clastic and carbonate Gur'yanovka Formation rocks have red and lilac colors in the middle of the Biya Section. The appearance of this bright coloration (purple, brown, lilac, and red) in the northern and eastern parts of the Uymen'-Lebed' facies zone was previously interpreted (Krivchikov et al., 1976; Sennikov, 1962) as the principal or at least significant criterion to separate the Gur'yanovka Formation from the overlying Chebor Formation. Note, however, that lilac rock coloration appears also at the base of the Gur'yanovka Formation stratotype described above, near the Bura mouth, and red rocks occur at the top of the Gur'yanovka Formation in the Gur'yanovka Glade section (see above). Almost all limestones in the Biya section are clayey or less often sandy. Some are detrital organic varieties (coquina), with detritus of various fauna groups, as well as limestone with algal rolls, 1–2 cm in diameter. Siltstone are fine- or medium-grained, rounded, well or moderately sorted, with either clay or lime cement. The cross sections of some sandstone members display signatures of submarine sandbars eroded by currents, with channels incised to 25 cm depths. Such channels were apparently formed below storm wave base (SWB), at sea depths more than 30 m. The presence of sandstone layers with perfectly smooth surfaces in the Biya Section indicates deposition in relatively deep water, below SWB (deeper than 30 m).

The Gur'yanovka Formation fauna assemblages in the Biya Section include predominant brachiopods, less frequent or few ostracods and conodonts, and sporadic trilobites, nautiloids, tabulates, and bryozoans. Brachiopods show the highest taxonomic diversity in the Gur'yanovka Formation oryctocenoses, while conodonts and trilobites are represented by few taxa. Limestones in the section almost lack corals and are free from algal calyptrae, which is evidence of sunlight shortage on the sea floor; together with the lack of clay lithologies, this indicates clear water and medium shelf depths exceeding 30–80 m in the disphotic zone. Facies of other Gur'yanovka Formation rocks in the section suggest deposition environments below fair-weather wave base but within SWB (10–30 m) or greater sea depths (50–100 m) for different section parts.

## **4.2. EASTERN GORNY ALTAI (Teletskoe Lakeside facies zone)**

### **4.2.1. AREA OF IOGACH VILLAGE**

#### **Tozodov Section**

##### ***Chronostratigraphic subdivisions of the International Stratigraphic Scale:***

***Regional stratigraphic subdivisions:*** Kuibyshevo, Kostinsky, Bugryshikha regional stages (horizons).

***Local lithostratigraphic subdivisions:*** Tozodov Body.


***Fauna:*** trilobites, ostracods, brachiopods, orthoceratids, bryozoans.

The Tozodov Section is located on the right side of the eponymous brook (right tributary of the Iogach Rv.) in two road quarries, one km from the creek's mouth (Figs 68–70). The total thickness of the exposed part of the section is more than 150 m. The thickness of the non-red part of the Tozodov Section as a possible lithological analogue of the often green-gray Karasa Formation (upper half of the Stretinka Group), is at least 120 m. This interval should be considered as a stand-alone gray terrigenous Tozodov Body. The thickness of red-colored part of the Tozodov Section, which was previously correlated with the Chebor Formation (Uymen'-Lebed' facies zone), is at least 30 m.


### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

The ichnofossils Gordia Emmons established in the seventh member are interpreted as biomarkers of relatively deep-water facies (Mangano, Droser, 2004). The ostracods *Egorovella* sp. were found in the third member of the Tozodov Section, while the eighth member contained diverse trilobites, namely *Asaphus knyrkoi* F. Schmidt, *Asaphus striatus* Brogger, *Lonchodomas rostratus* (Sars), *Pliomera fischeri* (Eichwald), characteristic of the middle and lower parts of the Upper Ordovician of Baltoscandia (Ivantsov, 2003). The faunas reported from this member also include the bryozoans *Dianulites ramosiformis* Jaroshinskaja.





**Fig. 68.** General view of the Tozodov Section (lower part).



**Fig. 69.** General view of the Tozodov Section (middle and upper parts).

System	Series	Stage	Formation	Member No.	Thickness, m	Lithology		Nautiloids	Trilobites	Brachiopods	Ostracods	Gastropods	Bryozoans	Ichthyofossils
Ordovician	Middle	Darniian	Tozodov											
	Upper	Katian	Logach											
		Sandbian												

Fig. 70. Lithology and ranges of fossil taxa from the Tozodov Section.



### 4.3. NORTHERN GORNY ALTAI (Biya-Katun' facies zone)

#### 4.3.1. AREA OF KAMLAK VILLAGE

##### Kamlak Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Upper Cambrian (Furongian) and Lower Ordovician (Tremadocian).

**Regional stratigraphic subdivisions:** Ordovician – Takoshkin Regional stage (Horizon).


**Local lithostratigraphic subdivisions:** Kamlak Formation.

**Zone:** *Iapetognathus* conodont Zone, *osloensis-hyperboreus* graptolite Zone.

**Fauna:** conodonts, graptolites, trilobites, brachiopods, chitinozoans, problematics.

Tremadocian strata are most completely represented and best constrained by faunas near Kamlak Village thus making a tie for Tremadocian correlations throughout the Gorny Altai. A small (6×7 km) Kamlak graben is filled with terrigenous-carbonate sediments folded in an asymmetric fold cut by faults. According to lithology and fauna patterns, the stratotype section includes three units of unequal thicknesses: The Lower, Middle, and Upper Kamlak subformations (Figs 71–73). The subdivision of the Kamlak Section into subformations is based on lithology and faunal control, mainly from trilobites. Members are aligned according to lithology and trilobite, graptolite, and brachiopod assemblages.

The Lower Kamlak Subformation crops out in the middle reaches of the Maly Kamlak River where the base of the Kamlak Formation contacts, along a fault, with the volcanic rocks of the Middle Cambrian Ust'-Sema Formation. The total thickness of the Lower Kamlak Subformation is about 120 m. The Lower and Middle Kamlak subformations are divided by a fault. The total thickness of the Middle Kamlak Subformation is about 440 m. The Middle and Upper Kamlak subformations are



**Fig. 71.** Sketch map of the Kamlak area.

**Fig. 72.** Limestone of the Member 10 of the Middle Kamlak Subformation (Kamlak Section).



System	Series	Stage	Horizon	Formation	Subformation	Zone	Member No.	Thickness, m	Lithology
Ordovician	Lower	Tremadocian	Takoshkin (= Upper Tayanza)	Kamlak	Upper	osloensis-ramosus	14	>30	Sandstone: gray, dark gray, or greenish-gray, outsized, polymictic.
							13	330	Conglomerate: fine and medium, with poorly sorted quartzite, quartz, granite, volcanic, sandstone, and limestone pebbles of high to medium roundness.
							12	70	Siltstone: brown-red, grading toward member top into gray and greenish-gray siltstone, with fine sandstone layers and limestone lenses in member upper part
							11	8	Limestone: black and gray, clayey, with black, gray, and yellowish-gray siltstone interbeds.
							10	130	Siltstone: brown-red, with lenses and layers of red nodular limestone.
							9	165	Siltstone and mudstone, brown-red.
							8	35	Fine and medium conglomerate.
							7	>20	Siltstone: gray or greenish-gray.
							6	>200	Siltstone and mudstone, brown-red, with fine sandstone interbeds and limestone lenses.
							5	115-170	Conglomerate: fine and medium, with poorly sorted quartzite, quartz, granite, volcanic, and limestone pebbles of high or medium roundness, grading on strike into gravelstone and sandstone; conglomerate encloses scarce lenses of reddish-gray sandy limestone.
							4	170	Siltstone and mudstone, brown-red.
							3	29	Sandstone: gray or reddish-gray, fine to coarse, with angular clasts of brown-red siltstone in detrital material.
							2	35	Conglomerate: fine, grading on strike into sandstone.
							1	59	Sandstone: gray or reddish-gray, fine to coarse, with siltstone interbeds.
							10	15	Limestone: yellow and grey, lumpy
							9	250	Limestone: light gray, massive, crystalline, with lenses and layers (1-3 m) of reddish and grayish-brown limestone in lower layers.
							8	31	Siltstone: lilac or brown-red.
							7	8	Conglomerate: fine to medium.
							6	12	Sandstone: variegated, rather quartz-feldspar, outsized, with minor amounts of disseminated "jasperoids".
							5	31	Conglomerate: fine to medium, compositionally similar to that in member 3.
Cambrian	Upper	Batyrbaiian	Lower-Middle Tayanza		Middle	C. caseyi	4	22	Siltstone: lilac-gray, brownish-gray or greenish-gray.
							3	30	Conglomerate: fine to medium, with well rounded but poorly sorted pebbles of diverse quartzite, porphyry, less often schists, gneisses, and limestone in a limy-sandy cement.
							2	34	Limestone: light gray, massive, crystalline, with 2-4 m thick layers and lenses of purple-red limestone at member base.
							1	>20	Siltstone: gray or purple-gray, with 1-3 cm thick silty sandstone interbeds.
							7	>2	Conglomerate: fine, with well- and rather well-rounded pebble of black and red quartzite and granite in a sandy-clayey-limy cement.
							6	60	Siltstone and slightly calcareous mudstone: lilac or brown-red, with layers of gray fine massive rather quartz-feldspar sandstone.
							5	6	Limestone: pinkish-gray, massive, locally thin-bedded.
							4	18	Siltstone: lilac-gray, calcareous, of uncertain bedding.
							3	6-8	Limestone: lilac- or pinkish-gray, of uncertain banding, crinoidal.
							2	17	Siltstone and mudstone: purple-gray and brown-red, with 3-4 cm pebble of dark gray limestone and black quartzite.
		Dobry			Lower	Proconodontus	1	6	Limestone: light gray, locally pinkish- or brownish-gray, thick-bedded.

Fig. 73. Lithology and ranges of fossil taxa from the Kamlak Section.



Member No.	Thickness, m	Graptolites	Conodonts	Trilobites	Brachiopods	Chitinozoans and problematic fossils
14	>30					
13	330					
12	70					
11	8					
10	130					
9	165					
8	35					
7	>20					
6	>200					
5	115-170					
4	170					
3	29					
2	35					
1	59					
10	15					
9	250					
8	31					
7	8					
6	12					
5	31					
4	22					
3	30					
2	34					
1	>20					
7	>2					
6	60					
5	6					
4	18					
3	6-8					
2	17					
1	6					

Fig. 73. The end.

divided by a fault. The Upper Kamlak Subformation has a total thickness of about 1400 m and is of a Late Tremadocian age according to the faunas.

According to B.D. Erdtmann who looked through the data, some graptolite forms found in member 12 upper Kamlak Subformation and identified as *Triograptus osloensis* Monsen and *Aletograptus hyperboreus* Obut et Sob. may belong to genus *Psigraptus* Jackson. These forms are rhabdosomes buried in sandstone, and their preservation leaves unknown the intravital direction (upward or horizontal?) of branches off the sicula. On the other hand, the Altai morphs lack occluded autothecae typical of *Psigraptus* and thus may be transitional from the *Triograptus* Monsen and *Aletograptus* genera to *Psigraptus* Jackson. The graptolite assemblage from locs S-7662, S-7664, and S-7661 generally corresponds to the *osloensis* - *ramosus* Zone.

In the year 2007 P. Mannik collected limestone samples from the Kamlak Section. Chemical proceeding revealed conodonts (provisional identification): from the lower part of the Middle Kamlak Formation, member 9, sample N 10 and from the upper part of the Middle Kamlak Formation, member 16, sample N 9 – *Variobiliconus* sp.

The composite section of the Kamlak Formation can be correlated with the International Stratigraphic Chart on the basis of few species of conodonts and graptolites. The conodont species *Oneotodus datsonensis* Druce et Jones found in members 9 and 10 in the lower half of the Middle Kamlak Subformation is known in the Datsonian and lowermost Warendian groups of Australia that span a stratigraphic interval of the *Cordylodus proavus*, *Hirsutodontus simplex*, *Cordylodus prolindstromi* and *Cordylodus lindstromi* zones (Dubinina, 2000). One may expect to find a transitional assemblage between the *Cordylodus lindstromi* and *lapetognathus fluctivagus* zones in the Lower Middle Kamlak Subformation. The graptolite assemblage found in member 12 of the Upper Kamlak Subformation corresponds to the *osloensis*-*ramosus* Zone. Earlier data on trilobites from the Lower Kamlak Subformation defined its Early Tremadocian age (Ermikov et al., 1979; Petrunina et al., 1984). The present view is that the Lower Kamlak Subformation must rather correlate with the uppermost Cambrian, proceeding from the revised conodont-based age and stratigraphy of the Ordovician-Silurian boundary strata (Webby et al., 2004) and from the correlation of the Mansian and Loparian regional stages of the Siberian Platform with the Late Cambrian (Kanygin et al., 2007). The trilobite and graptolite data suggest a Late Tremadocian age of the Upper Kamlak Subformation. Thus, the Middle Kamlak Subformation may correlate with the Lower Tremadocian. The trilobite assemblage from the Middle Kamlak Subformation differs from those in both the Lower and Upper Kamlak subformations.

#### 4.4. NORTH-WESTERN GORNY ALTAI (Anui-Chuya facies zone)

##### 4.4.1. AREA OF UST'-KAN VILLAGE

###### Chakyr-Azratkan (Chakyr-Elanda) Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Katian.

**Regional stratigraphic subdivisions:** Tekhten' Regional stage (Horizone).

**Local lithostratigraphic subdivisions:** Tekhten' Formation.

**Fauna:** alge, tabulate corals, stromatoporoids, trilobites, brachiopods, graptolites.

One of the studied sections crops out on the watershed ridge between the Chakyr and Elanda rivers 4.8 km (azimuth 185) from the Elanda mountain (elevation: 1663.7 m). The massive gray limestone with rare poorly preserved corals are overlain by terrigenous strata (Figs 74, 75). Two uppermost carbonate members in this section can be attributed to typical outcrops of the Chakyr beds with characteristically abundant corals.

The tabulate corals established on the Chakyr-Azratkan watershed in the upper member of platy limestone (being part of the stratotype section of the local stratigraphic unit) include: *Nyctopora altaica* Dz., *N. tchakyrensis* Dz., *Lyopora altaica* Dz., *Eofletcheria kovalevskiyi* Dz., *Calapoecia baragashiensis* Dz. and *C. lebediensis* Dz.

The brachiopod assemblages *Onniella* cf. *chancharica* Severg., *Sowerbyella* sp. were encountered along with corals in carbonates, while graptolites *Pseudoclimacograptus* sp., *Lincograptus* sp. and *Desmograptus* sp., trilobites *Sceptaspis* cf. *unica* Petrun., *Lonchodomas* sp., *Homotelus* sp., *Ceraurinus* sp., *Eorobergia* cf. *crassilimbata* Petrun., and the brachiopods *Dactylogonia subgeniculata* Severg., *Rostricellula* sp. are reported from the under- and overlying calcareous-terrigenous rocks.


The youngest carbonate member (thickness: 80 m) is featured as a solitary ridge descending into the Azratkan valley, where at the foot of its left side limestone become massive and bear many algal structures forming separate calyptrae and their grouping both in intravital and displaced positions (Sennikov et al., 2001; Sennikov et al., 2017).

The fauna found in the upper two members of the laminated limestone which constitute the second integral part of the stratotype of the Chakyr Beds section along the strike of the right bank of Azratkan (Elanda) Section (Fig. 76) included tabulate corals *Calapoecia lebediensis* Dz., *C. altaica* Dz., *C. anticostensis* Bill., *Eofletcheria kovalevskiyi* Dz., *Billingsaria* sp., *Fletcheriella altaica* Dz., *Nyctopora asratkanensis* Dz., *N. altaica* Dz., *Nyctopora* sp., *Catenipora elandensis* Dz. and *Eofletcheria sokolovi* Dz., heliolitids *Chaetetes tchakyrensis* Dz. (Sennikov et al., 2001). In addition to these, formerly from this composite section on the Chakyr Rv. (Dzyubo, 1960a,b, 1966; Dzyubo, Mironova, 1960; Cherepnina, 1960; Stratigraphic..., 1975) reported the tabulate *Nyctopora minimalis* (Radugin),



**Fig. 74.** Sketch map of the Chakyr-Azratkan area.

*Grewingia semilunatum* (Scheffen), *Catenipora kuruensis* Sok., *Paliphyllum primarium* Soshk. and rugose corals *Brachielasma altaica* Tcherepn. The listed tabulate genera from the Chakyr beds included (besides the above mentioned) those identified earlier *Saffordophyllum* Bass., *Foerstephyllum* Bass., *Reuschia* Kiaer, *Fletcheriella* Sok., and also heliolitids – *Wormsipora* Sok., *Stelliporella* Went., *Proheliolites* Kiaer (Dzyubo, 1966). In the past, this taxonomic diversity and representativeness in corals in the section once served as the basis for identification of the Chakyr Formation as a new local straton (Chakyr limestone, Chakyr beds) (Stratigraphic..., 1975; V. Sennikov, Sennikov, 1982; Sennikov et al., 2001, 2017).



System	Series	Stage	Formation	Member No	Thickness, m	Lithology	Trilobites	Brachio-pods	Tabulate corals	Algae
Ordovician	Upper	Katian	Tekhten'	16	>30	Alternating limestone bedded (0.3-0.5 m), grey and siltstone yellow, clayey-calcareous (1-5 cm) with algae and crinoids.				
				15	3-5	Limestone: grey-yellow, clayey.				
				14	50	Limestone; black, clayey, bedded (0.2-0.3 m), fine-grained, with tabulate corals, brachiopods, crinoids.				
				13	25	Limestone: gray, massive, fine-grained, pelitomorphic.				
				12	60	Sodded interval.				
				11	65	Siltstone: black, clayey with intercalates of sandstone and limestone. In the middle part of the member 2-5 cm layers of calcareous sandstone brownish, thin-flaggy, and sandy limestone.				
				10	30	Sodded interval.				
				9	115	Limestone: light-grey, massive, pelitomorphic, with intercalates of crinoid limestone, gray.				
				8	10	Sodded interval.				
				7	90	Alternating (in lower part of the member) 0.1-1.0 m, 10 m quartz-feldspar or calcareous sandstone: gray, brownish, greenish- and yellow-gray, fine-, medium-grained, and clayey siltstones: gray, black and greenish-gray.				
				6	50	Quartz-feldspar sandstone and silty sandstone: greenish-, snuffy-gray and gray, bedded (2-7 cm), fine- and medium-grained, with rare layers (3-7 cm) of siltstone greenish-gray, clayey. In the lower part of the member - lenses (5x15 cm) of limestone gray, organic.				
				5	20	Repeated intercalation of siltstone: greenish-gray, clayey (0.2 m) with sandstone: gray, fine-grained (3-10 cm), and rare limestone: gray, dark-gray detrital (up to 10 cm).				
				4	180	Sandstone: gray and brownish-gray, fine-grained, calcareous (varying degree) sandstone; thin-flaggy with rare 1-3 cm layers of siltstone dark-gray, clayey.				
				3	160	Alternating siltstone black and dark-gray (0.5 m) and sandstone gray-brownish, fine-grained (5-10 cm). In the upper part of the member number sandstone layers reduce.				
				2	60	Siltstone: gray, greenish-gray, clayey, thin-foliated. In the lower part of the member small "pebbles" of siltstone light-colored. In the middle and upper parts of the member - rare 2-5 cm layers of sandstone gray, fine-grained and slightly calcareous sandstone brownish, fine-grained.				
				1	>5	Quartz-feldspar sandstone: gray, fine-grained, flaggy.				

**Fig. 75.** Lithology and ranges of fossil taxa from the Chakyr-Azratkan (Chakyr-Elanda) Section.

System	Series	Stage	Formation	Member No.	Thickness, m	Lithology	Trilobites	Brachiopods	Graptolites	Tabulate corals and helioliths	Algae	Stromatolites
Ordovician	Upper	Katian	Tekhten'	19	80	Limestone: gray, platy. Laterally changed by massive-bedded.	<ul style="list-style-type: none"> <li>■ <i>Paracybeloides</i> sp.</li> <li>■ <i>Remopleurides</i> sp.</li> <li>■ <i>Calyptaulax</i> sp.</li> </ul>	<ul style="list-style-type: none"> <li>■ <i>Omniella cf. chancharica</i> Sev.</li> <li>■ <i>Sowerbyella</i> sp.</li> </ul>	<ul style="list-style-type: none"> <li>■ <i>Lincograptus</i> sp.</li> <li>■ <i>Desmograptus</i> sp.</li> </ul>	<ul style="list-style-type: none"> <li>■ <i>Nyctopora altaica</i> Dz.</li> <li>■ <i>Catenipora</i> sp.</li> <li>■ <i>Nyctopora altaica</i> Dz.</li> <li>■ <i>Nyctopora ichakysensis</i> Dz.</li> <li>■ <i>Lyopora altaica</i> Dz.</li> <li>■ <i>Eofletcheria kovalevskyi</i> Dz.</li> <li>■ <i>Calapocia baragashensis</i> Dz.</li> <li>■ <i>Calapocia lebediensis</i> Dz.</li> </ul>	<ul style="list-style-type: none"> <li>■ <i>Stromatolite</i></li> <li>■ <i>Cyanobacteria</i></li> <li>■ <i>Hedstroemia</i> sp.</li> </ul>	<ul style="list-style-type: none"> <li>■ <i>Clathrodictyon</i> (?) <i>kirgisicum amzassensis</i> V. Khalif.</li> </ul>
				18	20	Marl: yellow- and cream-brown, bedded						
				17	20	Sandstone: brownish-gray, limy, fine-grained, fine-flaggy.						
				16	80	Alternating 5-15 cm layers of sandstone brown-gray, limy, fine-grained and siltstone calcareous-clayey, fine-foliated.						
				15	60	Siltstone: dark-gray and black, clayey, foliated.						
				14	50	Sodded interval.						
				13	>20	Siltstone: yellow-gray, calcareous-clayey, fine-foliated.						
				12	110	Limestone: gray, algal, massive, sometimes platy (0,1-0,5 m). In the upper part of the member - black limestone.						
				11	15	Sodded interval. Cropping rocks from members 9 and 10.						
				10	10	Sandstone: brownish, quartz-feldspar, fine and medium-grained.						
				9	5	Siltstone and mudstone: black and dark-gray, clayey.						
				8	30	Sandstone: brownish, polymictic, mainly quartzitic, fine- and medium-grained, flaggy.						
				7	40	Sandstone: dark-gray and yellow-gray, polymictic, fine-grained.						
				6	5	Siltstone: greenish-gray and dark-gray.						
				5	40	Sandstone: brownish-gray, polymictic, fine and medium-grained.						
				4	10	Siltstone: gray, limy grading into clayey limestones.						
				3	8	Alternating siltstone dark-gray and sandstone yellow-gray, fine-grained, polymictic.						
				2	15	Sandstone: brownish, fine-grained, quartz-feldspar, flaggy.						
				1	50	Limestone: light-gray, massive.						

Fig. 76. Lithology and ranges of fossil taxa from the Azratkan (Elanda) Section.


#### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

Members in the Chakyr-Azratkan (Chakyr-Elanda) Section with abundant corals should be termed "coral meadows". The organogenic structures found on the left bank of the Azratkan (Elanda) River are composed of calcareous blue-green algae (Cyanobacteria), which, being enveloped (wrapped) in the bacterial film, formed organic micro-structures in the form of algal crusts, irregular hemispheres (1-2 cm), isolated from each other calyptrae (micro-domes) of 3-5, rarely up to 7 cm in diameter and 1-2 cm in height composed successively layered bacterial mats covers (Fig. 77).

Cyanobacteria (calyptraes) together with corals formed coral-algae meadows, covering large areas of the bottom. Localities occupied by extensively developed cyanobacterial micro-structures are characterized by significantly lower corals density, than the areas with sparse calyptraes.

Not only cyanae "build" calyptraes on the basis of their own colonies (Luchinina, 1973), rather their activity co-existing (symbiosis) with bacteria provides the basis for the formation of organomineral structures (stromatolites). In theses structures the "capture" of sediment particles and the biomineralization of the colony mats of calcareous blue-green filamentous algae take place along with the minerals precipitation on the surface of such tissue. Stromatolite





**Fig. 77.** Microbial-algae structures in limestone of the Tekhten' Formation (Chakyr-Azratkan Section).

a - g – calyptrae, h – stromatolite's structure, i – calcareous algae *Hedstroemia* sp.

structures were also recorded in the studied section. Besides cyanobacteria, calcareous red algae (Rhodophyta) devoid of a bacterial membrane were found in the sections. Red algae are a characteristic element of organogenic buildups (Brooke, Riding, 1998; Luchinina, Terleev, 2004). The red algae *Hedstroemia* sp. revealed among the latter was close to *Hedstroemia aequalis* Hoeg were identified in the Kozhim Rv. Section in the Urals (Chuvashov et al., 1993).

### **Ebogon Section**

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Sandbian, Katian.

**Regional stratigraphic subdivisions:** Bugryshikha, Khankhara, Tekhten' regional stages (horizons).

**Local lithostratigraphic subdivisions:** Bugryshikha, Khankhara, Tekhten' formations.

**Zone:** *clingani* graptolite Zone.

**Fauna:** graptolites, trilobites, brachiopods, ostracods, bryozoans, gastropods.

The section on the Ebogon and Elanda watershed was studied by several researchers (Sennikov, Vinkman, Kononov, 1959; Kononov, 1964). The area is bounded by the Ebogon and Elanda watershed and complicated by a series of small ridges.

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology		Graptolites	Brachiopods	Trilobites	Other groups
Ordovician	Upper	Katian	Tekhten	?	24	8		Limestone, platy, gray.	<i>Climacograptus</i> sp., <i>Dicranograptus</i> sp., <i>Diplograptus</i> sp., <i>Glyptograptus</i> sp., <i>Pseudoclimacograptus sharenbergi</i> (Lapworth), <i>Dicellograptus</i> sp.	<i>Orhambonites jaboganicum</i> Severg., <i>Orhambonites</i> sp., <i>Strophomena</i> sp., <i>Apatomorphia</i> sp., <i>Rostrocellula</i> sp., <i>Skenidioides costatus</i> sparsis Severg., <i>Multicostella</i> ( <i>Chaulistomella</i> ) <i>iniquistrata</i> Coop., <i>Isophragma ricevillense</i> Coop., <i>Plectrothis</i> aff. <i>tensis</i> Coop., <i>Leptellina</i> aff. <i>magna</i> (Rukav.), <i>Orhambonites</i> sp.	<i>Homotelus</i> sp., <i>Lonchodomas</i> sp., <i>Calyptraulax jaboganicus</i> Petrun., <i>Bronteops</i> sp., <i>Bronteopsis transversalis</i> Petrun., <i>Carriackia chancharica</i> Petrun., <i>Pseudosphærocochus</i> sp., <i>Helioimeris exotica</i> Petrun., <i>Tornquistia</i> sp., <i>Lonchodomas rostratus</i> (Sars), <i>Glaphurella sibirica</i> Petrun., <i>Sphaeræcochus</i> sp., <i>Cerninus amplius</i> Petrun., <i>Raymondella</i> sp., <i>Remophorella</i> sp., <i>Isotelioides ebogensis</i> Petrun., <i>Remophorella</i> sp., <i>Otarionella</i> cf. <i>koksoriana</i> Korol., <i>Calyptraulax</i> sp.	Crinoids
					23	30		Sodded interval. In crops - rocks from members 19 and 20.				
Ordovician	Sandbian	Khanakhara	Bugryshikhka	?	22	40		Siltstone clayey dark-gray and black.				
					21	80		Sandstone fine-grained, polymictic, mainly quartz, greenish.				
					20	160		Alternating sandstone polymictic, fine-grained, gray, silty sandstone and clayey siltstone. Sandstone sometimes significantly quartz with greenish color, especially in the upper part of the member.				
					19	3		Clayey limestone gray with crinoid stems (up to 50 % of rock).				
					18	125		Sandstone quartz and polymictic, fine-medium-grained, gray. In the upper part of the member appear brownish-gray silty sandstone.				
					17	8		Clayey limestone gray, replaced by calcareous siltstone.				
					16	100		Sandstone polymictic, mainly quartz, fine-grained, greenish-gray. In the upper 1 m layer of the member gray clayey siltstone.				
					15	55		Alternating siltstone yellowish- and greenish-gray and silty sandstone with boudines (3 x 5 cm) of sandstone fine-grained and limestone gray.				
					14	90		Clayey siltstone dark-gray and greenish-gray. In the upper part of the member rock color change to lighter and yellowish.				
					13	50		Alternating siltstone and silty sandstone greenish-gray, with boudines (3 x 3 cm) sandstone fine-grained.				
					12	5		Sandstone fine-grained gray.				
					11	80		Clayey limestone gray, on the strike changed by calcareous siltstone. Algal bioherms (1 x 1.5 m).				
					10	10		Calcareous siltstone dark gray.				
					9	40		Alternating (20-30 cm) sandstone fine-grained and silty sandstone gray.				
					8	8		Calcareous siltstone gray, with pelitomorphic limestone nodules.				
					7	1.5		Limestone oolitic, gray.				
					6	30		Clayey siltstone gray and greenish-gray, with layers of sandstone polymictic, fine-grained, gray.				
					5	5		Calcareous siltstone gray, with limestone nodules (30-50 cm).				
					4	10		Calcareous mudstone yellowish gray.				
					3	160		Clayey siltstone dark gray, in the upper part of the member sandstone fine-grained gray.				
					2	60		Sandstone fine-grained and siltstone dirty yellow-gray.				
					1	65		Sandstone medium grained, mainly quartz, with floating gravel gray.				

Fig. 78. Lithology and ranges of fossil taxa from the Ebogon Section.



The initial point of the section is located on one of these ridges, descending from the spurs (1663.7 m), 4.2 km from the elevation mark (azimuth 270°). A bed-by-bed description provided below concerns with only the topmost part of the section. The thickness of the lower part of the section totaling 525 m thick is characterized by alternation of sandstone and siltstone with rare thin (up to 1.5 m) intercalations of limestone. Given that the faulting affected the saddle on the crest, the upper part of the section is stratigraphically directed upsection and down the crest along the azimuth 250° (Fig. 78).

The section established along the ridge axis. The Katian trilobites were found 100-150 m north of the section line in the localities U-1671 и U-1672, in the deposits corresponding to the lower half of the 7th member. In addition to the above mentioned taxa these fauna finds included trilobites *Jabogonellus* sp., *Ceraurinella* sp., *Raymondella* sp., *Isoteloides* sp., *Pharostoma* sp., and *Calyptaulax* sp., brachiopods *Chaulistomella inaquistriata* (Coop.), *Isophragma zicevillense* Coop., *Apatomorpha altaica* Severgina, *Rostricellula* sp., *Plectorthis* aff. *tensis* Coop., *Leptellina* aff. *magna* (Rukav.), *Oniella altaica* Severgina, *Altaeorthis uscutchevi* Severgina, *Sivorthis jaboganicum* (Severg.) and Katian graptolites (clingani Subzone, *quadrimucronatus* Zone) *Dicranograptus clingani* Carruthers, *Climacograptus* sp., *Glyptograptus* sp., as well as ostracods, bryozoan, gastropods and crinoids.

#### 4.4.2. AREA OF UST'-MUTA VILLAGE

##### Marchetyonok Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** ? Tremadocian.

**Regional stratigraphic subdivisions:** Takoshkin and Tuloi regional stages (horizons).

**Local lithostratigraphic subdivisions:** Marcheta Formation.

Ordovician sections of the marine genesis in the vicinity of Ust'-Muta Village occur in separate fault blocks on the left bank of the lower Marcheta River (Figs 79, 80). The section (S-012) of the Marcheta Formation (Zasur'ya Group) is located on the left bank of the Marchetyonok Brook (Marcheta left tributary), 1350 m far from its mouth, and comprises.

The section of the Marcheta Formation (S-002) crops out on the left bank of the Marcheta River (left tributary of the Muta River), 900 m far from the Marchetyonok Brook inflow. The section begins at the floodplane terrace, then follows the second ridge from the Marchetyonok value, reaches the Marcheta/Marchetyonok divide, and ends at the top of 1201.0 m mountain. It comprises siltstone, mudstone, chert (Fig. 80).

Fig. 79. Sketch map of the Marcheta area.




Fig. 80. Violet-reddish mudstone from the Marchetyonok Section.

### **Marcheta-1 Section**

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Tremadocian, Floian.

**Regional stratigraphic subdivisions:** Tuloi and Lebed' regional stages (horizons).

**Local lithostratigraphic subdivisions:** Marcheta Formation.

**Zone:** *proteus* conodont Zone.

**Fauna:** conodonts, radiolarians, demosponges.

#### ***Peculiarities in facies and sedimentary environments.***

Member 6 of the Marcheta-1 Section composed of massive chert was formed on the mounds in the zone with underwater hydrotherms activity (?hydrothermal quartzites, ?jasperoids) (Figs 81, 82).



**Fig. 81.** Massive chert (?hydrothermal quartzites, ?jasperoids) from the Marcheta-1 Section (Member 6).

### **Marcheta-2 Section**

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Floian.

**Regional stratigraphic subdivisions:** Tuloi Regional stage (Horizon).

**Local lithostratigraphic subdivisions:** Marcheta Formation.

**Zone:** *elegans* conodont Zone.

**Fauna:** conodonts, radiolarians, demosponges.

Another section of the Marcheta Formation (S-013) (Fig. 83) crops out 300 m westward of the Marcheta-1 Section, opposite a spring on the left bank of the Marcheta River. The section begins 100 m far from the hill toe. It exposes upper strata of the Marcheta-1 Section but with slightly different lithologies and different member thicknesses.

All members, about 360 m of total thickness, belong to the Marcheta Formation.

Lithologies, especially in siliceous members, grade rapidly into one another along the strike in the three Marcheta sections: 6 m of red chert pass into red low-silica mudstone and siltstone within a distance of 30-50 m and then into gray mudstone and siltstone 50-70 m away. Color changes still more rapidly from red or lilac to sea-green, gray, dark olive, etc. Loaf-shaped concretions and traces of clay flowage found in the Marchetyonok Section and syndepositional breccia in the Marcheta-2 Section are typical of slope facies. Taking into account the marine origin of the Zasukh'yevskaya sediments, one may assume that the Marchetyonok Section formed on a seamount slope and the Marcheta-1 and Marcheta-2 sections were deposited on the sea floor in a vicinity of such a slope.

#### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

The lithofacies characteristics and low formation rate of the Zasukh'yevskaya Group (including the Marcheta Formation) (Sennikov et al., 2002, 2003, 2004; 2011b) suggest that these rocks formed in an oceanic environment (below the CCCD – Calcium Carbonate Compensation Depth) at depths corresponding to 4000–5000 m in the present-day oceans.



System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Radiolarians	Conodonts	Siliceous sponge spicules
Ordovician	Lower	Tremadocian	Marcheta	proteus	35	50	Mudstone: dark olive-green clayey.	Gen. et sp. indet. ■ ■ ■ ■ ■ Paracordylodus gracilis Lindstrom Oistodus gracilis Lindstrom Oistodus sp. Paroistodus cf. proteus (Lindstrom) Cornuodus longibasis Lindstrom Acontiodus reclinator Lindstrom Acontiodus (?) sp. Oncontodus sp.		
					34	2	Mudstone: reddish-gray siliceous-argillaceous.			
					33	30	Sandstone: green and dirty gray fine.			
					32	10	Mudstone: green and dirty gray silty sandstone grading upward into siltstone and then to member 32.			
					31	30	Sandstone: green or dirty gray, poorly sorted, fine massive, of uncertain bedding.			
					30	1.5	Siltstone: lilac, clayey.			
					29	35	Sandstone: green, poorly sorted, fine.			
					28	15	Mudstone: dark olive-gray, clayey.			
					27	10	Siltstone: dirty gray, clayey.			
					26	8	Mudstone: reddish brown, siliceous-argillaceous.			
					25	5	Siltstone: light-green, clayey.			
					24	110	Mudstone and fine sandstone: green.			
					23	117	Mudstone: gray or dark olive-gray, clayey.			
					22	3	Mudstone: dark red, argillaceous and siliceous-argillaceous.			
					21	28	Mudstone: bright green, siliceous-argillaceous.			
					20	0.7	Chert (possibly, hornfelsed): red or locally yellowish, with radiolarians and siliceous sponge spicules.			
					19	34	Mudstone: green, clayey.			
					18	5	Siltstone or silty sandstone: bright green.			
					17	60	Rocks similar to those in member 11, with gray-green or sea-green hues in upper layers.			
					16	3	Mudstone: dark olive-green, clayey.			
					15	20	Rocks similar to those in member 11, with sporadic 1-3 cm long lenses of green clayey siltstone.			
					14	4	Mudstone: dark olive, clayey.			
					13	8	Rocks similar to those in member 11.			
					12	2	Mudstone: dark olive-green, clayey, lumpy, and sandstone, fine.			
					11	0.5	Sandstone, locally gravelstone: greenish-gray, coarse.			
					10	1.1	Mudstone: lilac, clayey.			
					9	0.1	Mudstone: lilac with gray hue, clayey, with small lenses of gray chert.			
					8	0.3	Mudstone: lilac with gray hue.			
					7	23	Mudstone: sea-green, uncertainly bedded, massive.			
					6	2-6	Chert (hydrothermal jaspellite?): red, uncertainly bedded, massive, largely impregnated with ore minerals (hematite); member is lens-shaped: about 40 m long and 0 to 2-6 m thick.			
					5	7	Mudstone: sea-green, clayey, uncertainly bedded, massive, containing a 10-15 cm thick layer with 10 x 5 cm lenses or beads of red rocks from overlying member 6.			
					4	8	Rocks similar to those in member 2 but not banded.			
					3	10	Mudstone: cream-colored or brown, clayey, massive or lumpy, of uncertain bedding.			
					2	25	Mudstone: dark olive-gray or light greenish, siliceous-argillaceous, with thin (0.5 cm) banding produced by shade variation and high silica contents.			
					1	70	Mudstone: light green or dark olive, clayey, slightly siliceous, thin-banded, flaggy (1-2 cm) in lower 10 m and massive or lumpy above.			
				Marcheta	?					
					?					
					?					
					?					
					?					
					?					

Fig. 82. Lithology and ranges of fossil taxa from the Marcheta-1 Section.

Considering the possibly great CCCD in the Paleozoic, the first estimate of the paleobasin depth in the Altai segment of the Paleosian Ocean equals 1500–2000 m. Such depths, shallow on the ocean scale, are indirectly confirmed by the dominance of volcanic rock samples with the characteristics of paleoceanic uplift basalts. For bioindicator control of this value, let us analyze the taxonomic diversity of the faunal fossil assemblages in the Zashur'ya Group, the density of taphocoenosis, their morphologic complexity and skeleton preservation. Apart from the above-mentioned conodonts, the Zashur'ya Group chert contains radiolarians and sponge spicules. Microfossils have not been found in the white, gray, or green chert. Siliceous sponges spicules are relatively rare in the Zashur'ya Group chert and do not form large accumulations; therefore, most of these chert is radiolarite, not spongolite radiolarite. Most of the faunal fossils are associated with the red chert; a smaller proportion, with the brown-red and lilac ones. The terrigenous rocks

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Radiolarians	Conodonts	Siliceous sponge spicules
Ordovician	Lower	Floian	Marcheta	elegans	12	3	Mudstone: lilac, clayey, wavy-bedded, lumpy.	Gen. et sp. indet. ■  ■ <i>Juanognathus</i> (?) sp. ■ <i>Parioistodus</i> cf. <i>paralellus</i> (Pander) ■ <i>Parioistodus</i> sp. ■ <i>Drepanoistodus</i> sp.  Gen. et sp. indet. ■		
					11	70	Randomly interbedded fine sandstone, siltstone, and mudstone, greenish-gray to dirty dark olive-gray; a 2-m lens of lilac clayey mudstone in upper layers shows 1-3 mm banding and isometric 1-2 cm patches.			
					10	60	Mudstone, massive, and siltstone, greenish-gray or dirty gray, with sporadic nodules of greenish-gray fine sandstone.			
					9	12	Mudstone: lilac or green-gray, massive, lumpy; rocks change color rapidly both laterally and upwards: within 30 m along strike they pass into lilac-red clayey mudstone with siliceous nodules; a lens of red chert, 2-3 m long and 2-3 m thick.			
					8	5	Mudstone: greenish-gray, clayey, massive.			
					7	12	Mudstone: red-lilac or red-lilac-gray, clayey, of uncertain bedding, with round nodules of more siliceous rocks; rocks at 8 m above member base show weakly undulate 2 cm banding.			
					6	10	Mudstone: lilac or gray, clayey, lumpy; colors are randomly distributed and change rapidly from lilac to gray both laterally and upwards.			
					5	3	Syn depositional breccias in a matrix of dark olive-gray mudstone, with 1-3 cm clasts (occasionally 5-7 cm) of lilac, gray, or reddish mudstone more siliceous than the matrix; clasts often consist of thinly (3-5 mm) interbedded lilac and gray rocks. Clasts occasionally occupy up to 50 % of rock volume, but become progressively less abundant up the layer.			
					4	6	Sandstone: greenish-gray (to sea-green), fine.			
					3	9	Mudstone: silver-gray, clayey.			
					2	17	Mudstone: variegated, lumpy; rocks are mainly of dirty gray and dark colors; lilac and dirty lilac rocks occur in bedding-parallel lenses (patches) with either diffuse or sharp boundaries, looking like clasts (up to 20 cm); there are also <3 cm patches (clasts?) of red, bright lilac or green rocks; very thin bedding of 0.5 mm occurs occasionally.			
					1	150	Mudstone: pale dark olive-gray, clayey, massive.			

Fig. 83. Lithology and ranges of fossil taxa from the Marcheta-2 Section.

of the studied stratigraphic unit contain neither benthic nor pelagic fossils. The absence of benthos (except siliceous sponges) or widespread planktonic organisms such as graptolites and chitinozoans suggests that the paleobasin was fairly deep during the formation of the siliceous sediments of the Zasur'ya Group. The radiolarian taxonomic diversity in the Zasur'ya Group is low (several species within two genera in the open nomenclature); the morphology is simple; taphocoenosis density is low to medium (>10 to 10–100 specimens/cm<sup>2</sup> of the rock thin section or chip); the skeleton preservation is >10 % to 10–25%. The conodont fragments in the Zasur'ya Group show medium preservation; their taxonomic diversity in various zonal time-stratigraphic units is estimated as medium (3–5 species within 2–3 genera); the taphocoenosis density, as high (up to 10 specimens/locality). Considering the data on present-day basins, the high taxonomic diversity of radiolarians in the Zasur'ya Group may suggest that their communities lived in the lower part of the cool-water zone (200–500 m) and in the cold-water zone (500–5000 m). In the Charysh succession, their taphocoenosis are of medium density; in the other successions, a low density. This suggests that the upper distribution limit of the radiolarians is 200–300 m in the Charysh succession and over 300–500 m in the other successions. The poor and very poor skeleton preservation of the studied radiolarian assemblage suggests that the dead radiolarians covered considerable distances (300–700 m and more) before reaching the paleobasin bottom. Although the calculations are arbitrary, it is estimated on the basis of bioindicators that the Zasur'ya Group formed at depths of 500–1000 m in the Charysh area and at depths of 700–1200 m in the other areas. This is shallower than the first estimated depth of the paleoceanic rises in the Altai segment of the Paleoasian Ocean (1500–2000 m).



### Tekhten' Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Katian, Hirnantian, Rhuddanian, Aeronian.

**Regional stratigraphic subdivisions:** Tekhten', Listvyanka, Vtorye Utyosy and Syrovaty regional stages (horizons).

**Local lithostratigraphic subdivisions:** Tekhten', Vtorye Utyosy, Syrovaty formations.

**Fauna:** tabulate corals, rugose corals, trilobites, brachiopods, crinoids, graptolites.

The stratotype section of the Upper Ordovician Tekhten' Formation (Tekhten' Regional Stage) and the overlying Lower Silurian Vtorye Utyosy Formation is located on the right bank of the Tekhten' Brook, the right tributary of the Muta River, near Ust'-Muta Village (Fig. 84). The formation, in a much smaller stratigraphic volume, was formerly named the Dietken Formation according to Dietken, wrong respelling of an Altai name used in topographic maps. The documented section begins on the right side of a small ravine, 400 m from point 1181.0 m at azimuth 210°. The main section rises from the hill toe to point 1181.0 m and continues to the top at 1451.0 m. Members 1 and 2 and lower Member 3 crop out 200 m to the north of the section origin where the section begins with Member 3 (Figs 85, 86).

The section continues as far as the Muta/Anui river divide and then along the left side of the Surta Ravine (left bank of the Anui River near Bely Anui Village) as exposed alternating sandy-silty rocks with rare limestone layers containing rather representative Llandoveryan corals and brachiopods.

Some members of the section are clearly traceable




Fig. 84. Sketch map of the Ust'-Muta Village area.



Fig. 85. General view of the Tekhten' Section (middle part).

System		Series		Upper		Katian		Tekhten'		Rhuddanian		Syrtoat		Lithology		Tabulate corals		Rugose corals		Brachiopods		Trilobites		Graptolites	
Member No.	Thickness, m																								
14	>80													Interbedded (0.3-0.5 m) clayey siltstone and fine limy sandstone, greenish-gray; member lower half contains Early Silurian brachiopods; upper half of the member encloses gray limestone layers varying on strike from 5 to 10 m thick. The limestone contains a layer replete with fossils: numerous Silurian rugose corals.											
13	80													Sandstone: dirty yellow or pale lilac, rather quartz, polymictic, fine, occasionally with layers of dirty green clayey siltstone.											
12	15													Coarse clastic rocks, in three units: (i) lower 5 m of reddish-gray fine conglomerate (with quartz, limestone, siltstone, and sandstone pebble), (ii) middle 5 m of gravelstone, and (iii) upper 5 m of coarse polymictic rather quartz sandstone.											
11	5													Sodded interval.											
10	15													Siltstone: light gray, highly calcareous.											
9	30													Sandstone: dirty yellow, highly calcareous, polymictic, fine, with up to 30 cm thick layers of calcareous siltstone in the middle which locally grades on strike into clayey limestone.											
8	120													Sandstone: light gray or reddish-brown, quartz, medium and coarse.											
7	15													Siltstone: greenish, calcareous.											
5-6	120													Interbedded gray, yellowish-gray or greenish-gray clayey siltstone and fine polymictic limy sandstone, with a 5-m thick layer of black siltstone, and layers of quartz sandstone nearby, at member base; in most of middle layers, polymictic sandstone gives way to quartz sandstone; sandstone layers generally predominate and contain floating pebble and gravel; gravel is locally abundant enough for the rock to be classified as gravelstone.											
4	>30													Limestone: black, eluvial.											
3	80													Limestone: gray, slightly clayey, massive, containing 0.3 x 1.0 algal-biohermal buildups.											
2	40													Limestone: black, pelitic.											
1	>15													Interbedded siltstone and silty sandstone, dark olive-gray.											

Fig. 86. Lithology and ranges of fossil taxa from the Tekhten' Section.



southeastwards and northwestwards providing additional faunal evidence. For instance, black flaggy limestone, an equivalent of Member 2 on the right side of the upper Shiroky Ravine (a small ravine across the right Tekhten' watershed), 470 m far from point 1251.0 m at azimuth 35° (loc. 2040, 2041) contains Late Ordovician rugose corals *Grewinkia altaica* (Tcherepn.), *Parabrachielasma lebediensis* Tcherepn., *Ditoecholasma kanica* (Tcherepn.), tabulate corals *Cyrtophyllum kaninensis* Dziubo, *C. samyshiensis* Dziubo. Equivalent of Member 10 on the southern slope of point 1402.0 m, (loc. 142) trilobites *Acernaspis* (*Eskaspis*) *superciliexcelsis* Howells, *Acer.* (*Escaspis*) *becsciensis* Lesp. et Leten., *Podowrinella* cf. *striatonensis* Clarks et al., Warburgenellinae, (loc. 2039) trilobites *Acernaspis* (*Eskaspis*) *superciliexcelsis* Howells, *Acer.* (*Escaspis*) *xynon* Howells, Warburgenellinae, brachiopods *Isorthis prima* Walm. et Boucot, *Leptaena* cf. *haverfordensis* Bancr., *Atrypa* (?) *lindstromi* Wenjuk., *Eospirigerina* (?) sp., rugose corals *Cyathactis* sp., *Holophragma* sp., as well as dendroid graptolites were collected.

Members 1 through 9 belong to the Tekhten' Formation, Members 10 through 13 belong to the Vtorye Utyosy Formation, and Member 14 belongs to the Syrovaty Formation. The thickness of the Tekhten' Formation in the section is about 470 m, the Vtorye Utyosy Formation is 115 m thick, and the incomplete thickness of the Syrovaty Formation is over 80 m.

### **Muta Section**

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Katian, Hirnantian.

**Regional stratigraphic subdivisions:** Tekhten' Regional stage (Horizon).

**Local lithostratigraphic subdivisions:** Tekhten' Formation.

**Zone:** *supernus* graptolite Zone.

**Fauna:** stromatoporoids, tabulate corals, rugose corals, trilobites, crinoids, graptolites.

A sequence of lithologies different from the Tekhten'-type Section of the same Upper Ordovician age occurs 3 km to the west of the Tekhten' Brook. The section exposed at the Muta site originates 2.5 km uphill from Ust'-Muta Village on the side of a ravine that crosses the Muta left watershed (Figs 87, 88).

### **Peculiarities in facies, faunal assemblages and sedimentary environments.**

Carbonate members in the Marcheta Section are interpreted as uplifted reef structures. They belong to the central parts of reef uplifts.



**Fig. 87.** General view of the Muta Section (upper part).





## 4.5. WESTERN GORNY ALTAI (Charysh–Inya facies zone)

### 4.5.1. AREA OF UST'-CHAGYRKA VILLAGE

#### Barany-1 Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** ? Tremadocian, Floian, Dapingian, Lower Darriwilian.

**Regional stratigraphic subdivisions:** Takoshkin, Tuloi (=Lebed'), Kuibyshevo and Kostinsky regional stages (horizons).

**Local lithostratigraphic subdivisions:** Suetka, Voskresenka and Bugryshikha formations.

**Fauna:** conodonts, trilobites, brachiopods, gastropods, chitinozoans.

The Voskresenka Formation in the vicinity of Ust'-Chagyrka Village crops out along the Barany, Voskresenka, and Tachalov brooks, the left tributaries of the Chagyrka River (Figs 89, 90). An outcrop on the right side of the Barany Brook, 300 m from 500.3 m mountain at azimuth 235° exposes lilac to red siltstone and sandstone of the Suetka Formation (Gorny Altai Group) overstepped, with a sharp angular unconformity, by (stratotype section of Voskresenka Formation, P-78032) (Figs 90–92).




Fig. 90. Sketch map of the Barany area.




Fig. 89. Sketch map of the Ust'-Chagyrka area.




Fig. 91. General view of the exposure of the Barany-1 and Barany-2 sections.

System	Series	Stage	Formation	Member No.	Thickness, m	Lithology	Brachio-pods	Trilobites	Conodonts	Other groups
Ordovician		Middle	Darriwilian	Bugryshikha	8	>5	Sodded interval; talus bears fine conglomerate of light gray (to 80 %) well rounded pebble and dark gray (to 20 %) quartzite.	Archaeorthis altaica Sev.		
					7	90	Sodded interval; talus bears rare clasts of greenish-gray and yellowish-gray siltstone.			
					6	8	Limestone: gray and dark gray, sandy, laminated, flaggy.			
					5	205	Siltstone: greenish-gray and yellowish-gray, sandy, laminated, flaggy.			
					4	2	Sandstone: limy, grading along strike into sandy limestone.			
					3	25	Siltstone: yellowish greenish-gray, flaggy.			
					2	5	Sandstone: quartzitic, coarse, grading along strike into gravelstone and fine conglomerate.			
					1	40	Siltstone: green, massive, with scarce floating sandstone pebble; green siltstone gives way along strike (right side of the Tachalov Brook) to lilac-red, violet, green, and yellowish-gray siltstone with lenses and thin layers of medium to boulder conglomerate with pebbles mostly of Gorny Altai Group rocks or sporadically fine red quartzite; calcareous siltstone.			
							<ul style="list-style-type: none"> <li>■ <i>Idiostrophia</i> cf. <i>costata</i> Ulr. et Coop.</li> <li>■ <i>Idiostrophia</i> sp.</li> <li>■ Gen. et sp. indet.</li> <li>■ <i>Plectocamara costata</i> Coop.</li> <li>■ <i>Ateleasma batunensis</i> Sev.</li> <li>■ <i>Idiostrophia costata</i> Ulr. et Coop.</li> </ul>			
							<ul style="list-style-type: none"> <li>■ <i>Illaeus</i> sp.</li> <li>■ <i>Ceraurina</i> <i>frequens</i> Tchug.</li> <li>■ <i>Eorobergia bipunctata</i> Tchug.</li> <li>■ <i>Pliomerullus amplissimus</i> Petrun. sp.n.</li> <li>■ <i>Kolymella plana</i> (Tchug.)</li> <li>■ <i>Pliomerops parasiensis</i> Petrun. sp.n.</li> <li>■ <i>Bathyrullus nonnufus</i> Tchug.</li> <li>■ <i>Glaphurus altaicus</i> Web.</li> <li>■ <i>Eccoptochile tchagryica</i> Petrun. sp.n.</li> <li>■ <i>Acodus eletsicus</i> Tolm.</li> <li>■ <i>Naimanodus degiarevi</i> Tolm.</li> <li>■ <i>Triangulodus lerapintinensis</i> (Crespin)</li> <li>■ <i>Trangshanodus langshanensis</i> An</li> <li>■ <i>Semiacontiodus</i> ? <i>infusianensis</i> (An et Ding)</li> <li>■ <i>Protopanderodus</i> sp.</li> <li>■ <i>Parapanderodus striatus</i> (Graves et Ellisson)</li> <li>■ <i>Juanognathus jaanussoni</i> Serpagli</li> <li>■ <i>Periodon</i> cf. <i>P. falbellum</i> (Lindstrom)</li> <li>■ <i>Drepanoistodus subrectus</i> (Br. et M.)</li> <li>■ <i>Panderodus</i> ? <i>nogami</i> (Lee)</li> <li>■ <i>Cooperognathus</i> sp.</li> </ul>			
							■ Graptolites			

Fig. 92. Lithology and ranges of fossil taxa from the Barany-1 Section.

The member top is truncated by a fault. The total thickness of the section is about 380 m.

Members 1 through 6 belong to the Voskresenka Formation and members 7 and 8 belong to the overlying Bugryshikha Formation. The Voskresenka Formation in the Barany Section is 285 m thick and the Bugryshikha Formation is 95 m (incomplete thickness).

Gastropods and crinoids were collected from sandy limestone equivalent to member 4 on the right side of the Tachalov Brook (loc. S-7736) and gastropods were found on the right side of the Voskresenka Brook (loc. S-7726). Chitinozoans extracted by acid dissolution of rocks were identified as *Desmochitina minor* Eisenack and *Desmochitina* sp. at loc. S-7726 and *Desmochitina rhenana* Eisenack and *Conochitina* sp. at loc. S-7736. These species indicate an Ordovician age of the host sediment (most likely Tremadocian-Darriwilian, judging by their distribution in the Gorny Altai). Gastropods from S-7726 identified as *Temnodiscus* sp. are similar, according to V.I. Byaly, to the *Temnodiscus* sp. known from the variegated member of the Lugovoi Formation in the Biryusa River catchment (Siberian craton) correlated with the Nyaya regional stage (Early Tremadocian). Thus, the fauna-bearing sandy limestone can be conventionally assigned a Tremadocian age.



### Voskresenka-1 Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Darriwilian, Sandbian, Katian, Hirnantian.

**Regional stratigraphic subdivisions:** Bugryshikha, Khankhara and Tekhten' regional stages (horizons).

**Local lithostratigraphic subdivisions:** Bugryshikha, Khankhara and Tekhten' formations.

**Zones:** *coelatus*, *teretiusculus*, *serratulus*, *multidens* graptolite zones.

**Fauna:** tabulate corals, trilobites, brachiopods, ostracods, crinoids, bryozoans, gastropods, graptolites, polychaets, siliceous sponges, conodonts.

The most complete and stratigraphically continuous Ordovician section crops out of near Ust'-Chagyrka Village, in the Tachalov/Voskresenka brook divide (Fig. 93), 200 m from 530.3 m mountain at azimuth 15° and on northeastward down the divide. The rocks show monoclinical bedding to a dip of 75–80°. The section (S-78115) consists of (Fig. 94):

Members 1 through 10 in S-78115 belong to the Bugryshikha Formation, members 11 through 14 belong to the Khankhara Formation, and members 15 through 17 belong to the Tekhten' Formation. Thus, the Bugryshikha Formation in the Voskresenka-1 Section totals a thickness of about 140 m, the Khankhara Formation is more than 60 m thick, and the Tekhten' Formation is over 115 m.

The S-78115 section includes the graptolite zones: *coelatus* (lower half of member 6), *teretiusculus* (upper half of member 6 and lower half of member 7), *serratulus* (upper half of member 7), and *multidens* (member 8).




Fig. 93. Sketch map of the Voskresenka-Thachalov area.

System		Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Brachio-pods	Trilobites	Graptolites	Other groups
Ordovician	Upper	Sandbian	Khanakhara	Tekhten'		17	25	Limestone: gray or dark gray, massive, with crinoids.	<i>Plectorthis</i> aff. <i>altatus</i> Sev. <i>Leptellina</i> cf. <i>tennessensis</i> Uir. et Coop. <i>Anoplambonites grayae sibirica</i> Sev. <i>Hesperorthis</i> sp. <i>Leptellina</i> sp. <i>Hesperorthis</i> ex gr. <i>markovae</i> Rozm. <i>Isophragma</i> sp. <i>Skenidioides</i> sp. <i>Rosticellula</i> ex gr. <i>lenaensis</i> (Nikit.) <i>Lonchodomas</i> (Foliopyge) cf. <i>tardus</i> Petrun. <i>Lonchodomas</i> cf. <i>ichakyrensis</i> Petrun. <i>Bronzeopsis</i> sp. <i>Remopleurides</i> sp. <i>Encrinurides</i> sp. <i>Calyptauia</i> sp. <i>Pliomeros</i> sp. <i>Nileus</i> sp.	<i>Dicellograptus</i> aff. <i>moellensis</i> Caruthers <i>Amplexograptus coelatus</i> (Lapworth) <i>Cryptograptus tricinctus insectiformis</i> Ruedemann <i>Glossograptus limbatus</i> (Hopkinson) <i>Pseudoclimacograptus scharenbergi</i> (Lapworth) <i>Diplograptus multidentatus</i> Elies et Wood <i>Hustedograptus teretiusculus</i> (Hisinger) <i>Cryptograptus tricornis</i> (Caruthers) <i>Amplexograptus serratus</i> (Hall) <i>Acrograptus serratus</i> (Hall) <i>Climacograptus aff. perexcalatus</i> (Lapworth) <i>Amplexograptus perexcalatus</i> (Lapworth)	<i>Scelocodonts</i> <i>Gastropods</i> <i>Bryozoans</i> <i>Ostracods</i> <i>Tabulate corals</i> <i>Crinoids</i>	
						16	10	Silicite (chert): greenish-yellow or greenish-gray, massive, locally thinly banded (1-3 cm); with well or quite well preserved radiolarians, siliceous sponge spicules, and conodonts.				
						15	90	Sandstone: gray or dirty yellow to gray, rather quartz, fine, with rare thin (1-5 cm) interbeds of clayey siltstone, with graptolites at 15 and 40 meters above member base.				
						14	18	Silty sandstone and clayey siltstone, greenish-gray; with trilobites (2 m above member base).				
						13	22	Mudstone, clayey, and siltstone, dark gray or greenish-gray; with graptolites (8 meter above member base).				
						12	16	Siltstone: black, clayey, strongly foliated.				
						11	5	Limestone: gray or dark gray, crystalline; with brachiopods (2 m above member base).				
						10	4.5	Clayey mudstone and siltstone, gray or greenish-gray; with brachiopods (1 m above member base).				
						9	20	Sandstone: dirty brown, fine to medium; with brachiopods (5 and 18 meters above member base).				
		Dartwillian	Bugryshikha				8	22	Siltstone: pale yellowish-green or gray, clayey, strongly foliated; with graptolites (1 m above member base).	<i>Dicellograptus</i> aff. <i>moellensis</i> Caruthers <i>Amplexograptus coelatus</i> (Lapworth) <i>Cryptograptus tricinctus insectiformis</i> Ruedemann <i>Glossograptus limbatus</i> (Hopkinson) <i>Pseudoclimacograptus scharenbergi</i> (Lapworth) <i>Diplograptus multidentatus</i> Elies et Wood <i>Hustedograptus teretiusculus</i> (Hisinger) <i>Cryptograptus tricornis</i> (Caruthers) <i>Amplexograptus serratus</i> (Hall) <i>Acrograptus serratus</i> (Hall) <i>Climacograptus aff. perexcalatus</i> (Lapworth) <i>Amplexograptus perexcalatus</i> (Lapworth)	<i>Scelocodonts</i> <i>Gastropods</i> <i>Bryozoans</i> <i>Ostracods</i> <i>Tabulate corals</i> <i>Crinoids</i>	
							7	18	Siltstone: black or locally gray, clayey, with uneven shear surfaces; with graptolites (4 m meter above member base).			
							6	52	Interbedded fine clayey sandstone and siltstone, yellowish-gray or greenish-gray; with graptolites (2 m above member base).			
							5	2	Sandstone: dirty yellow, rather quartz, medium and coarse, clayey-carbonate cement.			
						4	1	Sandstone: dirty yellow, rather quartz, medium and coarse, clayey cement.	<i>Dicellograptus</i> aff. <i>moellensis</i> Caruthers <i>Amplexograptus coelatus</i> (Lapworth) <i>Cryptograptus tricinctus insectiformis</i> Ruedemann <i>Glossograptus limbatus</i> (Hopkinson) <i>Pseudoclimacograptus scharenbergi</i> (Lapworth) <i>Diplograptus multidentatus</i> Elies et Wood <i>Hustedograptus teretiusculus</i> (Hisinger) <i>Cryptograptus tricornis</i> (Caruthers) <i>Amplexograptus serratus</i> (Hall) <i>Acrograptus serratus</i> (Hall) <i>Climacograptus aff. perexcalatus</i> (Lapworth) <i>Amplexograptus perexcalatus</i> (Lapworth)	<i>Scelocodonts</i> <i>Gastropods</i> <i>Bryozoans</i> <i>Ostracods</i> <i>Tabulate corals</i> <i>Crinoids</i>		
						3	0.5	Sandstone: dirty yellow, rather quartz, medium and coarse, with well rounded floating quartz and quartzite pebble of 3-5 in diameter, siliceous-clayey cement.				
						2	7	Conglomerate: brown and dark gray, fine to medium, with well rounded quartz and quartzite pebble of 1-3 cm to 5-7 cm in diameter.				
						1	>10	Siltstone, dirty yellow and greenish-gray, thinly (3-5 cm) interbedded with mudstone or less often with fine polymictic sandstone, with sporadic floating quartz and quartzite pebble of 1-3 cm in diameter.				



### Tachalov Section

**Chronostratigraphic subdivisions of the International Stratigraphic Scale:** Katian, Hirnantian.

**Regional stratigraphic subdivisions:** Khankhara and Tekhten' regional stages (horizons).

**Local lithostratigraphic subdivisions:** Khankhara and Tekhten' formations.

**Zones:** *bicornis* and *supernus* (lower *supernus*, middle *ornatus* and upper *pacificus* subzones) graptolite zones and *A. ordovicicus* conodont Zone.

**Fauna:** tabulate corals, brachiopods, graptolites, polychaetes, conodonts, radiolarians, and chitinozoans.

The section occurs on the left side of the Tachalov Brook, left tributary of the Chagyrka River, near Ust'-Chagyrka Village where equivalents of the upper Voskresenka-1 Section (members 15 through 17) are exposed, namely (Figs 95–97):

Basal member 1a of the section belongs to the Khankhara Formation and members 1b through 3 belong to the Tekhten' Formation. The latter has an incomplete thickness of 130 m in the section.

The section includes *bicornis* and *supernus* (with lower *supernus*, middle *ornatus* and upper *pacificus* subzones) graptolite zones and *A. ordovicicus* conodont Zone.

### *Peculiarities in facies, faunal assemblages and sedimentary environments.*

The chert in the Tachalov Section and especially Voskresenka and Barany-2 sections is often banded and variously colored (owing to the codeposition of silica with metal hydroxides); this may suggest absent agitation of near-bottom water, which is enriched in silica, including biogenic silica. Consequently, such chert formed in basin bottom depressions in relative isolation not only from wave action (depths over 50–100 m) but also from near-bottom currents. In the Tachalov Section, the siliceous rocks (radiolarites) are underlain by well-sorted fine-grained sandstone interbedded with siltstone unaffected by wave action. Lenses of fine detrital limestone reflect the transport of bioclastic matter by near-bottom currents and its accumulation in microdepressions at the basin bottom. The sandstone and siltstone contain only pelagic fauna (graptolites,




Fig. 95. Sketch map of the Tachalov area.



















































































Fig. 96. General view of the Tachalov Section.

System		Series		Stage		Formation		Zone		Member No.	Thickness, m	Lithology	Radio-larians	Conodonts	Graptolites	Chitino-zoans	Tabulate corals	Ostracods	Brachio-pods	Scoleco-donts																																																																											
Ordovician		Upper		Katian		Tektien'		supenus-ornatus - pacificus		2	15																																																																																				
								~		3	0 - 40																																																																																				
												Silty sandstone and sandstone, rather quartz, fine, greenish to dark olive-gray; with bicornis zone graptolites (first meter above member base).		75		Sandstone: dirty yellowish-gray, rather quartz, fine, with layers of siltstone and 5-10 cm thick and up to 10 m long lenses of detrital limestone; member contains graptolites (first meter above member base).	Limestone: gray or light gray, massive.																																																																														

**Fig. 97.** Lithology and ranges of fossil taxa from the Tachalov Section.

chitinozoans), and the limestone lenses contain pelagic conodonts (*A. ordovicicus* Zone), benthic ostracods, and polychaetes. Conodont assemblages from the chert and limestone are very close in their taxonomic composition. The conodont fragments in the Tachalov Section show medium preservation, high taxonomic diversity, and medium taphocoenosis density (10–100 specimens/locality). In the studied successions of the upper Tekhten Formation in the Tachalov–Voskresenka area, the siliceous rocks (radiolarites) are immediately overlain by unstratified gray limestone. Outcrops of such carbonates are rounded in plan and lenticular in section; the lenticular shape is traced for a distance of 10–20 m (Barany-2 Section) to 100–150 m (Voskresenka and Tachalov sections). The massive outcrops of these carbonate units suggest that such limestone may be of extremely shallow-water algal-bioherm origin. Such proximity between the chert and limestone poses a question: either the chert with radiolarians (radiolarites) are not of deep-water origin or the limestone are not of shallow-water origin (200 meters and more) and do not have an allochthonous-bioherm character. The uncertain genesis of the chert and carbonate rock units suggested that a 200–250 m rapid regression (a local uplift of the basin bottom or a general sea level fall) had occurred before the chert stopped accumulating and the carbonates started to form. According to the global data and numerous reconstructions of T–R cycles, the World Ocean level fell abruptly by 80–100 m in the late Ordovician owing to the global glaciation (Brenchley, 2004; Haq and Schutter, 2008; Nielsen, 2004). No local orogeny which could have led to a large-scale bottom uplift in the Late Ordovician Altai basin was observed (Sennikov, 2006a, b). All this casts doubt upon the shallow-water origin of the carbonate sediments in the upper Tekhten' Formation in the Tachalov – Voskresenka area. Such massive gray organic-clastic limestone in the upper units of the Tachalov and Voskresenka sections, where poorly sorted bioclasts are embedded in a micritic groundmass and 30–80 % of the volume is accounted for by crinoidal and stromatoporoid(?) fragments as well as ostracod and brachiopod fragments and shells, should be assigned to bioclastic



wackestone-packstones (Varaksina and Sennikov, 2006). The above allochthonous wackestone-packstones with a large amount of micritic groundmass might have formed in large depressions at a considerable distance from shallow-water areas with autochthonous carbonate accumulation. In general, the carbonates crowning the Tachalov, Voskresenka, and Barany-2 sections might have formed in local closed depressions with respect to the relatively deepwater parts of the shelf (150–300 m). The chert in the Tachalov – Voskresenka and Barany-2 sections contain rich radiolarian assemblages (5–6 species within four genera), forming medium- and high-density taphocoenosis (10–100 and over 100 specimens/cm<sup>2</sup> of the rock thin section or chip). With the modern parameters for estimating the depths of warm-water radiolarian communities and their maximum population density, all this yields an average depth of 150–250 m. The good radiolarian preservation in the chert of the Tachalov Section (75–100 %) and the good radiolarian preservation in the Barany-2 Section (50–75 %) suggest that the radiolarian skeletons almost immediately reached the paleobasin bottom, without intense dissolution, and only in the sediment they dissolved very slightly. The above data suggest that the cherts (radiolarites) in the Tachalov and Barany-2 sections accumulated at depths of ~250–300 m.

#### **Voskresenka-2 Section**

***Chronostratigraphic subdivisions of the International stratigraphic scale:*** Katian.

***Regional stratigraphic subdivisions:*** Khankhara Regional stage (Horizon).

***Local lithostratigraphic subdivisions:*** Khankhara Formation.

***Zones:*** *caudatus* graptolite Zone.

***Fauna:*** graptolites.

Dirty yellow-gray fine sandstone of member 1 in the Tachalov Section extend along strike to the right side of the Voskresenka Brook where the Voskresenka-2 Section, stratigraphically higher than S-78117 and lower than S-7225. The Voskresenka-2 Section belongs to the Khankhara Formation and spans the *caudatus* graptolite Zone.

#### **Barany-2 Section**

***Chronostratigraphic subdivisions of the International stratigraphic scale:*** Katian, Hirnantian.

***Regional stratigraphic subdivisions:*** Tekhten', Listvyanka and Vtorye Utyosy regional stages (horizons).

***Local lithostratigraphic subdivisions:*** Tekhten', Listvyanka and Vtorye Utyosy formations.

***Zones:*** *persculptus* graptolite Zone.

***Fauna:*** trilobites, graptolites, radiolarians, siliceous sponges.

The Barany-2 Section (S-832 = S-0525) is located near Ust'-Chagyrka Village, on the right side of the lower Barany Brook where it enters the Charysh River valley. The section includes (Figs 98, 99):

Members 1 through 5 belong to the Tekhten' Formation and Member 6 belongs to the Vtorye Utyosy Formation. The incomplete thickness of the Tekhten' Formation in the section is about 20 m.

#### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

See above description for the Tachalov Section.



**Fig. 98.** Massive chert in the Tekhten' Formation, Barany-2 Section.







Ordovician							Lithology	Radio-larians	Grapto-lites	Other groups		
System	Series	Stage	Formation	Zone	Member No.	Thickness, m						
	Upper	Hirnantian	Tekhten'	Vityozhskiy	persculptus	6	5		Siltstone and mudstone: dark silver-gray, foliated.	<i>Borisella subulata</i> (Web. et Bl.) ■ <i>Entactinidae</i> gen. et sp. indet. ■ <i>Kalimnaspheera</i> sp. ■	Gen. et sp. indet. ■  <i>Normalograptus</i> ex gr. <i>persculptus</i> (Salter) ■     ■ Gen. et sp. indet.	Trilobites ■
						5	2-10		Limestone: gray, massive.			
						4	3-5		Silicite (chert): greenish-gray, gray, brown, or yellowish-dirty gray, banded, alternating with dirty gray or silver-gray siltstone.			
						3	1-3		Sandstone: dirty green-gray, polymictic, fine, including flat 3-5 cm thick and 20-30 cm long loaf-shaped lenses of gray or dark gray fine-grained bioclastic limestone; member is lens-shaped and pinches out.			
						2	10		Siltstone and mudstone: dirty green or gray, alternating with gray or greenish-gray silicite (chert).			
						1	20		Sandstone and siltstone: gray or dark olive-greenish-gray, polymictic, fine.			

Fig. 99. Lithology and ranges of fossil taxa from the Barany-2 Section.

#### 4.5.2. AREA OF MARALIKHA VILLAGE

##### Pichuzhikha Section

**Chronostratigraphic subdivisions of the International stratigraphic scale:** Floian, Dapingian, Dariwillian.

**Regional stratigraphic subdivisions:** Tuloi, Kuibyshevo, Kostinsky and Bugryshikha regional stages (horizons).

**Local lithostratigraphic subdivisions:** Voskresenka Formation.

**Zones:** *densus*, *gibberulus* graptolite zones.

**Fauna:** graptolites.

Stratigraphically higher strata of the Voskresenka Formation crop out on the right side of the Pichuzhikha Brook, 200 m far from the place where it enters the Charysh River terrace. The section from the water surface upwards includes (Figs 100–102):


The top of the section is truncated by a fault. Member 1 is conventionally equated to the *approximatus* Zone; Member 2 and lower 20 m, Member 3 correlate with the *balticus* Subzone of the *densus* Zone; upper 45 m, Member 3 make up the *densus* Subzone of the *densus* Zone; lower 30 m, Member 4 are conventionally correlated to the *angustifolius elongatus* Zone and its upper 40 m form the *gibberulus* Zone; Member 5 may correspond to the top of the Voskresenka Formation and may be assigned a middle Darriwilian age by analogy with the stratotype section of the Voskresenka Formation near Ust'-Chagyrka Village. Members 1 through 5 belong to the Voskresenka Formation and Member 6 may belong to the Bugryshikha Formation. The total thickness of the Voskresenka Formation in the Pichuzhikha Section is 180 m.



***Peculiarities in facies, faunal assemblages and sedimentary environments.***

Rocks composing the Pichuzhikha Section deposited on a large seamount, which is proved by the extremely thin Voskresenka Formation as compared to other sections of this formation. Given that the seamount was significantly removed from the provenance area of terrigenous material supplied from land, therefore only thin terrigenous sediments (dark-colored mudstone and siltstone) deposited on its top. The seamount was quite steep giving way to underwater landslide down the slope. The top of this seamount located at depths not exceeding the depth of storm waves action (below 50 m) was eroding due to landslide activity and affiliated water motion. Gravelite formed at the foot of the studied seamount in the form of thick beds with "curlers" and "twists" (Maralikha Section see below), created by underwater landslides, regularly descending from the slope of such seamount.

**Fig. 100.** Sketch map of the Maralikha area.



**Fig. 101.** Siltstone in the Voskresenka Formation, Pichuzhikha Section.








**Fig. 103.** General view of the middle part of the Maralikha Section.




**Fig. 104.** Lithology and ranges of fossil taxa from the Maralikha Section.

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Graptolites	Other groups
Ordovician	Middle	Darrivillian	Voskresenka	austrodentatus	11	250	Alternation of the siltstones and greenish-grey fine-grained sandstones. 50% of rocks are vesicular, with rare "twisting" sandy-siltstones comprised 3-5% of rock.	<ul style="list-style-type: none"> <li>Callograptus sp.</li> <li>Pseudograptus angustus Mu et Lee</li> <li>Isograptus caduceus (Salter)</li> <li>Isograptus cognatus (Harris et Thomas)</li> <li>Pseudisograptus manubriatus (T.S. Hall)</li> <li>Phyllograptus anna Hall</li> <li>Isograptus caduceus nanus (Ruedemann)</li> <li>Undulograptus sinodentatus (Mu et Lee)</li> <li>Undulograptus sinensis Obut et Sob.</li> <li>Acrograptus sp.</li> <li>Acrograptus compressus (Harris et Thomas)</li> <li>Acrograptus sinicus (Mu et Lee)</li> <li>Undulograptus austrodentatus (Harris et Kable)</li> </ul>	<ul style="list-style-type: none"> <li>Nautiloids</li> <li>Tritobites</li> </ul>
					10	120	Siltstones: grey vesicular with "twisting" structures occupied up to 90% of rock, upwards to the top of the bed reduced to 10%. Rare coarse-grained quartz sandstone pebbles were observed.		
					9	140	Siltstones: greenish and grey vesicular with "twisting" siltstones and fine-grained sandstones occupied up to 50 % of rock. Pebbles (from 5 to 15 cm in diameter) of fine-gained quartz sandstones have been observed.		
					8	80	Siltstones: vesicular with "twisting" structures of the same composition, occupied up to 50% of the rock. Their size in the lower part of the bed 1-3 cm to 0.5 m, gradually enlarged to up to 1-3 m the top of the bed. Rare angular white and bull quartz pebbles (up to 1 cm in diameter) were observed.		
					7	20	Siltstones: grey, fine schistose into large plates.		
					6	80	Siltstones: grey with single fine (5-7 cm, rarely 10 cm) light-grey to light-greenish grey, fine-grained polymictic sandstone intercalates. Siltstones possess fine cross-bedding.		
					5	0.5-1	Conglomerates: grey with middle to coarse pebbles and nodules. Pebbles are 2-5 cm rarely up to 12 cm in diameter. Pebbles are sorted due to clasts size, round, oriented by flatten surfaces according to bedding, other orientation was not observed. Pebbles occupied up to 80% of rock. In composition pebbles are represented by 50% dark-grey limestones, 35% siltstones and 15% mudstones. Mudstones are also observed as fragments of the "folded" intercalates (10-15 cm length, thickness 1-3 cm). Matrix consisted of fine-middle-grained sandstones and siltstones.		
					4	65	Alternation of grey fine-grained polymictic mudstones and sandstones.		
					3	25	Alternation of the vesicular siltstones and fine-grained polymictic bedded sandstones.		
					2	15	Siltstones: vesicular with "twisting"(from 3 cm to 1 m)siltstones and fine-grained sandstones, non oriented, occupied up 50 % of rock.		
					1	25	Alternation of greenish-gray siltstones and fine-grained polymictic sandstones.		
				? angustifolius elongatus - gibberulus				<ul style="list-style-type: none"> <li>Pseudotrigraptus cf. ensiformis (Hall)</li> <li>Glossograptus ex. gr. acanthus Elles et Wood</li> <li>Pseudotrigraptus angustus Mu et Lee</li> <li>Pendeograptus cf. pendens (Elles)</li> <li>Expansograptus suecicus (Tullberg)</li> <li>Acrograptus cognatus (Harris et Thomas)</li> <li>Expansograptus extensus (Hall)</li> <li>Isograptus caduceus imitatus Harris</li> <li>Paraglossograptus sp.</li> <li>Phyllograptus anna Hall</li> <li>Isograptus caduceus nanus (Ruedemann)</li> <li>Expansograptus taimyrensis Obut et Sob.</li> <li>Expansograptus hirsutus (Salter)</li> <li>Pseudotrigraptus sp.</li> <li>Loganograptus logani (Hall)</li> <li>Cardiograptus sp.</li> </ul>	<ul style="list-style-type: none"> <li>Conodonts</li> <li>Crinoids</li> <li>Brachiopods</li> </ul>





**Fig. 105.** Lithological peculiarities in the terrigenous rocks of the Maralikha Section (a-f) and in similar Silurian rocks (g).



**Fig. 106.** Diagram (in modern coordinates) for directions of non-lithified movement during underwater sliding in the Maralikha Section (A – member 6, B – member 8)



### ***Peculiarities in facies, faunal assemblages and sedimentary environments.***

The second half of the Maralikha Section is composed of rocks related to gravelite representing a specific lithological type – clastic sandy-aleuritic and argillite (siltstone) sediments. Such deposits formed on the slope of the seamount and slide down the slope as underwater landslides before the layers formed became lithified. As a result, some parts of the disrupted layer began to roll down the slope, acquiring gradually rounded shape with a more elongated axis across the direction of the movement. Some "curlers" and "twists" have internal structure in the form "spheres" insert into each other. The sediments formation at the slope foot followed by their compaction such "curlers" and "twists" became flattened. Such landslides may have been triggered by be giant storms, or evolved in places of abrupt terrain changes in the offshore parts inherent in transition areas (from shallow to deeper or redeepened shelf). One shouldn't rule out another most likely cause of underwater landslides as earthquakes. Regular appearance of units with "curlers" and "twists" in the Maralikha Section allows a most plausible assumption of a sharply defined crest on the shelf. Alternatively, this may be the presence of huge (up to 10 m in diameter) "curlers" and "twists" which can be attributed to olistostromes.

### **4.5.3. AREA OF BUGRYSHIKHA VILLAGE**

#### **Gora Altai Section**

***Chronostratigraphic subdivisions of the International stratigraphic scale:*** Darriwilian.

***Regional stratigraphic subdivisions:*** Bugryshikha Regional stage (Horizon).

***Local lithostratigraphic subdivisions:*** Bugryshikha Formation.

***Fauna:*** trilobites, brachiopods.

Middle Ordovician strata in the lower part of the composite stratotype section of the Bugryshikha Formation crop out near Bugryshikha Village, on the left side of the Belaya River and along the right and left sides of the lower Bugryshikha River (left tributary of the Belaya River). The lowermost member of the Bugryshikha Formation is exposed on Gora Altai (Fig. 107) that descends in a bluff into the Belaya River on its left side below the Bugryshikha inflow. The documented section (S-071) includes the following members, listed downstream (Fig. 108):

From the first and third members of the Gora Altai Section directly in the coastal cliff of Belaya River (locs 216 and 216a) trilobites *Lonchodomas sagittatus* Levit., *Nileus tengriensis* Web., *Homotelus angustus* Petrun., brachiopods *Archaeorthis altaica* Severg., *Glyptorthis primus* Severg., *Atelelasma subdorsoconvexum* Severg., *Ujukites tarlykensis* Andreeva have been found. From the rocks analogue to the third bed of the described section, on the flatten top of the Gora Altai (loc. 803) brachiopods *Ujukites tarlykensis* Andreeva were collected, and at the western slope of the Gora Altai, near Bugryshikha Village (loc. 178) brachiopods *Glyptorthis primus* Severg. Possibly, at the same stratigraphic level on the right bank of the Bugryshikha River, near Bugryshikha Village (loc. F-10) brachiopods *Atelelasma subdorsoconvexum* Severg. in Rozman. have been recovered.



**Fig. 107.** General view of the exposure along left bank of Belaya River, the Gora Altai Section.

System	Series	Stage	Formation	Member No.	Thickness, m	Lithology	Trilobites	Brachio-pods
Ordovician	Middle	Darriwilian	Bugryshikha	6	>100	Siltstone and sandstone: fine, lumpy, wave-bedded.	<ul style="list-style-type: none"> <li>■ <i>Lonchodomas communis</i> Lev.</li> <li>■ <i>Homotelus inferus</i> Lev.</li> <li>■ <i>Ampix</i> sp.</li> </ul>	
				5	40	Mudstone and siltstone: dark gray or black, thin-bedded; frequent pyrite crystals indicate deposition in anoxic conditions; siltstone contains small (5 cm long and 2 cm thick) mudstone lenses.	<ul style="list-style-type: none"> <li>■ <i>Lonchodomas sagittatus</i> Levit.</li> <li>■ <i>Nileus tengriensis</i> Web.</li> <li>■ <i>Homotelus angustus</i> Petrun.</li> </ul>	
				4	>10	Silty sandstone and siltstone: silver-gray or dark gray, lumpy, with traces of soft sediment slumping (up to 2-3 cm long and 1 cm wide tongues).		
				3	50	Mudstone and siltstone: gray, slightly calcareous, with conglomerate-like layers (lenses) with few floating carbonate concretions (3-5 cm in diameter) of dark gray clayey limestone.		<ul style="list-style-type: none"> <li>■ <i>Archaeorthis altaica</i> Severg.</li> <li>■ <i>Glyptorthis primus</i> Severg.</li> <li>■ <i>Ateleasma subdorskonvexum</i> Severg.</li> <li>■ <i>Ujukites tarykensis</i> Andreeva</li> </ul>
				2	0.2	Syn depositional breccia (size of clasts 1-2 cm, rarely 3 cm) of siltstone, mudstone, and fine sandstone in a siltstone matrix; clasts occupy up to 80 % of rock volume.		
				1	100	Mudstone, siltstone, and less often fine sandstone, gray.		

**Fig. 108.** Lithology and ranges of fossil taxa from the Gora Altai Section.

All members, of a total thickness of 300 m, belong to the lower Bugryshikha Formation.

The Gora Altai Section is extended upwards with the Bugryshikha Section, 1.5 km far to the northwest, on the other side of the same synclinal fold.

### Bugryshikha Section

**Chronostratigraphic subdivisions of the International stratigraphic scale:** Darriwilian.

**Regional stratigraphic subdivisions:** Bugryshikha Regional stage (Horizon).

**Local lithostratigraphic subdivisions:** Bugryshikha Formation.

**Zones:** *teretiusculus* graptolite Zone.

**Fauna:** trilobites, brachiopods, graptolites.

The Bugryshikha Section occurs northwest of the Gora Altai Section, 150 m from 580.8 m mountain at azimuth 190°. The documented section includes the following members, listed downhill toward Bugryshikha Village (Fig. 109):

All members belong to the lower half of the Bugryshikha Formation. The section is 850 m thick, and the total thickness of the composite stratotype section of the Bugryshikha Formation is 1000 m (without the lowermost and



System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Graptolites	Brachio-pods	Other groups
Ordovician	Middle	Darriwilian	Bugryshikha	<i>teretiusculus</i>	15	24	Siltstone: black.			
					14	390	Siltstone or less often fine sandstone: light greenish-gray.			
					13	65	Siltstone: brownish, greenish or dark olive-gray.			
					12	20	Siltstone: dark gray.			
					11	50	Interbedded siltstone and black fine sandstone.			
					10	20	Sandstone: light gray, fine.			
					9	>5	Mudstone and siltstone: black, clayey.			
					8	35	Sodded interval.			
					7	20	Siltstone: dark green to dark gray.			
					6	22	Sandstone and fine siltstone, dark gray or black, more rarely greenish-gray.			
					5	30	Sandstone and siltstone: light gray, polymictic, fine, gray.			
					4	5	Sodded interval.			
					3	145	Sandstone and siltstone: dark green to light green or brown, fine.			
					2	5	Sodded interval.			
					1	20	Siltstone and mudstone, variegated.			

Fig. 109. Lithology and ranges of fossil taxa from the Bugryshikha Section.

uppermost strata). Findings of graptolites in member 5 correspond to the *teretiusculus* Zone. According to graptolite, trilobite, and brachiopod assemblages, the Bugryshikha Formation in its stratotype section correlates with the Upper Darriwilian – Lower Sandbian.

The Gora Altai Section is extended upwards by the Malaya Uskuchevka Section located 1 km far to the southeast on the other side of the same anticlinal fold, on the right side of the Belaya River.

#### Malaya Uskuchevka Section

**Chronostratigraphic subdivisions of the International stratigraphic scale:** Darriwilian, Sandbian.

**Regional stratigraphic subdivisions:** Bugryshikha and Khankhara regional stages (horizons).

**Local lithostratigraphic subdivisions:** Bugryshikha and Khankhara formations.

**Zones:** *multidens* graptolite Zone (*antiquus lineatus*, *wilsoni* subzones).

**Fauna:** trilobites, brachiopods, graptolites.

A large part of the Bugryshikha and Khankhara formations crops out on the right side of the Malaya Uskuchevka River (right tributary of the Belaya River), uphill from the roadway. The section (S-8351) may be considered a parastratotype of the Khankhara Formation as its members exactly match those in the parastratotype section along the Bolshaya Uskuchevka River (Fig. 110):

System	Series	Stage	Horizon	Formation	Zone	Member No.	Thickness, m	Lithology
Ordovician	Upper	Sandbian	Bugryshikha	Bugryshikha	wilsoni	26	10	Mudstone: greenish-gray, limy, with sporadic carbonate concretions (to 0.5 mm in diameter).
						25	3	Limestone: gray and dark gray, slightly argillaceous.
						24	40	Mudstone: silver-gray, limy, strongly foliated.
						23	15	Siltstone: yellowish-dirty gray; limy cement.
						22	10	Sandstone: greenish-gray, highly calcareous, fine.
						21	20	Sandstone: yellowish-dirty gray, slightly calcareous, polymictic, with rare gray limestone concretions (to 5 cm in diameter) in upper layers.
						20	10	Limestone: gray, oolitic (oolites of <1 mm in diameter), flaggy in lower layers and massive in upper layers.
						19	40	Mudstone: bluish-gray, clayey, slightly calcareous, with "loaves" of gray limestone, from 3-5 cm to 10-15 cm in diameter, in lower layers.
						18	20	Sandstone: greenish-gray, rather quartz, polymictic, fine, with rare (at 1-3 m) clayey siltstone interbeds.
						17	40	Interbedded greenish-gray siltstone and clayey mudstone.
						16	15	Sandstone: greenish-gray, rather quartz, polymictic, fine.
						15	40	Mudstone: greenish-gray, locally dark gray, clayey, strongly foliated.
						14	10	Siltstone: light gray.
						13	30	Mudstone: dark gray, clayey.
						12	30	Siltstone: greenish-dirty gray to yellowish-gray; clayey cement.
						11	40	Mudstone: silver-gray clayey.
						10	100	Mudstone: bluish-gray, clayey.
						9	-6	Sandstone: rather quartz, medium-grained, polymictic, with sporadic 1 cm floating quartz pebble of low or medium roundness.
						8	> 1	Conglomerate: fine to medium, unsorted, with 0.5 to 3 cm, rarely to 5 cm quartz or quartzite pebbles of medium roundness.
						7	80	Mudstone: dark gray, locally almost black, clayey.
						6	25	Sandstone: greenish-gray, polymictic, fine.
						5	70	Interbedded siltstone and dark gray clayey mudstone.
						4	150	Mudstone: bluish-gray, clayey, massive, locally cavernous.
						3	200	Siltstone: dark silver-gray, nearly black, thick-bedded (10-20 cm), often massive, with scarce light bluish-gray interbeds.
						2	70	Siltstone: light gray or gray dark gray in upper layers, with black interbeds, clayey.
						1	>50	Siltstone: gray, light gray or locally bluish, clayey.

Fig. 110. Lithology and ranges of fossil taxa from the Malaya Uskuchevka Section.



Members 1 through 19 belong to the Bugryshikha Formation; members 20 through 26 belong to the Khankhara Formation. The incomplete thickness of the Bugryshikha Formation in the section is 980 m (without basal layers) and that of the Khankhara Formation exceeds 110 m. Trilobite and brachiopod assemblages indicate a Late Darriwilian and Early Sandbian age of the section. Graptolites correspond to the *multidens* Zone, including the *antiquus lineatus* Subzone at loc. S-7645 and the *wilsoni* Subzone at other localities.

Member No.	Thickness, m	Graptolites	Brachiopods	Trilobites
26	10			
25	3			
24	40			
23	15			
22	10			
21	20			
20	10			
19	40			
18	20			
17	40			
16	15			
15	40			
14	10			
13	30			
12	30			
11	40			
10	100			
9	~6			
8	> 1			
7	80			
6	25			
5	70			
4	150			
3	200			
2	70			
1	>50			

Fig. 110. The end.

#### 4.5.4. AREA OF KRASNOSHEKOV VILLAGE

##### Batun Section

**Chronostratigraphic subdivisions of the International stratigraphic scale:** Darriwilian.

**Regional stratigraphic subdivisions:** Kostinsky Regional stage (Horizon).

**Local lithostratigraphic subdivisions:** Voskresenka Formation.

**Zones:** *dentatus*, *balhaschensis* graptolite zones.

**Fauna:** graptolites, conodonts, trilobites, brachiopods.

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Graptolites	Conodonts	Trilobites	Brachiopods
Ordovician	Middle	Darriwilian	Voskresenka	balhaschensis	4	10	Limestone: yellow and yellow-gray, sandy and crystalline.	<ul style="list-style-type: none"> <li>Undulograptus sp.</li> <li>Undulograptus dentatus (Brong.)</li> <li>Dichograptidae</li> <li>Glossograptus fimbriatus (Hopk.)</li> <li>Glossograptus cf. acanthus Elles et Wood</li> <li>Pseudotrigonograptus sp.</li> <li>Isograptus (Nich.)</li> <li>Isograptus gibberulus</li> <li>Pseudisograptus manubriatus (T.S. Hall)</li> <li>Expansoagraptus extensus Obut et Zub.</li> <li>Expansoagraptus sp.</li> <li>Expansoagraptus balhaschensis (Keller)</li> </ul>	<ul style="list-style-type: none"> <li>Eopliacognathus pseudopliatus (Vitra)</li> <li>Periodon aculeatus Had.</li> <li>Pseudobolodina sp.</li> <li>Paroistodus onghialis (Serg.)</li> <li>Protopanderodus rectus (Lindstr.)</li> <li>Scolopodus giganteus Sw. et Bergstr.</li> </ul>	<ul style="list-style-type: none"> <li>Ceratinella cf. frequens Tschug.</li> <li>Bathyrrellus nonnullus Tschug.</li> <li>Camickia sp.</li> <li>Pliomera fischeri astatica Tschug.</li> <li>Kolymella aff. plana (Tschug.)</li> <li>Pliomerellus amplissimus Petrun.</li> <li>Pl. cf. jacuticus Tschug.</li> <li>Carollinites sp.</li> <li>Glaphurus alticus Weber</li> <li>Raymondaspis sp.</li> </ul>	<ul style="list-style-type: none"> <li>Trondorthis sibirica Severg.</li> <li>Orthidium fimbriatum Cooper</li> <li>Isophragma orientale Andreeva</li> <li>Idiostrophia costata Cooper</li> <li>Archaeorthis altica Severgina</li> <li>Ateleasma batunensis Severgina</li> </ul>
				dentatus	3	100	Siltstone: dark-gray, almost black, clayey.				
					2	20	Limestone: gray, sandy and crystalline.				
					1	~80	Sandstone: gray, greenish- and brown-gray, polymictic, mainly quartzitic, fine and medium-grained.				

The Batun Section occurs on the west of Krasnoshekovo Village, near former Kostinsky mine. Despite the poorly exposed section, it produce relatively diverse fossil groups. The Batun Section is the stratotype for the Kostinsky Horizon (Fig. 111).

##### Suetka Section

**Chronostratigraphic subdivisions of the International stratigraphic scale:** Katian.

**Regional stratigraphic subdivisions:** Tekhten' Regional stage (Horizon).

**Local lithostratigraphic subdivisions:** Siliceous-terrigenous Body.

**Zones:** *supernus* graptolite Zone.

**Fauna:** graptolites, conodonts, radiolarians.

The Suetka Section (loc. S-8223=S-0515-0518) occurs on the right side of the Suetka River, 2 km upstream of Suetka Village, on the southern slope of 323.6 m mountain (Fig. 112). The section consists of (Figs 113, 114):

The whole section, with a total thickness of about 70 m, belongs to the siliceous-terrigenous sequence and spans the *supernus* graptolite Zone.

**Peculiarities in facies, faunal assemblages and sedimentary environments.**

The chert beds in the Suetka Section (Maralikha – Suetka area) are up to 3 m thick. The chert is massive, gray, dark gray, and black, usually banded owing to lighter strips (0.1-1.0 cm) against a darker background. Also, light brown-yellow chert is observed, which show banding owing to color tints (1-3 cm). Conodonts (*A. ordovicicus* Zone) in the Suetka succession were found in lenticular beds of olistostrome limestone in the lower part of the succession, where graptolites (*supernus* – *ojsuensis* zones) were also found in siliceous mudstone. The conodont assemblage from the carbonate lenses is dominated by *Periodon grandis* (Ethington),

**Fig. 111.** Lithology and ranges of fossil taxa from the Batun Section.



in the Suetka Section can be called radiolarites. The high taxonomic diversity of the radiolarian assemblages with shells of complex morphology may indicate dwelling depths of 75–500 m. The high-density taphocoenosis propose dwelling depths of 150–250 m. The excellent preservation of the radiolarian skeletons in the lower part of the succession (like in



**Fig. 113.** Lithology and ranges of fossil taxa from the Suetka Section.



**Fig. 114.** Lithological peculiarities of the the chert and limestone of the Suetka Section.

a – c, e – chert; d – massive sitstone; f – lens of limestone.

the Tachalov one) proved that they did not dissolve and the lower distribution limit of the radiolarian communities was located near the basin bottom. The well-preserved radiolarian skeletons in the upper part of Suetka Section may suggest that the shells dissolved for a short time before reaching the paleobasin bottom, 100–150 m below their dwelling place. According to the bioindicator analysis, the chert from the lower part of Suetka Section might have formed at depths of 150–250 m, but the conodonts from the Suetka and Tachalov sections suggest depths of 300–350 m, which is more probable. Therefore, the paleobasin depths for the upper Suetka succession might have been 350–500 m. Note that graptolites were found only in the lower part of the succession. The absence of graptolites in the Altai successions with deep-water oceanic rocks indirectly confirms the relatively deep-water origin of the upper Suetka Section.



#### 4.5.5. AREA OF CHINETA VILLAGE

## Chineta Section

*Chronostratigraphic subdivisions of the International stratigraphic scale: Sandbian, Katian.*

**Regional stratigraphic subdivisions:** Bugryshikha, Khankhara, Tekhten' and Listvyanka regional stages (horizons).

**Local lithostratigraphic subdivisions:** Bugryshikha, Khankhara, Tekhten' and Vtorye Utyosy formations.

**Zones:** *teretiusculus*, *wilsoni* and *clingani* graptolite zones.

**Fauna:** trilobites, brachiopods, graptolites, conodonts.

A section that spans the Middle-Upper Ordovician Bugryshikha, Khankhara, and Tekhten' formations and the Lower Silurian Vtorye Utyosy Formation crops out at the northeastern end of Chineta Village, on the right side of the Inya River (Fig. 115). The documented section includes the following members, listed down the southern slope of 609.3 m mountain (Fig. 116):

On the southern slope of 609.3 m mountain, the section is extended with a repetition of Member 4 (see above), at least 100 m thick, composed of interbedded siltstone and fine sandstone, quite often slightly calcareous; in upper part there is an interbed of light gray fine to medium limy sandstone grading into sandy limestone; limestone contains trilobites and brachiopods (loc. S-812, 90 m far from point 609.3 m at azimuth 145°).

Equivalents of Member 2 on the other side of the fold north of point 609.3 m contain graptolites *Diplograptus* sp., *Dicellograptus* sp. (loc. 2302/1).

Members 1 and 2 belong to the Bugryshikha Formation, members 3 through 4 belong to the Khankhara Formation, Member 5 marks the base of the Tekhten' Formation, and Member 6 correlates with the Vtoroye Utyosy Formation.

Loc. S-795 in Member 2 may correspond to the *teretiusculus* graptolite Zone and loc. 2308 and 2308/1 in Member 4 apparently belongs to the *wilsoni* and *clingani* zones.

The incomplete thickness of the Bugryshikha Formation in the section is at least 220 m, the Khankhara Formation is more than 120 m thick, the incomplete thickness of the Tekhten' Formation is at least 40 m, and that of the Vtorye Utyosy Formation exceeds 30 m.

## Burovlyanka Section

*Chronostratigraphic subdivisions of the International stratigraphic scale:* Katian, Hirnantian, Rhuddanian, Aeronian stages.

**Regional stratigraphic subdivisions:** Tekhten', Listvyanka and Vtorrye Utyosy regional stages (horizons).

**Local lithostratigraphic subdivisions:** Tekhten' and Vtorye Utyosy formations.

**Zones:** *supernus-ornatus-ojsuensis*, *persculptus*, *acuminatus*, *sibiricus*, *cyphus* graptolite zones.


**Fauna:** trilobites, brachiopods, graptolites, conodonts, crinoids, algae.

Upper Ordovician and Lower Silurian strata crop out on the left bank of the Inya River opposite Chineta Village, on the divide of the Burovlyanka and Listvyanka brooks (Inya left tributaries) and along the left side of the Listvyanka Brook (Fig. 117). The fragment of section S-882 exposed on the southern slope of 591 m mountain, right upward from the Burovlyanka floodplain includes (Figs 118, 119).

Section S-833 runs parallel to S-822 on the southwestern slope of 635.4 m mountain, from the Burovlyanka Brook floodplain toward the Burovlyanka/Listvyanka divide and consists of:

On the left bank of the Listvyanka Brook, in 700 m upstream from its mouth from the equivalent of the upper part of Member 4 from S-833 section represented by siltstone (loc. S-824-2/10). Middle Llandoveryan *triangulatus*, *gregarius* zones graptolites were identified. Among them: *Demirastrites triangulatus* (Harkness), *Hedrograptus rectangularis* (McCoy), *Hedrograptus* sp., *Rastrites longispinus* Perner and *Monograptus* sp.

Siltstone in equivalents of Member 6 of S-833 on the left bank of the Listvyanka Brook, 500 m upstream of its mouth, contains Upper Llandoveryan graptolites of the *halli* Zone (loc. S-824-1/1), 120 m stratigraphically



**Fig. 115.** Sketch map of the Chineta area.

System	Series	Stage	Formation	Zone	Member No.	Thickness, m	Lithology	Graptolites	Trilobites	Conodonts	Brachiopods
Ordovician	Upper	Sandbian	Bugryshikha	teretiusculus	1	100	Interbedded mudstone, siltstone, and less often fine sandstone, bluish-greenish and silver-gray.	<i>Metaclimacograptus</i> sp. ■ <i>Hustedograptus</i> ex gr. <i>teretiusculus</i> (Hisinger) ■ <i>Climacograptus</i> sp. ■ <i>Dicranograptus</i> sp. ■ <i>Leptograptus</i> sp. ■ <i>Pseudoclimacograptus scharenbergi</i> (Lapworth) ■			
					2	120	Interbedded mudstone and siltstone, bluish-greenish and silver-gray.				
					3	3	Sandstone, light gray, limy, grading into sandy limestone.				
					4	120	Mudstone and siltstone, less often fine sandstone: silver-gray or brownish, slightly calcareous and clayey, with sporadic 0.5 m thick layers of highly calcareous siltstone grading into clayey-sandy limestone; bluish-greenish-gray siltstone contains trilobites.	<i>Diplograptus</i> sp. ■ <i>Orthograptus</i> sp. ■ <i>Diplograptus compactus</i> Elles et Wood ■ <i>Glyptograptus euglyphus</i> (Lapworth) ■ <i>Climacograptus tubiferus</i> Lapworth ■ <i>Dicellograptus</i> sp. ■			
					5	40	Limestone: gray, massive.				
					6	30	Mudstone: silver-gray or black, clayey.				
		Katian	Khankhara	wilsoni	?						
					?						
					?						
					?						
		Hirnantian	Tekhten'		?						
					?						
		Rhuddanian	Vitye Utyosy		?						
					?						

Fig. 116. Lithology and ranges of fossil taxa from the Chineta Section.







Fig. 117. Sketch map of the Burovlyanka area.

System		Series		Stage		Formation		Zone		Member No.		Thickness, m		Lithology		Graptolites		Trilobites		Other groups	
Ordovician	Upper	Silurian		Llandovery		Vtorye Utyosy		acumi-angustus-sibiricus cyphus		6 (4)		250		Siltstone and mudstone: dark greenish-yellowish-gray, foliated.		<ul style="list-style-type: none"> <li>■ <i>Glyptograptus</i> sp.</li> <li>■ <i>Normalograptus</i> sp.</li> <li>■ <i>Normalograptus</i> cf. <i>longifolius</i> Manck</li> <li>■ <i>Normalograptus ornatus ornatus</i> Elles et Wood</li> </ul>					
		Rhuddanian												Limestone: gray or black, clayey, flaggy.		<ul style="list-style-type: none"> <li>■ <i>Parakidograptus acuminatus</i> (Nicholson)</li> <li>■ <i>Hydrograptus</i> sp.</li> <li>■ <i>Parakidograptus</i> sp.</li> <li>■ <i>Parakidograptus moyeriensis</i> (Obut)</li> <li>■ <i>Metabolograptus sibiricus</i> (Obut)</li> <li>■ <i>Dimorphograptus</i> sp.</li> <li>■ <i>Cystograptus</i> sp.</li> <li>■ <i>Rhaphidograptus maslovi</i> Obut et Sob.</li> <li>■ <i>Metaclimacograptus</i> sp.</li> <li>■ <i>Coronograptus angustus</i> Obut</li> <li>■ <i>Pseudoclimacograptus</i> sp.</li> </ul>					
	Lower	Hirnantian				persculptus		5b 5a		10 2.1 0.8		50-190		Siltstone, silty sandstone, and fine limy-clayey sandstone, greenish-gray, locally black.		<ul style="list-style-type: none"> <li>■ <i>Appendispino-graptus supernus</i> (Elles et Wood)</li> <li>■ <i>Normalograptus</i> sp.</li> <li>■ <i>Normalograptus</i> (Koren et Mikh.)</li> <li>■ <i>Dicellograptus ornatus ornatus</i> (Salter)</li> <li>■ <i>Normalograptus persculptus</i> (Salter)</li> <li>■ <i>Normalograptus</i> sp.</li> <li>■ <i>Parakidograptus lorraineis</i></li> <li>■ <i>Glyptograptus</i> (Ruedemann)</li> <li>■ <i>Pseudoclimacograptus</i> sp.</li> <li>■ <i>Glyptograptus</i> sp.</li> <li>■ <i>Orthograptus</i> sp.</li> <li>■ <i>Normalograptus</i> sp.</li> </ul>					
		Tekhten'				supernus-onatus ojsuensis		4		2		2		Limestone: gray, algal-biohermal, in a greenish-gray silt-sandstone matrix; 1 x 2 m carbonate bioherms occupy 80-90 % of member volume.		<ul style="list-style-type: none"> <li>■ <i>Elles et Wood</i></li> <li>■ <i>Climacograptus acantus</i></li> </ul>					
Katian								?		2		10		Limestone: gray, algal-biohermal, with mud patches; 3x5 m bioherms occupy 80-90 % of member volume and occur in massive limestone.							
								1		>250				Limestone: gray or light gray, massive, with sporadic poorly preserved crinoids in upper layers.				<ul style="list-style-type: none"> <li>■ Crinoids</li> <li>■ Conodonts</li> </ul>			

Fig. 118. Lithology and ranges of fossil taxa from the Burovlyanka Section.



**Fig. 119.** Lithological peculiarities of limestone of the Burovlyanka Section.

a - b – algae bioherms in the terrigenous matrix (siltstone); c – Dalmanitina member clayey thin-bedded limestone; d - e – Dalmanitina member massive limestone; f – siltstone and sandstone.

higher than loc. S-824-2/10 with graptolites of the *triangulatus* and *gregarius* zones: *Stimulograptus halli* (Barrande), *Monograptus* sp., *Paradiversograptus capillaris* (Carruthers) and *Glyptograptus tamariscus* (Niholson).

Siltstone in equivalents of Member 6 of S-833 on the left bank of the Inya River, 100 m upstream of an island near the Listvyanka Brook inflow (loc. S-8341) contains graptolites of the Upper Llandoveryan *guerichi* Zone: *Spirograptus guerichi* Loydell, Storch et Melchin, *Rastrites* sp., *Stimulograptus halli* (Barrande), *Oktavites planus* (Barrande), *Paradiversograptus runcinatus* (Lapworth), *Paradiversograptus capillaris* (Carruthers), *Agetograptus tenuissimus* Sennikov, *Hedrograptus* sp., *Petalograptus* sp.

All six members of S-822 section and members 1–3 of S-833 section belong to Tekhten' Formation, and members 4 and 5 of S-833 section – to Vtorye Utyosy Formation, Member 6 of S-833 section to the Syrovaty Formation. Thickness of the Tekhten' Formation in the composite Burovlyanka Section (incomplete) is more than 330 m, Vtorye Utyosy Formation – about 420 m, Syrovaty Formation – more than 100 m.



## CONCLUSIONS

This book contains the paleontological, biostratigraphic, lithologic and biofacies data on the key Ordovician sections from five large parts of the Gorny Altai - Uymen'-Lebed', Teletskoe Lakeside, Biya-Katun', Anui-Chuya and Charysh-Inya facies zones. In addition to the above mentioned facies zones, the Ordovician sedimentary sections with a specific set of formations could be distinguished. These are interpreted either as separate paleo-offshore area of the unique Altai shelf paleobasin (Milovanoka and Vydrikha facies zones) or as fragments of another paleobasin (Ulagan facies zone). We carried out detailed studies and bed-by-bed descriptions of the reference sections from the Milovanoka, Vydrikha and Ulagan facies zones. Some of the paleontological-stratigraphic data obtained from them could be correctly aligned with those from the five main structural facies zone. However, most sequences from the Milovanoka, Vydrikha and Ulagan zones still require further study of litho- and biostratigraphy in order to generate paleogeographic and geodynamics reconstructions.

Authors hope that readers can get an overview of the structure of the Altai Ordovician sedimentary basins and the paleobiota inhabiting them. Geological materials from the Gorny Altai (South of West Siberia) could be a good example of the successive evolutionary development of sedimentary basins and the entire biosphere during the Ordovician period of the Earth's history.

## Acknowledgements

*The study was supported by Trofimuk Institute of Petroleum Geology and Geophysics SB RAS, and Russian Foundation for Basic Research (RFBR, project N 19-05-20060 - "13-th International Symposium on the Ordovician System"). This is contribution to IGSP 653 Project "The onset of the Great Ordovician Biodiversification Event".*

## REFERENCES

- Achab A. (1989.) Ordovician chitinozoan zonation of Quebec and Western Newfoundland // *Journal Paleontology*. Vol. 63, N 1. P. 14–24.
- Andreeva O.N. (1985). Middle Ordovician brachiopods and trilobites from Tuva and Altai // *Paleontologicheskii Zhurnal*. Issue 2. P. 38–47.
- Barnes C.R., Fortey R.A., and Williams S.H. (1996). The pattern of Global Bio-Event during the Ordovician period. In: *Global Events and Event Stratigraphy in the Phanerozoic. Results of the International Interdisciplinary Cooperation in the IGGP-Project 216 "Global Biological Events in Earth History"*. Springer-Verlag, Berlin, Heidelberg. P. 139–172.
- Bergström S., Finney S., Chen Xu, Palson Ch., Wang Zhi-hao, and Grahn Y. (2000). A proposed global boundary stratotype for the base of the Upper Series of the Ordovician System: The Fagelsang section, Scania, southern Sweden // *Episodes*. Vol. 23, N 2. P. 102–109.
- Bergström S., Lofgren A., and Maletz J. (2004). The GSSP of the Second (Upper) Stage of the Lower Ordovician Series: Diabasbrottet at Hunneberg, Province of Västergötland, Southern Sweden // *Episodes*. Vol. 27, N 4. P. 265–272.
- Bergström S., Finney S., Chen Xu, Goldman D., and Leslie S.A. (2006). Three new Ordovician global stage names // *Lethaia*. Vol. 39. P. 287–288.
- Bergström S.M., Chen X., Gutiérrez-Marco J.C., and Dronov A. (2009). The new chronostratigraphic classification of the Ordovician System and its relations to major regional series and stages and to  $\delta^{13}\text{C}$  chemostratigraphy // *Lethaia*. Vol. 42. P. 97–107.
- Bergström J., Pärnaste H., and Zhiyi Zhou. (2013). Trilobites and biofacies in the Early-Middle Ordovician of Baltica and a brief comparison with the Yangtze Plate // *Estonian Journal of Earth Sciences*. Vol. 62, N 4. P. 205–230.
- Brenchley P.J. (2004). End Ordovician Glaciation. In: *The Great Ordovician Biodiversification Event*. Webby B., Paris F., Droser M.L. and Percival I.G. (Eds). Columbia University Press. New York. P. 81–83.
- Brooke C., Riding R. (1998). Ordovician and Silurian coralline red algae // *Lethaia*. Vol. 31. P. 185–195.
- Bukolova E.V. (2011). The Middle Ordovician litho- and biostratigraphy of northwestern Gorny Altai // *Russian Geology and Geophysics*. Vol. 52, N 9. P. 979–990.
- Buslov M.M., Fujiwara Y., Safonova I.Yu., Okada Sh., and Semakov N.N. (2000). The junction zone of the Gorny Altai and Rudny Altai terranes: structure and evolution // *Russian Geology and Geophysics*. Vol. 41, N 3. P. 377–390.
- Buslov M.M., Safonova I.Yu., and Bobrov V.A. (1999). Exotic terrane of Late Cambrian-Early Ordovician oceanic crust in the north-western Gorny Altai (Zasurin Formation): Structural position and geochemistry // *Doklady Akademii nauk*. Vol. 368. Issue 5. P. 650–654.
- Chen Xu, Rong Jiayu, Fan Junxuan, Zhan Renbin, Mitchell Ch.E., Harper D.A.T., Melchin M.J., Peng Ping'an, Finney S.C., and Wang Xiaofeng. (2006). The Global Boundary Stratotype Section and Point (GSSP) for the base of the Hirnantian Stage (the uppermost of the Ordovician System) // *Episodes*. Vol. 29, no. 3. P. 183–196.
- Cherepnina S.K. [Черепнина С.К.] (1960). Subclass Tetracoralla (Rugosa). In: *Paleozoic biostratigraphy of the Altai-Sayan mountainous area*. L.L. Khal'fin (Ed.). Vol. I. Lower Paleozoic. Trudy SNIIGGiMS. Issue 19. Novosibirsk: SNIIGGiMS Publ. House. P. 361–362. (In Russian). [Подкласс Tetracoralla (Rugosa). Тетракораллы // Биостратиграфия палеозоя Саяно-Алтайской горной области. Т. I. Нижний палеозой / Под ред. Л.Л. Халфина: Тр. СНИИГГиМС. Вып. 19. Новосибирск: СНИИГГиМС]
- Chuvashov B.I., Luchinina V.A., Shuiskii V.P., Shaikin I.M., Barchenko O.I., Ishchenko A.A., Saltovskaya V.D., and Shirshova D.I. [Чувашов Б.И., Лучинина В.А., Шуйский В.П., Шайкин И.М., Барченко О.И., Ищенко А.А., Салтовская В.Д., Ширшова Д.И.] (1987). Fossil calcareous algae. Morphology, systematics, methods of studies. Novosibirsk: Nauka. 225 p. [Ископаемые известковые водоросли. Морфология, систематика, методы изучения. Новосибирск: Наука.]
- Chuvashov B.I., Shuysky V.P., and Ivanova R.M. (1993). Stratigraphical and facies complexes of the Paleozoic calcareous algae of the Urals // *Bollettino della Societa Paleontologica Italiana*. T. 1. Spec. Vol. C. 93.
- Contributions to Siberian IGGP 410/421 Joint Meeting. Ordovician-Lower Carboniferous succession, eustatic (sedimentary) cycles, event levels and palaeogeography of Southern West Siberia (Guidebook of excursion). (2001). Yolkin E.A., Sennikov N.V., Petrunina Z.E., Bakharev N.K., Izokh N.G., Klets A.G., Alekseenko A.A., Anastasieva S.A., Ermikov V.D., Gladkikh L.A., Gutak Ya.M., Khlebnikova T.V., Mamlin A.N., Obut O.T., Rodina O.A. and Tokarev V.N. Novosibirsk: Institute of Petroleum Geology, SB RAS. P. 8–92.
- Cooper R.A., Nowlan G.S., and Williams S.H. (2001). Global Statatotype Section and Point for base of the Ordovician System // *Episodes*. Vol. 24, N 1. P. 19–28.
- Cooper R.A., Maletz J., Taylor L., Zalasiewicz J. (2004). Graptolites: Patterns of diversity across Paleolatitudes // *The Great Ordovician Biodiversification Event*. Webby B., Paris F., Droser M.L. and Percival I.G. (Eds). Columbia University Press. New York. P. 281–293.
- Correlative stratigraphic charts. (1964). Trudy SNIIGGiMS, supl. Issue 29. Data on stratigraphy of the Sayan-Altai Folded Area. (In Russian). [Корреляционные стратиграфические схемы. (Тр. СНИИГГиМС. Приложение к вып. 29). Материалы по стратиграфии Саяно-Алтайской складчатой области. Новосибирск: Изд-во СНИИГГиМС. 9 прил.]



- Decisions of the All-Union stratigraphic meeting on Precambrian, Paleozoic and Quaternary Systems of Middle Siberia, Novosibirsk, 1979. (1983).* Pt. I. Upper Proterozoic and Lower Paleozoic. Novosibirsk. 215 p. (In Russian). [Решения Всесоюз. стратигр. совещ. по докембрию, палеозою и четвертичной системе Средней Сибири, Новосибирск, 1979). Ч. I. Верхний протерозой и нижний палеозой. Новосибирск.]
- Decisions of the Interdepartmental meeting on development of the uniform stratigraphic charts for Siberia. (1959).* Moscow: Gosgeoltekhizdat. 91 p. (In Russian). [Решения Межведомственного совещания по разработке унифицированных стратиграфических схем Сибири. М.: Госгеолтехиздат.]
- Dobretsov N.L. (2003).* Evolution of structures of the Urals, Kazakhstan, Tien Shan, and Altai-Sayan region within the Ural-Mongolian fold belt (Paleoasian ocean) // *Russian Geology and Geophysics*. Vol. 44, N 1–2. P. 3–26.
- Dobretsov N.L. (2011).* Early Paleozoic tectonics and geodynamics of Central Asia: role of mantle plumes // *Russian Geology and Geophysics*. Vol. 52, N 12. P. 1539–1552.
- Documents on stratigraphy of the Sayan-Altai Folded Area. (1964).* Trudy SNIIGGiMS. Issue. 29. Novosibirsk: SNIIGGiMS Publ. House. 161 p. [Материалы по стратиграфии Саяно-Алтайской складчатой области. Тр. СНИИГГиМС. Вып. 29. Новосибирск: Изд-во СНИИГГиМС.]
- Dubinina S.V. [Дубинина С.В.] (2000).* Conodonts and zonal stratigraphy of the Cambrian-Ordovician boundary deposits. Trudy GIN RAN. N 517. Moscow: Nauka. 239 p. (In Russian). [Конodontы и зональная стратиграфия пограничных отложений кембрия и ордовика. М.: Наука.]
- Dzyubo P.S. [Дзюбо П.С.] (1960a).* Hydrozoa Chaetetida. In: *Paleozoic biostratigraphy of the Altai-Sayan mountainous area. L.L. Khalfin (Ed.). Vol. I. Lower Paleozoic. Trudy SNIIGGiMS. Issue 19. Novosibirsk: SNIIGGiMS Publ. House. P. 373–374. (In Russian).* [Группа Hydrozoa Chaetetida. Биостратиграфия палеозоя Саяно-Алтайской горной области. Т. I. Нижний палеозой. Под ред. Л.Л. Халфина. Труды СНИИГГиМС. Вып. 19. Новосибирск: СНИИГГиМС.]
- Dzyubo P.S. [Дзюбо П.С.] (1960b).* Anthozoa Heliolitida. In: *Paleozoic biostratigraphy of the Altai-Sayan mountainous area. L.L. Khalfin (Ed.). Vol. I. Lower Paleozoic. Trudy SNIIGGiMS. Issue 19. Novosibirsk: SNIIGGiMS Publ. House. P. 382–387. (In Russian).* [Группа Anthozoa Heliolitida // Биостратиграфия палеозоя Саяно-Алтайской горной области. Т. I. Нижний палеозой. Под ред. Л.Л. Халфина. Труды СНИИГГиМС. Вып. 19. Новосибирск: СНИИГГиМС.]
- Dzyubo P.S. [Дзюбо П.С.] (1966).* Tabulate corals and heliolitids from Ordovician of the Gorny Altai and Gornaya Shoriya. Abstract of PhD dissertation. Tomsk: Tomsk University Press. 25 p. (In Russian). [Табуляты и гелиолитиды ордовика Горного Алтая и Горной Шории. Автореф. канд. дисс. Томск: Изд-во Томского ун-та.]
- Dzyubo P.S. and Mironova N.V. [Дзюбо П.С., Миронова Н.В.] (1960).* Subclass Tabulates. In: *L.L. Khalfin (Ed.) Paleozoic biostratigraphy of the Altai-Sayan mountainous area. Vol. I. Lower Paleozoic. Trudy SNIIGGiMS. Issue 19. Novosibirsk: SNIIGGiMS Publ. House. P. 374–382. (In Russian).* [Подкласс Tabulata. Биостратиграфия палеозоя Саяно-Алтайской горной области. Т. I. Нижний палеозой. Под ред. Л.Л. Халфина. Труды СНИИГГиМС. Вып. 19. Новосибирск: СНИИГГиМС.]
- Elles G.L. and Wood E.M.R. (1901-1918).* Monograph of British graptolites. Palaeontol. Soc. London. 539 p.
- Erdtmann B.-D. (1984).* Outline ecostratigraphic analysis of the Ordovician graptolite zones in Scandinavia in relation to the palaeogeographic disposition of the Japetus // *Geol. Palaeontol.* Vol. 18. P. 9–15.
- Erdtmann B.-D. (1986).* Early Ordovician eustatic cycles and their bearing on punctuations in early nematophorid (planktic) graptolite evolution // *Lect. Notes Earth Scien.* Vol. 8. P. 139–152.
- Erdtmann B.-D. and Miller J.F. (1981).* Eustatic control of lithofacies and biofacies changes near the base of the Tremadocian. Second Internat. Symposium on the Cambrian System. P. 78–81.
- Ermikov V.D., Zeyfert L.L., Petrunina Z.E., Puzyrev A.A., and Sennikov N.V. (1979).* Tremadocian of the northern part of the Gorny Altai // *Soviet Geology and Geophysics*. Vol. 20, N 11. P. 21–34.
- Gintsinger A.B. [Гинцингер А.Б.] (1958).* Ordovician, Silurian and Devonian stratigraphic chart for the Gorny Altai. Moscow: Gosgeoltekhizdat. Issue 61. P. 24–39. (In Russian). [Стратиграфическая схема ордовика, силура и девона Горного Алтая // Материалы по стратиграфии Западной Сибири. М.: Госгеолтехиздат.]
- Gintsinger A.B. [Гинцингер А.Б.] (1964).* Ordovician correlative stratigraphic chart for Altai, Salair and Gornaya Shoriya. Data on stratigraphy of the Sayan-Altai folded area. Novosibirsk: SNIIGGiMS Publ. House. Issue 29. P. 94–107. (In Russian). [Корреляционная стратиграфическая схема ордовика Алтая, Салаира и Горной Шории. Материалы по стратиграфии Саяно-Алтайской складчатой области. Новосибирск: Изд-во СНИИГГиМС.]
- Gintsinger A.B. and Sennikov V.M. [Гинцингер А.Б., Сенников В.М.] (1967).* Ordovician of the Altai-Sayan Area. In: *Paleozoic stratigraphy of the Central Siberia. Novosibirsk: Nauka. P. 74–86. (In Russian).* [Ордовик Алтае-Саянской области. Стратиграфия палеозоя Средней Сибири. Новосибирск: Наука.]
- Gintsinger A.B. [Гинцингер А.Б.] (1969).* Ordovician sections Kuznetsk Basin region margin of Salair. In: *Precambrian and Lower Paleozoic sections of the western part of Altai-Sayan folded area. Krasnoyarsk: Krasnoyarsk Publ. House. P. 189–209. (In Russian).* [Разрезы ордовика Прикузбасской окраины Салаира. Разрезы докембрия и нижнего палеозоя западной части Алтае-Саянской складчатой области. Красноярск: Красноярское книжн. изд.]
- Goldman D., Leslie S.A., Nolvak J., Young S., Bergström S.M., and Huff W.D. (2007).* The Global Stratotype Section and Point (GSSP) for the base of the Katian Stage of the Upper Ordovician Series at Black Knob Ridge, Southeastern Oklahoma, USA // *Episodes*. Vol. 30, N 4. P. 258–270.

- Gradstein F.M., Ogg J.G., Schmitz M.D., and Ogg G.M. (Editors). (2012). The Geologic Time Scale 2012. 2-volume set. Elsevier Science Ltd. 1176 p.
- Haq B.U. and Schutter S.R. (2008). A Chronology of Paleozoic Sea-Level Changes // *Science*. Vol. 322. P. 64-68.
- Ivantsov A.Yu. (2003). Ordovician Trilobites of the Subfamily Asaphinae of the Ladoga Glint // *Paleontological Journal*. Vol. 37, Suppl. 3. P. S229-S337.
- Iwata K., Sennikov N.V., Buslov M.M., Obut O.T., Shokal'skii S.P., Kuznetsov S.A., and Ermikov V.D. (1997). Late Cambrian-Early Ordovician age of basalt-siliceous-terrigenous Zasukh'yevskaya Formation (northwestern Gorny Altai) // *Russian Geology and Geophysics*. Vol. 38, N 9. P. 1463-1479.
- Izokh N.G., Obut O.T., Iwata K., and Sennikov N.V. [Изох Н.Г., Обут О.Т., Ивата К., Сенников Н.В.] (2003). Conodont associations in the Ordovician of the Altai-Sayan Folded Area // *Bull. of Tomsk State University. Ser. "Earth Sciences" (geology, geography, meteorology, geodesy)*. Supl. 3 (II), April 2003. Problems of geology and geography of Siberia: Contributions of the scientific conference dedicated to 125 anniversary of the TSU and 70<sup>th</sup> anniversary of the Geology-Geography Department, 2-4 April, 2003. P. 88-90. (In Russian). [Ассоциации конодонтов в ордовике Алтае-Саянской складчатой области // *Вестн. Том. гос. ун-та. Серия "Науки о земле"* (геология, география, метеорология, геодезия). Прил. № 3 (II), апрель 2003: Материалы научных конференций, симпозиумов, школ, проводимых в ТГУ. Проблемы геологии и географии Сибири. Материалы науч. конф., посвященной 125-летию основания Томского государственного университета и 70-летию образования геолого-географического факультета, 2-4 апреля 2003 г.]
- Izokh N.G., Sennikov N.V., and Obut O.T. [Изох Н.Г., Сенников Н.В., Обут О.Т.] (2005). Discovery of a new level in the zonal Ordovician conodont scale of the Altai-Sayan Folded area. *V.M. Podobina* (Ed.-in-Chief). *Evolution of Life on the Earth: Proceedings of the III Intern. Sympos.* November 1-3, Tomsk. Tomsk: TSU Publishing House. P. 125-127. (In Russian). [Находка нового уровня в зональной конодонтовой шкале ордовика Алтае-Саянской складчатой области. Эволюция жизни на Земле: Материалы III Междунар. симпоз. Томск: Изд-во Том. гос. ун-та.]
- Kaljo D. and Martma T. (2011). Carbon isotope trend in the Mirny creek area, NE Russia, its specific features and possible implications of the uppermost Ordovician stratigraphy. *Ordovician of the World*. Instituto Geologico y Minero de Espana. Madrid. P. 267-273.
- Kanygin A.V., Yadrenkina A.G., Timokhin A.V., Moskalenko T.A., and Sychev O.V. [Каныгин А.В., Ядренкина А.Г., Тимохин А.В., Москаленко Т.А., Сычев О.В.] (2007). Stratigraphy of oil and gas basins of Siberia. Ordovician of Siberian platform. Novosibirsk: Academic Publ. House "Geo". 403 p. (In Russian). [Стратиграфия нефтегазовых бассейнов Сибири. Ордовик Сибирской платформы. Новосибирск: Академическое изд-во "Гео".]
- Kanygin A.V., Sennikov N.V., Shurygin B.N., Volkova V.C., Grazhdankin D.V., and Postnikov A.A. [Каныгин А.В., Сенников Н.В., Шурыгин Б.Н., Волкова В.С., Гражданкин Д.В., Постников А.А.] (2010). Paleontology. Stratigraphy and paleobiogeography of Precambrian and Phanerozoic of Siberia. In: History of the development of the Institute of Geology and Geophysics of the Siberian Branch (Academy of Sciences of the USSR and the Russian Academy of Sciences) and its scientific schools. Novosibirsk: GEO. P. 101-152. (In Russian). [Палеонтология. Стратиграфия и палеобиогеография докембрия и фанерозоя Сибири // История развития института геологии и геофизики СО (АН СССР и РАН) и его научных направлений. Новосибирск: ГЕО.]
- Kanygin A.V., Yadrenkina A.G., Timokhin A.V., Moskalenko T.A., Gonta T.V., Stepanova N.I., Sychev O.V., and Maslova O.A. [Каныгин А.В., Ядренкина А.Г., Тимохин А.В., Москаленко Т.А., Гонта Т.В., Степанова Н.И., Сычев О.В., Маслова О.А.] (2017). Ordovician regional stratigraphic chart of the Siberian Platform (new version) // *Geology and mineral resources of Siberia*. N 5c. C. 4-57. (In Russian). [Региональная стратиграфическая схема ордовикских отложений Сибирской платформы (новая версия) // Геология и минерально-сырьевые ресурсы Сибири.]
- Koren T.N. [Корень Т.Н.] (2002). Problems of the General stratigraphic chart of Russia // *Regional geology and metallogeny*. N 15. C. 14-25. (In Russian). [Проблемы Общей стратиграфической шкалы ордовикской системы // Региональная геология и металлогения.]
- Krivchikov A.V., Petrunina Z.E., and Severgina L.G. [Кривчиков А.В., Петрунина З.Е., Севергина Л.Г.] (1976). Ordovician System. In: Geological structure and relief of the East Altai: Stratigraphy. Novosibirsk. P. 29-44. (In Russian). [Ордовикская система. Геологическое строение и рельеф Восточного Алтая: Стратиграфия. Новосибирск.]
- Kul'kov N.P. and Severgina L.G. [Кульков Н.П., Севергина Л.Г.] (1989). Stratigraphy and brachiopods of Ordovician and Silurian of the Gorny Altai. Novosibirsk: Nauka. 223 p. (In Russian). [Стратиграфия и брахиоподы ордовика и нижнего силура Горного Алтая. Новосибирск: Наука.]
- Kuznetsov V.A. [Кузнецов В.А.] (1948). Stratigraphic description. Silurian System. Gorny Altai // *Geology of the USSR*, Vol. XIV. West Siberia. Pt. I. Geological description. Moscow; Leningrad: State Publ. House for Geol. Literat. P. 157-165. (In Russian). [Стратиграфическое описание. Силурийская система. Горный Алтай // Геология СССР. Т. XIV. Западная Сибирь. Ч. I. Геологическое описание. М.; Л: Гос. изд. геол. лит.]
- Levitsky E.S. [Левицкий Е.С.] (1962). About a new trilobite genus *Cybelurus* // *Izvestiya vuzov. Geology and exploration*. N 7. P. 129-132. (In Russian). [О новом роде трилобитов – *Cybelurus* // Изв. вузов. Геология и разведка.]
- Levitsky E.S. [Левицкий Е.С.] (1963). Middle Ordovician trilobites from the north-west Gorny Altai and their stratigraphic significance. Abstract of the PhD dissertation. Moscow: Moscow Prospecting Institute. 23 p. (In Russian). [Трилобиты среднего ордовика северо-запада Горного Алтая и их стратиграфическое значение. Автореф. дис.... канд. геол.-мин. наук. М.: Моск. геол.-развед. ин-т.]



- Loydell D.K. (2012). Graptolite biozone correlation charts // Geological Magazine. Vol. 149. P. 124–132.
- Luchinina V.A. [Лучинина В.А.] (1973). Ecology of algae and microstructure of algal biostromes from Lower Cambrian of middle stream of Lena River // Environment and life in geological past. Late Precambrian and Paleozoic of Siberia. Trudy Inst. Geol. Geophys. Issue 169. P. 69–71. (In Russian). [Экология водорослей и микроструктура водорослевых биостромов нижнекембрийского среднего течения р. Лены // Среда и жизнь в геологическом прошлом. Поздний докембрий и палеозой Сибири. Тр. Ин-та геол. и геофиз. СО РАН.]
- Luchinina V.A. and Terleev A.A. [Лучинина В.А., Терлеев А.А.] (2004). Peculiarities in composition of dendrolites in the Lower Cambrian organogenic structures of the Siberian Platform // News of Paleontology and Stratigraphy. Issue 6–7. P. 43–57. Supplement to Journal Geologiya i Geofizika. Vol. 45. (In Russian). [Особенности состава дендролитов в нижнекембрийских органогенных постройках Сибирской платформы // Новости палеонтологии и стратиграфии. Выпуск 6–7. Приложение к журналу Геология и геофизика. Т. 45].
- Mangano M.G. and Droser M.L. (2004). The Ichnologic Record of the Ordovician Radiation. The Great Ordovician Biodiversification Event. Webby B., Paris F., Droser M.L. and Percival I.G. (Eds.) New York: Columbia University Press. P. 369–379.
- Männil R.M. [Мянниль Р.М.] (1966). History of the Baltica basin development during Ordovician. Tallinn: Valgus. 198 p. (In Russian). [История развития Балтийского бассейна в ордовике. Таллин: Изд-во Валгус.]
- Meidla T., Ainsaar L., and Truuver K. (2011). Ostracods in Baltoscandia through the Hirnantian crises. In: Ordovician of the World. Instituto Geologico y Minero de Espana. Madrid. P. 353–357.
- Melnikova L.M. (2010). Some Ostracodes from the Gur'yanovka Formation (Upper Ordovician) of Northeastern Gorny Altai // Paleontological Journal. Vol. 44, N 4. P. 399–408.
- Melnikova L.M. (2011). Ostracodes from the Tuloi and Karasu formations (Ordovician) of northeastern Gorny Altai // Paleontological Journal. Vol. 45, N 1. P. 60–64.
- Miller J.F. (1984). Cambrian and earliest Ordovician conodont evolution, biofacies and provincialism // Geol. Soc. America. Spec. Paper 196. P. 43–68.
- Mitchell C. E., Chen Xu, Bergström S.M., Zhang Yuandong, Wang Zhihao, Webby B.D., and Finney S.C. (1997). Definition of the global boundary stratotype for the Darriwilian Stage of the Ordovician system // Episodes. Vol. 20. P. 158–166.
- Mitchell C.E., Storch P., Holmden C., Melchin V.J., and Guriérrez Varco J.C. (2011). New stable isotope data and fossils the Hirnantian Stage in Bohemia and Spain: implications for correlation and paleoclimate. In: Ordovician of the World. Instituto Geologico y Minero de Espana. Madrid. P. 371–378.
- Moskalenko T.A. [Москаленко Т.А.] (1977). Ashgillian conodonts in the Gorny Altai. Problems of Ordovician and Silurian stratigraphy of Siberia. Novosibirsk: Nauka. Trudy of Inst. Geol. and Geophys. SB AS USSR. Issue 372. P. 74–83. (In Russian). [Ашгиллские конодонты на Горном Алтае. Проблемы стратиграфии ордовика и силура Сибири. Новосибирск: Наука. Труды ИГиГ СО АН СССР.]
- Munnecke A., Calner M., Harper D.T., and Servais T. (2010). Ordovician and Silurian sea-water chemistry, sea level, and climate: A synopsis // Palaeogeography, Palaeoclimatology, Palaeoecology. Vol. 296. P. 389–413.
- Naumenko A.I., Gutak Ya.M. (1982). Correlation of Ordovician deposits in the Ernant and Ulagan troughs // Soviet Geology and Geophysics. Vol. 23, N 4. P. 106–108.
- Nicoll R.S., Laurie J.R., Shergold J.H., and Nielsen A.T. (1992). Preliminary correlation of Latest Cambrian to Early Ordovician sea level events in Australia and Scandinavia. In: Global Perspectives on Ordovician geology. Netherlands. Balkema / Rotterdam / Brookfield. P. 381–394.
- Nielsen A.T. (2003). Ordovician Sea Level Changes: potential for global event stratigraphy // In: Albanesi G.I., Beresi M.S. and Peralta S.H. (Eds). INSUGEO, Serie Correlacion Geologica, 17. Ordovician from the Andes. P. 445–449.
- Nielsen A.T. (2004). Ordovician Sea Level Changes: A Baltoscandian Perspective. In: The Great Ordovician Biodiversification Event. Webby B., Paris F., Droser M.L. and Percival I.G. (Eds). Columbia University Press. New York. P. 84–93.
- Nielsen A.T. (2011). A re-calibrated revised sea level curve for the Ordovician Baltoscandia. In: Ordovician of the World. Instituto Geologico y Minero de Espana. Madrid. P. 399–401.
- Nikonov A.A. [Никонов А. А.] (1931). Sketch on geology and stratigraphy of the Tigerek-Chineta region in the Rudny Altai. Moscow; Leningrad: Geol. Publ. House of the Prospecting Management. Issue 28. P. 31–47. (In Russian). [Очерк геологии и стратиграфии Тигерекско-Чинетинского района в Рудном Алтае. М.; Л.: Геол. изд-во глав, геол.-разв. упр. (Тр. глав. ГРУ ВСНХ СССР. Вып. 28).]
- Noskov Yu.S. [Носков Ю.С.] (2007). Conglomerates with igneous rock pebbles of Gorny Altai. Novosibirsk: SNIIGGiMS Publ. House. 168 p. (In Russian). [Конгломераты с галькой магматических пород Горного Алтая. Новосибирск: Изд-во СНИИГГиМС.]
- Obut A.M. and Sobolevskaya R.F. [Обут А.М., Соболевская Р.Ф.] (1968). Llandoveryan graptolites from cores of drilling wells of Norilsk area. Moscow: Nauka. 136 p. (In Russian). [Граптолиты лландоверии в кернах буровых скважин Норильского района. (В кн. Обут А.М., Соболевская Р.Ф., Меркурьева А.П.). М.: Наука.]
- Obut O.T. and Semenova A.M. (2011). New data on Upper Ordovician radiolarians from the Gorny Altai (SW Siberia, Russia). In: Ordovician of the World. Instituto Geologico y Minero de Espana. Madrid. P. 403–407.
- Obut A.M. and Sennikov N.V. [Обут А.М., Сенников Н.В.] (1984). Graptolites and zonation of the Lower Ordovician of the

- Gorny Altai. Stratigraphy and fauna of Lower Ordovician of the Gorny Altai. Moscow: Nauka. P. 53–106. (In Russian). [Граптолиты и зональное расчленение нижнего ордовика Горного Алтая. Стратиграфия и фауна нижнего ордовика Горного Алтая. Москва: Изд-во Наука.]
- Pärnaste H., Bergström J. (2013).* The asaphid trilobite fauna: Its rise and fall in Baltica // *Palaeogeography, Palaeoclimatology, Palaeoecology*. Vol. 389. P. 64–77.
- Perfil'ev Yu.S. and Levitsky E.S.* [Перфильев Ю.С., Левицкий Е.С.] (1963). Bugryshikha Formation and its trilobites // *Bull. Mosc. Soc. Nature*. Vol. 38. Issue 2. P. 17–29. [Бугрышихинская свита и ее трилобиты // Бюл. Моск. об-ва испыт. природы.]
- Perfil'ev Yu.S.* [Перфильев Ю.С.] (1959). New data on Lower Paleozoic stratigraphy of the Gorny Altai // *Izvestiya of the High School. Geology and Prospecting*. N 11. P. 20–30. (In Russian). [Новые данные по стратиграфии нижнего палеозоя Горного Алтая // Изв. вузов. Геология и разведка.]
- Petrulina Z.E.* [Петрунина З.Е.] (1966). Tremadocian trilobites and biostratigraphy from the western part of Sayan-Altai mountainous area. Abstract of PhD dissertation. Alma-Ata: Institute of Geological Sciences AS Kazakh SSR. 30 p. (In Russian). [Трилобиты и биостратиграфия тремадока западной части Саяно-Алтайской горной области. Автореф. канд. дисс. Алма-Ата: Изд-во Ин-та геологических наук АН Казахской ССР.]
- Petrulina Z.E.* [Петрунина З.Е.] (1968). About biostratigraphic division of Tremadocian of the Western part of the Sayan-Altai mountainous area // *New data on geology and mineral resources of the West Siberia*. Tomsk. Issue 3. P. 87–89. (In Russian). [О биостратиграфическом расчленении тремадока западной части Саяно-Алтайской горной области // Новые данные по геологии и полезным ископаемым Западной Сибири. Томск.]
- Petrulina Z.E.* [Петрунина З.Е.] (1973). New Tremadocian trilobites genera and species from West Siberia // *New data on geology and mineral resources of the West Siberia*. Tomsk. Issue 8. P. 59–68. (In Russian). [Новые роды и виды тремадоковских трилобитов Западной Сибири // Новые данные по геологии и полезным ископаемым Западной Сибири. Томск.]
- Petrulina Z.E.* [Петрунина З.Е.] (1990). Some new Early Ordovician trilobites from the western part of the Altai- Sayan Mountainous Area // *New in Paleozoic paleontology and biostratigraphy of the Asian part of the USSR*. Novosibirsk: Nauka, Siberian Branch. Trudy of Inst. Geol. and Geophys. SB AS USSR. Issue 770. P. 21–58. (In Russian). [Некоторые новые раннеордовикские трилобиты западной части Алтае-Саянской горной области. Новое в палеонтологии и биостратиграфии палеозоя Азиатской части СССР. Новосибирск: Наука, Сиб. отд-ние. (Труды Ин-та геологии, геофизики СО АН СССР.)]
- Petrulina Z.E. and Severgina L.G.* [Петрунина З.Е., Севергина Л.Г.] (1962). On Ordovician stratigraphy of the North-Western Gorny Altai // *Materials on geology of the West Siberia // New data on paleontology and stratigraphy of the West Siberia*. Tomsk: Tomsk University Press. Issue 63. P. 81–93. (In Russian). [К стратиграфии ордовика северо-запада Горного Алтая // Материалы по геологии Западной Сибири. Новые данные по палеонтологии и стратиграфии Западной Сибири. Томск: Изд-во Том. гос. ун-та. (Тр. Том. гос. ун-та. Вып. 63).]
- Petrulina Z.E., Sennikov N.V., Ermikov V.D., Zeyfert L.L., Krivchikov A.V., Puzyrev A.A., Severgina L.G., Obut A.M., and Zaslavskaya N.M.* [Петрунина З.Е., Сенников Н.В., Ермиков В.Д., Зейферт Л.Л., Кривчиков А.В., Пузырев А.А., Севергина Л.Г., Обут А.М., Заславская Н.М.] (1984). Lower Ordovician stratigraphy and fauna of Mountain Altai. Moscow: Nauka Publ. House. 124 p. (In Russian). [Стратиграфия и фауна нижнего ордовика Горного Алтая. М.: Наука.]
- Resolutions of Interdepartmental Stratigraphic committee and its regular commissions. (2012).* St. Petersburg: VSEGEI Publ. House. Vol. 41. 44 p. (In Russian). [Постановления Межведомственного стратиграфического комитета и его постоянных комиссий. Санкт-Петербург: Изд-во ВСЕГЕИ.]
- Safonova I.Yu., Sennikov N.V., Komiya T., Bychkova Y.V., and Kurganskaya E.V. (2011).* Geochemical diversity in oceanic basalts hosted by the Zashur'ya accretionary complex, NW Russian Altai, Central Asia: Implications from trace elements and Nd isotopes // *Journal of Asian Earth Sciences*. Vol. 42. P. 191–207.
- Schonlaub H.P., Ferretti A., Gaggero L., Hammarlund E., Harper D.A.T., Histon K., Priewalder H., Spotl C., and Storch P. (2011).* The Late Ordovician glacial event in the Carnic Alps (Austria). In: *Ordovician of the World*. Instituto Geológico y Minero de Espana. Madrid. P. 515–526.
- Sennikov N.V.* [Сенников Н.В.] (1976). Graptolites and Lower Silurian stratigraphy of the Gorny Altai. Moscow: Nauka Publ. House. 270 p. (In Russian) [Граптолиты и стратиграфия нижнего силура Горного Алтая. М.: Наука.]
- Sennikov N.V.* [Сенников Н.В.] (1996). Paleozoic graptolites from the Middle Siberia (systematics, phylogeny, biochronology, biology, paleogeography). Novosibirsk: SB RAS, SRC UIGGM Press. 225 p. (In Russian). [Граптолиты палеозоя Средней Сибири (систематика, филогения, биохронология, биологическая природа, палеозоогеография). Новосибирск: Изд-во СО РАН, НИЦ ОИГГМ.]
- Sennikov N.V.* [Сенников Н.В.] (1998). Manifestation of the global Ordovician-Silurian biotic crisis in the Central Siberian graptolite communities // *Russian Geology and Geophysics*. Vol. 39, N5. P. 565–574.
- Sennikov N.V.* [Сенников Н.В.] (2006a). Geological events in Paleozoic of the Altai-Sayan Folded Area and their reflection in paleogeographic and sedimentary environments and paleobiota. In: *Evolution of the biosphere and biodiversity*. 70th anniversary A.Yu. Rosanov. Moscow: KMK Publ. House. P. 506–526. (In Russian). [Геологические события в палео-



зое Алтае-Саянской складчатой области и их отражение в палеогеографических и седиментационных обстановках и в палеобиотах. Эволюция биосферы и биоразнообразия. К 70-летию А.Ю. Розанова. М.: Товарищество научных изданий КМК.]

- Sennikov N.V. [Сенников Н.В.] (2006b). Ordovician events and palaeogeographic sedimentary environments in the Altai-Salair paleobasin (Western part of the Altai-Sayan Folded Area, Russia). Intern. Sympos. "Palaeogeography and Global Correlation of Ordovician Events". (IGCP 503 Project). Contributions. Novosibirsk, Russia. August 5-7, 2006. Novosibirsk: Publishing House of SB RAS, "Geo" Branch. P. 50–52.
- Sennikov N.V. [Сенников Н.В.] (2012). New data on graptolite zonation of Katian and Hirnantian (Ordovician) of Gorny Altai. Paleontology and stratigraphic boundaries. Proceedings of LVIII session of Paleontological society of RAS (2–6 April, 2012, St. Petersburg). St. Petersburg: VPO. P. 122–123. (In Russian). [Новые данные по зональному расчленению катинского и хирнантского ярусов (ордовик) Горного Алтая по граптолитам. Палеонтология и стратиграфические границы. Мат. LVIII сессии Палеонтол. общества при РАН (2–6 апреля 2012 г., Санкт-Петербург). СПб.: ВПО.]
- Sennikov N.V. [Сенников Н.В.] (2013). Graptolite zones in Ordovician of Russia. V.I. Krasnov (Ed.) Regional stratigraphy of Late Precambrian and Paleozoic of Siberia: digest of scientific papers Novosibirsk: SNIIGiMS Publ. House. P. 45–58. (In Russian). [Зональные граптолитовые подразделения в ордовике России. Региональная стратиграфия позднего докембрия и палеозоя Сибири: Сб. науч. тр. / Под ред. В.И.Краснова. Новосибирск: СНИИГТиМС.]
- Sennikov N.V. and Obut O.T. [Сенников Н.В., Обут О.Т.] (2002). Testing the chitinozoan zonal succession according to the graptolite zonal scale (Ordovician, Gorny Altai and Moscow syncline) // News of Paleontology and Stratigraphy. Issue 5. P. 51–63. Supplement to Journal Geologiya i Geofizika. Vol. 43. (In Russian). [Тестирование зональной последовательности по хитинозоям на базе граптолитовых зон (ордовик, Горный Алтай и Московская синеклиза) // Новости палеонтологии и стратиграфии. Выпуск 5. Приложение к журналу Геология и геофизика. Том 43.]
- Sennikov N.V. and Bukolova E.V. [Сенников Н.В., Буколова Е.В.] (2010). Dynamics of graptolite taxonomic diversity in Altai and Salair Ordovician basin. Evolution of organic world and biotic crisis. LVI session of Paleontological society. St. Petersburg: Paleontological institute Publ. House. P. 100–102. (In Russian). [Динамика таксономического разнообразия граптолитов в Алтайском и Салаирском ордовикских бассейнах. Эволюция органического мира и биотические кризисы. LVI сессия Палеонтологического общества. Санкт-Петербург: Изд-во ПИН.]
- Sennikov N.V. and Ainsaar L. [Сенников Н.В., Айнсаар Л.] (2012). The first data on carbon isotopes in Hirnantian sediments of Gorny Altai. A.I. Zhamoida (Ed-in-chief). Paleozoic of Russia: regional stratigraphy, paleontology, geo- and bioevents. Proceedings of III All-Russian meeting: Upper Paleozoic of Russia: regional stratigraphy, paleontology, geo- and bioevents. St. Petersburg: VSEGEI. P. 205–207. (In Russian). [Первые данные по изотопам углерода в отложениях хирнантского яруса Горного Алтая. Палеозой России: региональная стратиграфия, палеонтология, гео- и биособытия. Материалы III Всероссийского совещания «Верхний палеозой России: региональная стратиграфия, палеонтология, гео- и биособытия», 24–28 сентября 2012 г., Санкт-Петербург / Отв. ред. А.И. Жамойда. СПб.: Изд-во ВСЕГЕИ.]
- Sennikov N.V. and Tolmacheva T.Yu. [Сенников Н.В., Толмачева Т.Ю.] (2013). Minutes of the meeting of the Ordovician and Silurian Systems Commission. Resolutions of the Interdepartmental Stratigraphic Committee and its standing committees. St. Petersburg: VSEGEI Publ. House. Issue 42. P. 19–20. (In Russian). [Протокол заседания Комиссии по ордовикской и силурийской системам. Постановления Межведомственного стратиграфического комитета и его постоянных комиссий. Санкт-Петербург: Изд-во ВСЕГЕИ.]
- Sennikov N.V., Puzyrev A.A., and Russkikh V.G. [Сенников Н.В., Пузырев А.А., Русских В.Г.] (1979). Ordovician and Silurian of the Ust'-Chagyrka Village area. Problems of stratigraphy and tectonics of Siberia. (Trudy Inst. Geol. Geophys. SB AS USSR). Novosibirsk. P. 30–45. (In Russian). [Ордовик и нижний силур района с.Усть-Чагырки (Горный Алтай). Проблемы стратиграфии и тектоники Сибири. (Труды Ин-та геологии и геофизики СО АН СССР. Новосибирск.)
- Sennikov N.V., Ermikov V.D., Petrunina Z.E., Puzyrev A.A., and Severgina L.G. (1982). On the age of the basal strata of the Ordovician to Middle Devonian complex in the Northwestern Altai region // Soviet Geology and Geophysics. Vol. 23, N 8. P. 51–56.
- Sennikov N.V., Petrunina Z.E., Gladkikh L.A., Ermikov V.D., Zinov'eva T.V., Mamlin A.N., and Shokal'skii S.P. (1984). New Ordovician-Silurian boundary cross-sections in Gorny Altai // Soviet Geology and Geophysics. Vol. 25, N 7. P. 18–22.
- Sennikov N.V., Petrunina Z.E., and Savitsky V.R. [Сенников Н.В., Петрунина З.Е., Савицкий В.Р.] (1995). About Ordovician Savel'evka and Bulukhta formations of the Gorny Altai. In: New data on geology and natural resources from the western part of the Altai-Sayan Area. Novokuznetsk: GGP "Zapsibgeologia". P. 63–66. (In Russian). [О савельевской и булуктинской свитах ордовика Горного Алтая. Новые данные о геологии и полезных ископаемых западной части Алтае-Саянской области. Новокузнецк: ГГП "Запсибгеология".]
- Sennikov N.V., Kazansky A.Yu., Ermikov V.D., Iwata K., Mikhaltsov N.E., Obut O.T., Izokh N.G., and Khlebnikova T.V. [Сенников Н.В., Казанский А.Ю., Ермаков В.Д., Ивата К., Михальцов Н.Э., Обут О.Т., Изох Н.Г., Хлебникова Т.В.] (2000). Comparative analysis of sedimentary, paleozoogeographic and paleoclimatic data on Ordovician and Silurian of Altai and Tuva. 300 years of the Russian prospecting geological survey: history of the prospecting, geological structure

- and natural resources of Altai. Barnaul: Altai State University Publ. House. P. 195–200. (In Russian). [Сравнительный анализ седиментационных, палеозоогеографических и палеомагнитных данных по ордовику и силуру Алтая и Тувы. 300 лет горно-геологической службе России: история горнорудного дела, геологическое строение и полезные ископаемые Алтая. Барнаул: Изд-во Алтайского гос. ун-та.]
- Sennikov N.V., Petrunina Z.E., and Gladkikh L.A. [Сенников Н.В., Петрунина З.Е., Гладких Л.А.] (2001a). Litho- and biostratigraphical dissection of the Ashgillian deposits in the central part of Gorny Altai. *In: Actual problems of geology and mineragenesis Southern of Siberia*. Novosibirsk: “Geo” branch of SB RAS Publ. House. P. 135–144. (In Russian). [Лито- и биостратиграфическое расчленение ашгилла центральной части Горного Алтая. Актуальные вопросы геологии и минерагении юга Сибири. Новосибирск: Изд-во СО РАН, филиал “Гео”.]
- Sennikov N.V., Obut O.T., Khlebnikova T.V., Iwata K., and Ermikov V.D. [Сенников Н.В., Обут О.Т., Хлебникова Т.В., Ивата К., Ермиков В.Д.] (2001b). Structure and age of the volcanic-terrigenous Zasurya Formation from the central part of the Gorny Altai (Anui-Chuya zone, Marcheta River). *In: Actual problems of geology and mineragenesis Southern of Siberia. Materials of the scientific conference, 31 Oct. – 2 Nov. 2001, Elan Village, Novokuznetsk, Kemerovo region*. Novosibirsk: “Geo” branch of SB RAS Publ. House. P. 145–151. (In Russian). [Строение и возраст вулканогенно-кремнисто-терригенной засурьинской свиты в центральной части Горного Алтая (Ануйско-Чуйская зона, р. Марчета). Актуальные вопросы геологии и минерагении юга Сибири. Новосибирск: Изд-во СО РАН, филиал “Гео”.]
- Sennikov N.V., Iwata K., Ermikov V.D., Obut O.T., and Khlebnikova T.V. (2003). Palaeozoic oceanic sedimentation settings and fauna associations at the southern margin of the West Siberian Plate // *Russian Geology and Geophysics*. Vol. 44, N 1–2. P. 152–168.
- Sennikov N.V., Obut O.T., Iwata K., Khlebnikova T.V., and Ermikov V.D. (2004). Lithological Markers and Bio-indicators of Deep-water Environments during Paleozoic Siliceous Sedimentation (Gorny Altai Segment of the Paleo-Asian Ocean) // *Gondwana Research*. Vol. 7. N 3. P. 843–852.
- Sennikov N.V., Izokh N.G., Kazansky A.Yu., Petrunina Z.E., Kungurtsev L.V., Khlebnikova T.V., Mikhail'sov N.E., and Savitsky V.R. [Сенников Н.В., Изох Н.Г., Казанский А.Ю., Петрунина З.Е., Кунгурцев Л.В., Хлебникова Т.В., Михальцов Н.Э., Савицкий В.Р.] (2006a). New biostratigraphic and palaeomagnetic data on the Malinovka Series (Lower and Middle Ordovician, Tuva) // *News of Paleontology and Stratigraphy*. Issue 8. P. 27–41. Supplement to *Journal Geologiya i Geofizika*. Vol. 47. (In Russian). [Новые биостратиграфические и палеомагнитные данные по малиновской серии (нижний-средний ордовик, Тува) // *Новости палеонтологии и стратиграфии*. Вып. 8. Приложение к журналу «Геология и геофизика». Т. 47.]
- Sennikov N.V., Petrunina Z.E., Obut O.T., and Izokh N.G. (2006b). Ordovician Regional stage and zone of the western part of Altai-Sayan Folded Area (south of West Siberia) – as a tool for Regional and Global correlation of the Ordovician events. “Palaeogeography and Global Correlation of Ordovician Events” (IGCP 503 Project “Ordovician Palaeogeography and Palaeoclimate”): *Contrib. of Intern. Sympos.* Novosibirsk, Aug. 5–7, 2006. Novosibirsk: Academic Publishing House “Geo”. P. 57–60.
- Sennikov N.V., Yolkin E.A., Petrunina Z.E., Gladkikh L.A., Obut O.T., Izokh N.G., and Kipriyanova T.P. (2008). Ordovician-Silurian Biostratigraphy and Paleogeography of the Gorny Altai. Novosibirsk: Publishing House SB RAS. 156 p.
- Sennikov N.V., Obut O.T., and Bukolova E.V. (2011a). Ordovician regional chronostratigraphic scheme of the Gorny Altai / *In: Ordovician of the World. Gutierrez-Marco J.C., Rabano I., and Garcia-Bellido D.* (Eds). *Cuadernos del Museo Geominero*. N 14. Instituto Geologico y Minero de Espana, Madrid. P. 547–551.
- Sennikov N.V., Obut O.T., Bukolova E.V., and Tolmacheva T.Yu. (2011b). The depths of the Early Paleozoic sedimentary basins of the Paleasian Ocean: lithofacies and bioindicator estimates // *Russian Geology and Geophysics*. Vol. 52. Issue 10. P. 1171–1194.
- Sennikov N.V., Tolmacheva T.Yu., Izokh N.G., and Obut O.T. (2014). The Cambrian-Ordovician Boundary in Gorny Altai // *Doklady Earth Sciences* Vol. 457, N 1. P. 792–795.
- Sennikov N.V., Tolmacheva T.Yu., Obut O.T., Izokh N.G., and Lykova E.V. (2015a). Zonation of the Siberian Ordovician deposits based on pelagic groups of fauna // *Russian Geology and Geophysics*. Vol. 56. Issue 4. P. 594–610.
- Sennikov N.V., Ainsaar L., and Meidla T. (2015b). Hirnantian Isotope Carbon Excursion in Gorny Altai, southwestern Siberia // *Estonian Journal of Earth Sciences*. Vol. 64, N 3. p. 225–232.
- Sennikov N.V., Luchinina V.A., Obut O.T., Tokarev D.A., and Khabibulina R.A. (2017). Late Ordovician “coral-microbial-algal meadows” in the north-western part of Gorny Altai. *Contributions of Interekspo Geo-Siberia-2017: XIII International Conference “Subsurface management. Mining. New trends and techniques for prospecting, exploration and exploitation of mineral resources. Economy. Geoeology”*. – Novosibirsk: SSUGT, P.150–154. [Позднеордовикские «кораллово-микробально-водорослевые луга» в северо-западной части Горного Алтая. Интерэкспо ГЕО-Сибирь-2017. XIII Междунар. науч. конгр. и выставка, 17–21 апреля 2017 г., Новосибирск: Междунар. науч. конф. «Недропользование. Горное дело. Направления и технологии поиска, разведки и разработки месторождений полезных ископаемых. Экономика. Геоэкология»: сб. материалов в 4 т. Т 2, № 1. – Новосибирск: СГУГиТ.]
- Sennikov N.V., Obut O.T., Tolmacheva T.Yu., Lykova E.V., and Khabibulina R.A. (2018a). The Upper Ordovician of northeastern Gorny Altai: stratigraphy and deposition environments // *Russian Geology and Geophysics*. Vol. 59. Issue 1. P. 72–87.



- Sennikov N.V., Obut O.T., Izokh N.G., Kipriyanova T.P., Lykova E.V., Tolmacheva T.Yu., Khabibulina R.A. [Сенников Н.В., Обут О.Т., Изох Н.Г., Киприянова Т.П., Лыкова Е.В., Толмачева Т.Ю., Хабибулина Р.А.] (2018 b). Regional stratigraphical chart of the Ordovician of western part of the Altai-Sayan Folded Area (new version) // *Geology and mineral resources of Siberia*. N 7. P. 15-53. (In Russian). [Региональная стратиграфическая схема ордовикских отложений западной части Алтае-Саянской складчатой области (новая версия) // *Геология и минерально-сырьевые ресурсы Сибири*.]
- Sennikov V.M. [Сенников В.М.] (1962). Ordovician of the Uimen'-Lebed' Synclinorium of the Gorny Altai. Materials on regional geology of Siberia. Moscow: Gostekhgeolizdat. Issue 24. P. 116–133. (In Russian). [Ордовик Уйменско-Лебедского синклиория Горного Алтая. Материалы по региональной геологии Сибири. М.: Госгеолтехиздат. (Тр. СНИИГГиМС. Вып. 24).]
- Sennikov V.M. [Сенников В.М.] (1977). The history of development of structures in the south part of Altai-Sayan folded area in Ordovician. Barnaul: Altai Publ. House. 135 p. (Trudy SNIIGGiMS. Issue 201.) (In Russian). [История развития структур южной части Алтае-Саянской складчатой области в ордовике. Барнаул: Алтайское книжное изд-во. (Тр. СНИИГГиМС, вып. 201).]
- Sennikov V.M. and Sennikov N.V. (1982). The stratigraphy of Ordovician rocks in the Anui-Chuya synclinorium (Gornyi Altai region) // *Soviet Geology and Geophysics*. Vol. 23, N. 6. P. 13–20.
- Sennikov V.M., Vinkman M.K., and Kononov A.N. [Сенников В.М., Винкман М.К., Кононов А.Н.] (1959). Cambrian-Ordovician and Ordovician of the Gorny Altai. In: Materials on regional geology. Moscow: Gosgeoltekhizdat. P. 51–66. (In Russian). [Кембро-ордовик и ордовик Горного Алтая. Материалы по региональной геологии. М.: Госгеолтехиздат.]
- Severgina L.G. [Севергина Л.Г.] (1965). Brachiopods and Ordovician biostratigraphy of the western part of the Sayan-Altai Mountainous Area: Abstract of PhD dissertation. Tomsk: Tom. University Press. 23 p. (In Russian). [Брахииподы и биостратиграфия ордовика западной части Саяно-Алтайской Горной области: Автореф. дис.... канд. геол.-мин. наук. Томск: Изд-во Том. гос. ун-та.]
- Severgina L.G. [Севергина Л.Г.] (1968). Ordovician System. Altai-Sayan Area (western part). In: Geological structure of the USSR. Vol. I. Moscow: Nedra Press. P. 278–280. (In Russian). [Ордовикская система. Алтае-Саянская область (западная часть). Геологическое строение СССР. Т. I. М.: Недра.]
- Severgina L.G. [Севергина Л.Г.] (1972). Stratigraphic chart for the Middle and Upper Ordovician of the western Part of Sayan-Altai Area. Kemerovo: Kemerovo Press. P. 62–66. (In Russian). [Схема стратиграфии средне- и верхнеордовикских отложений западной части Саяно-Алтайской области. Кемерово: Кемеровское кн. изд-во. (Изв. Кузнецкого отдела Географического об-ва СССР. Вып. 1).]
- Severgina L.G. [Севергина Л.Г.] (1973). Biostratigraphy of the Ordovician deposits of the Old Paleozoic framework of the Kuznetsk Basin. In: Nature of Kuznetsk Basin. Novokuznetsk: USSR Kuznetsk Geographic Society Press. P. 170–181. (In Russian). [Биостратиграфия ордовикских отложений древнепалеозойского обрамления Кузбасса. Природа Кузбасса. Новокузнецк: Изд-во Кузнецкого отдела Географ. об-ва СССР.]
- Severgina L.G. [Севергина Л.Г.] (1978). Brachiopods and biostratigraphy of Upper Ordovician of Gorny Altai, Salair and Gornaya Shoriya. In: Fauna and biostratigraphy of Upper Ordovician and Silurian of Altai-Sayan area. Moscow: Nauka. P. 3–41. (In Russian). [Брахииподы и биостратиграфия верхнего ордовика Горного Алтая, Салаира и Горной Шории. Фауна и биостратиграфия верхнего ордовика и силура Алтае-Саянской области. М.: Наука.]
- Severgina L.G. [Севергина Л.Г.] (1984). Some Upper Ordovician (Ashgillian) brachiopods of Gorny Altai. In: Paleozoic paleontology and biostratigraphy of Siberia. Novosibirsk: Nauka. P. 39–48. (In Russian). [Некоторые верхнеордовикские (ашгиллские) брахиподы Горного Алтая. Палеонтология и биостратиграфия палеозоя Сибири. Новосибирск. Изд-во Наука.]
- Strachan I. (1996). A bibliographic index of British graptolites (Graptoloidea). Part 1. The Palaeontographical Society London. P. 1–40.
- Strachan I. (1997). A bibliographic index of British graptolites (Graptoloidea). Part 2. The Palaeontographical Society London. P. 41–155.
- Stratigraphic dictionary of the USSR. (1956). Moscow: Gosgeoltekhizdat. 1283 p. (In Russian). [Стратиграфический словарь СССР. М.: Госгеолтехиздат.]
- Stratigraphic dictionary of the USSR. Cambrian, Ordovician, Silurian, Devonian. (1975). Leningrad: Nedra Press. 622 p. (In Russian). [Стратиграфический словарь СССР. Кембрий, ордовик, силур, девон. Л.: Недра.]
- Stratigraphic dictionary of the USSR. New Paleozoic stratigraphic units for the USSR. (1991). Leningrad: Nedra Press. 555 p. (In Russian). [Стратиграфический словарь СССР. Новые стратиграфические подразделения палеозоя СССР. Л.: Недра.]
- Paleozoic stratigraphy of the Central Siberia (1967). Novosibirsk: Nauka. 251 p. (In Russian). [Стратиграфия палеозоя Средней Сибири. Новосибирск: Изд-во Наука.]
- The Geologic Time Scale 2012. (2012). / Editors: Gradstein F.M., Ogg J.G., Schmitz M.D., Ogg G.M. 2-volume set. Elsevier Science Ltd. 1176 p.
- Tikhonov V.I. [Тихонов В.И.] (1956). About stratigraphy of the “Anui-Chuya” and “Green-violet” formations of the Charysh

- Altai // Trudy VAGT. Issue 2. P. 34–45. (In Russian). [К стратиграфии “Ануйско-Чуйской” и “Зелено-фиолетовой” формаций Причарышского Алтая // Труды ВАГТ.]
- Underwood C.J., Crowley S.F., Marshall J.D., and Brenchley P.J. (1997). High-resolution carbon isotope stratigraphy of the basal Stratotype (Dob’s Linn, Scotland) and its global correlation // Journal of the Geological Society. Vol. 154. P. 709–718.
- Usov M.A. [Усов М.А.] (1936). Stages and cycles of tectonogenesis in the West-Siberian region. Tomsk, 209 p. (In Russian). [Фазы и циклы тектоногенеза Западно-Сибирского края. Томск.]
- Varaksina I.V. and Sennikov N.V. [Вараксина И.В., Сеников Н.В.] (2006). Lithology and carbonate rocks sedimentary environments in Ordovician of Gorny Altai and Salair (Siberia, Russia). International Symposium “Palaeogeography and Global Correlation of Ordovician Events”. (IGCP 503 Project). Contributions. Novosibirsk, Russia. August 5–7, 2006. Novosibirsk: Publishing House of SB RAS, “Geo” Branch. P. 67–71.
- Vinkman M.K. and Gintsinger A.B. [Винкман М.К., Гинцингер А.Б.] (1967). Type section of the Ordovician deposits in the north-east of the Anui-Chuya Synclinorium, Gorny Altai. Materials on the Siberian regional geology. Novosibirsk: SNIIGGiMS Publ. House. Issue 57. P. 161–167. (In Russian). [Типовой разрез ордовикских отложений на северо-востоке Ануйско-Чуйского синклинория в Горном Алтае. Материалы по региональной геологии Сибири. Новосибирск: Изд-во СНИИГГиМС. (Тр. СНИИГГиМС. Вып. 57).]
- Walliser O.H. (1986). Towards a more critical approach to bio-events. Global Bio-Events. A Critical Approach. Walliser O.H. (Ed.). Lecture Notes in Earth Sciences. Bhattacharji S., Friedman G.M., Neugebauer H.J. and Seilacher A. (Eds.). Vol. 8. Springer-Verlag. P. 5–16.
- Walliser O.H. (Ed.) (1996). Global Events and Event Stratigraphy in the Phanerozoic. Springer-Verlag, New York, 333 p.
- Wang Xiaofeng and Chen Xu. (2003). Ordovician chitinozoan diversification events in China // Science in China (Ser. D. Earth Sciences). Vol. 33, N 3. P. 210–215. (In Chinese).
- Wang Xiaofeng, Stouge S., Erdtmann B.-D., Chen Xiaohong, Li Zhihong, Wang Chuanshang, Zeng Qinglun, Zhou Zhiqiang, and Chen Huiming. (2005). A proposed GSSP for the base of the Middle Ordovician Series: the Huanghuachang section, Yichang, China // Episodes. Vol. 28, N 2. P. 105–117.
- Webby B., Cooper R., Bergström S.M., and Paris F. (2004). Stratigraphic Framework and Time Scales. In: The Great Ordovician Biodiversification Event. Webby B., Paris F., Droser M.L., and Percival I.G. (Eds). New York: Columbia University Press. P. 41–47.
- Yaroshinskaya A.M. [Ярошинская А.М.] (1960). Bryozoa. Paleozoic biostratigraphy of the Sayan-Altai mountainous area. Vol. I. Lower Paleozoic. L.L. Khalfin (Ed.). Novosibirsk: SNIIGGiMS Publ. House. Trudy SNIIGGiMS. Issue 19. P. 362, 393–400. [Тип Bryozoa. Мшанки. Биостратиграфия палеозоя Саяно-Алтайской горной области. Т. I. Нижний палеозой. Под ред. Л.Л. Халфина. Труды СНИИГГиМС. Вып. 19. Новосибирск: СНИИГГиМС.]
- Yolkin E.A., Sennikov N.V., Buslov M.M., Yazikov A.Yu., Gratsianova R.T., and Bakharev N.K. (1994). Paleogeographic reconstruction of the Western Altai-Sayan area in the Ordovician, Silurian, and Devonian and their geodynamic interpretation // Russian Geology and Geophysics. Vol. 35, N 7–8. P. 100–124.
- Zalasiewicz J.A., Taylor L., Rushton A.W.A., Loydell D.K., Rickards R.B., and Williams M. (2009). Graptolites in British Stratigraphy // Geological Magazine. Vol. 146, N 06. P. 785–850.



## PALEONTOLOGICAL PLATES

## Tabulate corals

### Plate 1

Tabulate corals from the localities in the northeastern part of Gorny Altai, the right bank of the Lebed' River, Lebed' Section.

*Collection of Raliya A. Khabibulina.*

Fig. 1. *Heliolites tchorparensis* Poltavceva:

The Lebed' section, Member 19, loc. Kh1013-2. Upper Ordovician, Chebor Formation.

1a – transverse thin section, 1b – longitudinal thin section.

Fig. 2. *Nyctopora nicholsoni* (Raduguin):

The Lebed' section, Member 7, loc. Kh1014-2. Upper Ordovician, Gur'yanovka Formation.

2a – transverse thin section, 2b – longitudinal thin section.

Fig. 3. *Calapoecia canadensis* Billings:

The Lebed' section, Member 9, loc. Kh1014-4. Upper Ordovician, Gur'yanovka Formation.








3a – transverse thin section, 3b – longitudinal thin section.

Fig. 4. *Tetradium* sp.:

The Lebed' section, Member 9, loc. Kh1214-9. Upper Ordovician. Gur'yanovka Formation.

4a – transverse thin section, 4b – longitudinal thin section.





## Tabulate corals

### Plate 2

Tabulate corals from the localities in the eastern part of Gorny Altai, Teletskoye Lakeside, the right bank of the Verkhniy Turochak River, left tributary of Iogach River.

*Collection of Raliya A. Khabibulina.*

Fig. 1. *Catenipora* sp.:

The Verkhniy Turochak Section, Member 4, loc. S-183. Upper Ordovician, Samysh Body.

1a – transverse thin section, 1b – longitudinal thin section.

Fig. 2. *Rhaphidophyllum ellipsoidalis* Preobrazhensky:

The Verkhniy Turochak Section, loc. 6116. Upper Ordovician, Samysh Body.

2a – transverse thin section, 2b – longitudinal thin section.

Fig. 3. *Nyctopora denticulate* Sokolov et Tesakov:

The Verkhniy Turochak Section, Member 4, loc. S-1754. Upper Ordovician, Samysh Body.





3a – transverse thin section, 3b – longitudinal thin section.

Fig. 4. *Cyrtophyllum bargastensis* Dziubo:

The Verkhniy Turochak Section, Member 4, loc. S-183. Upper Ordovician, Samysh Body.

4a – transverse thin section, 4b – longitudinal thin section





## Tabulate corals

### Plate 3

Tabulate corals from the localities in the eastern part of Gorny Altai, Teletskoye Lakeside (Verkhniy Turochak River), Biya and Lebed' rivers and northwestern part of Gorny Altai (Charysh River).

*Collection of Raliya A. Khabibulina.*

Fig. 1. *Reuschia aperta* Kiaer:

The right bank of Verkhniy Turochak River (left tributary of Iogach River). The Verkhniy Turochak Section, loc. 6116. Upper Ordovician, Samysh Body.

1a – transverse thin section, 1b – longitudinal thin section

Fig. 2. *Tetradium borealis* Tchernychev:

The right bank of the Biya River, loc. 7134. Upper Ordovician.

2a – transverse thin section, 2b – transverse thin section.

Fig. 3. *Mesofavosites* sp.:

The right bank of the Charysh River (near Ust'-Kan Village), loc. S-1615. Upper Ordovician- Lower Silurian.





3a – transverse thin section, 3b –longitudinal thin section.

Fig. 4. *Reuschia* sp.:

The right bank of the Lebed' River. Lebed' Section, Member 9, loc. Kh1214-13. Upper Ordovician, Gur'yanovka Formation.

4a – transverse thin section, 4b – longitudinal thin section.





## Brachiopods

### Plate 4

Brachiopods from the eastern part of Gorny Altai, Teletskoye Lakeside, the Tozodov Section. Tozodov Formation, Upper Ordovician, Sandbian Stage.

*Collection of Tatyana A. Shcherbanenko.*

Fig. 1. *Eoanastrophia lebediensis* (Severgina):

1 – specimen Tozodov-6/1, ventral exterior, natural-sized; 1a – ventral exterior; 1b – dorsal exterior; 1c – lateral view; 1d – anterior view; 1e – posterior view; all  $\times 3$ . The Tozodov Section, Member 6.

Figs 2-6. *Apatomorpha altaica* Severgina:

2 – specimen C-1440/2, ventral exterior, natural-sized. 2a – ventral exterior; 2b – dorsal exterior; 2c – lateral view; 2d – posterior view; all  $\times 3$ . The Tozodov Section, Member 3.

3 – specimen C-1440/3, ventral exterior, natural-sized. 3a – ventral exterior; 3b – dorsal exterior; 3c – lateral view; 3d – posterior view; all  $\times 2$ . The Tozodov Section, Member 3.

4 – specimen C-1441/4, internal mold of ventral valve,  $\times 3$ . The Tozodov Section, Member 7.

5 – specimen C-1442/5, internal mold of dorsal valve,  $\times 3$ . The Tozodov Section, Member 8.

6 – specimen C-1442/6, cardinal process,  $\times 3$ . The Tozodov Section, Member 8.


Figs 7-9. *Sivorthis friendsvillensis* (Cooper):

7 – specimen C-1440/7, ventral exterior, natural-sized. 7a – ventral exterior; 7b – dorsal exterior; 7c – lateral view; 7d – anterior view; 7e – posterior view; all  $\times 3$ . Tozodov Section, Member 3.

8 – specimen C-1442/8, internal mold of ventral valve  $\times 3$ . Tozodov Section, Member 8.

9 – specimen C-1442/9, internal mold of dorsal valve,  $\times 3$ . Tozodov Section, Member 8.





## Trilobites

### Plate 5

Trilobites from the localities in the northwestern Gorny Altai, Bugryshikha Formation, the Gora Altai Section. Specimens 1–6 are from Member 5, in 1 m from the bottom; specimens 7–11 are from Member 14, in 2 m from the bottom.

*Collection of Aleksander V. Timokhin.*

Figs 1, 4, 8 – *Homotelus inferus* Levitsky.


Figs 2, 5, 7, 10, 11 – *Lonchodomas rostratus* (Sars).

Fig. 3 – *Niellus* sp.

Fig. 6 – *Telephina* sp.

Fig. 9 – *Megistaspis* sp.





## Trilobites

### Plate 6

Trilobites from the localities in the northwestern Gorny Altai, Bugryshikha Formation, the Gora Altai Section. All specimens are from Member 5, in 3m from the bottom.

*Collection of Aleksander V. Timokhin.*

Figs 1, 3 – *Lonchodomas communis* Levitsky.

Figs 2, 9, 14 – *Lonchodomas rostratus* (Sars).


Fig. 4 – *Eorobergia insignis* Petrunina.

Fig. 5 – *Telephina* ex gr. *mobergi* Hadding.

Fig. 6 – Gen. nov. sp. indet.

Figs 7, 8, 12, 13 – *Homotelus inferus* Levitsky.

Figs 10, 11 – *Megistaspis* sp.





## Trilobites

### Plate 7

Trilobites from the localities in the northwestern Gorny Altai, Bugryshikha Formation, the Malaya Uskuchevka Section. All specimens are from Member 1.

*Collection of Aleksander V. Timokhin.*


Figs 1, 2, 11 – *Niellus* sp.

Figs 3, 5b, 7 – *Eorobergia* sp.

Figs 4, 10 – *Homotelus inferus* Levitsky.

Figs 5a, 6, 8 – *Telephina* sp.

Fig. 9 – *Lonchodomas rostratus* (Sars)



## Trilobites

### Plate 8

Trilobites from the localities in the northwestern Gorny Altai, Khankhara Formation, the Ebogon Section. All specimens are from Member 20.

*Collection of Aleksander V. Timokhin.*

Fig. 1 – *Bronteopsis transversal* Petrunina.

Figs 2, 6, 9, 10 – *Erratecrinuru brutoni* Owen.

Fig. 3 – *Bronteopsis gregaria* Raymond.


Fig. 4 – *Calyptaulax* sp.

Figs 5, 8 – *Niellus* sp.

Fig. 7 – *Lonchodomas rostratus* (Sars).

Figs 11, 12, 13, 14, 15, 16 – *Eorobergia* sp.





## Trilobites

### Plate 9


Trilobites from the localities in the northwestern Gorny Altai, Khankhara Formation, the Ebogon Section. All specimens are from Member 20.

*Collection of Aleksander V. Timokhin.*

Figs 1a, 1b, 2, 3, 4, 5, 6, 7, 8 – *Lonchodomas rostratus* (Sars).

Figs 9, 10, 11, 12, 14 – *Stygina minor* Skjeseth.

Figs 13, 15, 16, 17 – *Megistaspis* sp.





## Trilobites

### Plate 10

Trilobites from the localities in the northeastern Gorny Altai. Figs 1-6: Lebed' River, the Lebed' Section, Karasa Formation, Member 25; figs 8-12: Biya River, Samysh Body, loc. S-1618.

*Collection of Aleksander V. Timokhin.*

Figs 1, 3, 7, 12 – *Homotelus* sp.

Figs 2, 10 – *Lonchodomas rostratus* (Sars).


Fig. 4 – *Calyptaulax* sp.

Figs 5, 6 – *Iliaenus* sp.

Fig. 8 – *Eorobergia* sp.

Fig. 9 – *Megistaspis* sp.

Fig. 11 – *Stygina minor* Skjeseth



## Trilobites

### Plate 11

Trilobites from the localities in the northeastern Gorny Altai, Karasa Formation, the Yurok Section.  
*Collection of Aleksander V. Timokhin.*

Figs 1, 2 – *Cybelurus altaicus* Levitsky:  
Member 1, in 3 m from the bottom, loc. C-1118-1.

Figs 3, 4, – *Paracybeloides(?)* sp.:  
specimen 3 - Member 2, in 8 m from the bottom, loc. C-1118-2; specimen 4 – Member 3, loc. C-1116-c.

Fig. 5 – *Carolinites* sp.:  
Member 3, loc. C-1116-c.


Figs 6, 7 – *Robergia sparsa* Nikolaisen:  
specimen 6 – Member 3, loc. C-1116-c; specimen 7 - Member 2, in 5 m from the bottom, loc. C-1118-2,.

Figs 8, 9, 10, 11, 12 – *Lonchodomas rostratus* (Sars):  
specimen 8 - Member 3, loc. C-1116-c; specimens 9, 12 - Member 2, in 8 m from the bottom, loc. C-1118-2;  
specimen 10 - Member 3, loc. C-1116-a; specimen 11 – Member 3 , loc. C-1116-c.

Figs 13 – 16, *Raymondaspis* sp.  
specimen 14 - Member 2, loc. C-1118-2, 5 m from the bottom; specimen 15 – Member 4, in 5 m from the bottom, loc. C-1118-4; specimen 16 – Member 3 , loc. C-1116-c.

Fig. 17 – *Megistaspis (M.) polyphemus* Brogger:  
Member 1, in 3 m from the bottom, loc. C-1118-1.





## Trilobites

### Plate 12

Trilobites from the localities in the northeastern Gorny Altai. Figs 1-6, 8-16: upper part of the Karasa Formation, upper part of the Tuloi Section, loc. R-409b; fig 7: lower part of the Karasa Formation, the Yurok Section, Member 2, in 7,5 m from the bottom, loc. C-1118-2.

*Collection of Aleksander V. Timokhin.*

Figs 1, 2 – *Megistaspis (M.) polyphemus* Brogger.

Fig. 3 – *Paracybeloides* (?) sp.

Figs 4, 5 – *Robergia sparsa* Nikolaisen.

Fig. 6 – *Carolinites* sp.

Fig. 7 – *Niobe* sp.


Figs 8, 9, 10 – *Lonchodomas rostratus* (Sars).

Fig. 11 – *Agerina* sp.

Fig. 12 – *Raymondaspis* sp.

Fig. 13 – *Niellus* sp.

Figs 14 –16 – *Ampyx* sp.





## Trilobites

### Plate 13

Trilobites from the localities in the eastern Gorny Altai, Tozodov Formation, the Tozodov Section, members 7–8.

*Collection of Aleksander V. Timokhin.*

Figs 1, 4 – *Asaphus knyrkoi* F. Schmidt.


Figs 2, 3, 6, 17 – *Asaphus striatus* Brogger.

Figs 7, 8, 9, 10 – *Lonchodomas rostratus* (Sars).

Fig. 11 – *Pliomera fischeri* (Eichwald).

Figs 5, 14, 16, 18 – *Iliaenus* sp.

Figs 12, 13 – *Homotelus* sp.



## Ostracods

### Plate 14

Ostracods from the localities of the Teletskoe Lakeside area, northeastern part of Gorny Altai.  
*Collections of Taras V. Gonta.*

Figs 1–3. *Primitia* sp.: Samysh River, loc. S-1630, specimen 1 – left valve, lateral view; specimen 2 – left valve, lateral view; specimen 3 – right valve, lateral view.

Figs 4, 5. *Laccochilina* (*Laccochilina*) aff. *torosa* Kanygin: Tarlyk River, loc. S-1631, specimen 4 – male right valve, lateral view; specimen 5 – male left valve, lateral view.

Figs 6, 7. *Bolbina* sp.: Verkhniy Turochak River, loc. S-1754, specimen 6 – male left valve, lateral view, specimen 7 – female left valve, lateral view.

Figs 8, 9. *Rigidella* sp. 1: Tarlyk River, loc. S-1631, specimen 8 – left valve, lateral view; specimen 9 – right valve, lateral view.

Figs 10, 11. *Rigidella* sp. 2: Tarlyk River, loc. S-1631, specimens 10, 11 – left valves, lateral view.

Figs 12–14. *Steuslofia* sp.: Tarlyk River, loc. S-1631, specimens 12, 13, 14 – right valves, lateral view.


Figs 15, 16. *Egorovella* sp.: specimen 15 – Tozodov Brook, loc. S-1442, right valve, lateral view; specimen 16 – Tarlyk River, loc. S-1631, right valve, lateral view.

Figs 17, 18. *Pseudozygobolbina invisitata* Melnikova: Samysh River, loc. S-1630, specimens 17, 18 – left valves, lateral view.

Figs 19, 20. *Hallatina* aff. *orlovi* V. Ivanova: Tarlyk River, loc. S-1631, specimens 19, 20 – left valves, lateral view.

### Plate 14





## Graptolites

### Plate 15

Graptolites from the localities in the northeastern part of Gorny Altai. Scale bar 1 mm.

Collections of Elena V. Lykova, Nikolay V. Sennikov.

Fig. 1. *Eotetragraptus quadribrachiatus* (Hall):

The left bank of the Lebed' River, Pridorozhny Section, loc. B-1011a. Tuloi Formation, Dapingian, *E. hirundo* Zone, *I. caduceus-imitatus* Subzone.

Fig. 2. *Eotetragraptus harti* (T.S. Hall):

The right bank of the Tuloi River, Tuloi Section, loc. S-0724. Tuloi Formation, Dapingian, *E. hirundo* Zone.

Figs 3, 10. *Pseudoisograptus manubriatus* (T.S.Hall):

3 – the right bank of the Tuloi River, Tuloi Section, loc. B-1111. Tuloi Formation, Dapingian, *E. hirundo* Zone;

10 – right bank of the Tuloi River, Tuloi Section, loc. B-117. Tuloi Formation, Dapingian, *E. hirundo* Zone.

Figs 4, 5, 11, 12. *Corymbograptus deflexus* (Elles et Wood):

4, 12 – the right bank of the Tuloi River, Tuloi Section, loc. S-0724. Tuloi Formation, Dapingian, *E. hirundo*

Zone; 5, 11 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-107. Tuloi Formation, Dapingian,

*I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 6, 7. *Corymbograptus inflexus* (Chen et Xia):

6, 7 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-106. Tuloi Formation, Dapingian,

*I. gibberulus* Zone, *C. deflexus* subzone.

Figs 8, 9. *Expansograptus extensus* (Hall):

8 – the right bank of the Tuloi River, Tuloi Section, loc. B-118. Tuloi Formation, Dapingian, *E. hirundo* Zone; 9

– the left bank of the Lebed' River, Pridorozhny Section, loc. B-096. Tuloi Formation, Dapingian, *I. gibberulus*

Zone, *I. maximo-divergens* Subzone.

Figs 13, 14. *Pseudophyllograptus angustifolius angustifolius* Hall:

13, 14 – the right bank of the Biya River, Yurok Section, loc. S-1118-4/18. Tuloi Formation, Floian,

*Ps. angustifolius elongatus* Zone.







## Graptolites

### Plate 16

Graptolites from the localities in the northeastern part of Gorny Altai. Scale bar 1 mm.

Collections of Elena V. Lykova, Nikolay V. Sennikov.

Figs 1, 2, 6. *Pseudoisograptus manubriatus* (T.S.Hall):

1 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-096. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 2 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-109. Tuloi Formation, Dapingian, *E. hirundo* Zone, *I. caduceus-imitatus* Subzone; 6 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-106 a. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *C. deflexus* Subzone.

Figs 3, 4, 10. *Pseudoisograptus manubriatus janus* Cooper et Ni :

3 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-093. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 4, 10 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-096. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 7, 8, 9. *Isograptus gibberulus* (Nicholson):


7, 8 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-108. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 9 – the right bank of the Tuloi River, Tuloi Section, loc. B-117. Tuloi Formation, Dapingian, *E. hirundo* Zone.

Figs 5, 11, 12, 14, 15. *Pseudoisograptus manubriatus* (T.S.Hall):

5, 11, 12, 15 – the right bank of the Tuloi River, Tuloi Section, loc. S-0724. Tuloi Formation, Dapingian, *E. hirundo* Zone; 14 – the right bank of the Tuloi River, Tuloi Section, loc. B-117. Tuloi Formation, Dapingian, *E. hirundo* Zone.

Figs 13. *Isograptus divergens* (Harris)

13 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-107a. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.



## Graptolites

### Plate 17

Graptolites from the localities in the northeastern part of Gorny Altai. Scale bar 1 mm.

Collections of Elena V. Lykova, Nikolay V. Sennikov.

Figs 1, 2, 3, 11. *Isograptus gibberulus* (Nicholson):

1 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-096. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 2, 3 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -107a. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 11 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -108. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 4, 6-9, 13. *Isograptus divergens* (Harris):

4 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-109. Tuloi Formation, Dapingian, *E. hirundo* Zone, *I. caduceus-imitatus* Subzone; 6, 7, 9, 13 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -107a. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 8 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -093. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.


Figs 5. *Isograptus forcipiformis* (Ruedemann):

5 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-093. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 10, 12, 14. *Isograptus maximo-divergens* (Harris):

10, 14 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-109. Tuloi Formation, Dapingian, *E. hirundo* Zone, *I. caduceus-imitatus* Subzone; 12 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -093. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.





## Graptolites

### Plate 18

Graptolites from the localities in the northeastern part of Gorny Altai. Scale bar 1 mm.

Collections of Elena V. Lykova, Nikolay V. Sennikov.

Figs 1, 3. *Isograptus divergens* (Harris):

1 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-109. Tuloi Formation, Dapingian, *E. hirundo* Zone, *I. caduceus-imitatus* Subzone; 3 – the left bank of the Lebed' River, Pridorozhny Section, loc. B –107a. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 2, 4. *Isograptus walcottorum* Ruedemann:

2 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-096. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone; 4 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -093. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 5, 13. *Isograptus paraboloides* Tzaj :

5, 13 – the right bank of the Tuloi River, Tuloi Section, loc. B-118. Tuloi Formation, Dapingian, *E. hirundo* Zone.

Figs 6, 12. *Isograptus elegans* Tzaj:

6 – the right bank of the Tuloi River, Tuloi Section, loc. S-0724. Tuloi Formation, Dapingian, *E. hirundo* Zone; 12 – the left bank of the Lebed' River, Pridorozhny Section, loc. B -096. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Fig. 7. *Isograptus maximo-divergens* (Harris):

7 – the left bank of the Lebed' River, Pridorozhny Section, loc. B-108. Tuloi Formation, Dapingian, *I. gibberulus* Zone, *I. maximo-divergens* Subzone.

Figs 8, 9, 11. *Phyllograptus ilicifolius* Hall:

8, 9, 11 – the right bank of the Biya River, Yurok Section, loc. S-1118-4/5. Tuloi Formation, Floian, *Ps. angustifolius elongatus* Zone.

Fig. 10. *Phyllograptus anna longus* Ruedemann:

10 – the right bank of the Biya River, Yurok Section, loc. S-1118-4/5. Tuloi Formation, Floian, *Ps. angustifolius elongatus* Zone.







## Graptolites

### Plate 19

Graptolites from the localities in the western (Krasnoshchekovo and Maralikha villages, Charysh River) and northeastern (Tuloi, Lebed' and Biya rivers) parts of Gorny Altai. Scale bar 1 mm.

*Collections of Elena V. Lykova, Nikolay V. Sennikov.*

Figs 1, 2, 3, 4, 13. *Eoglyptograptus dentatus* (Brongniart):

1, 2, 3 – West of Krasnoshekovo Village, Batun Section, loc. S-813a. Voskresenka Formation, Darriwilian, *E. dentatus* Zone.

Figs 4, 13. *Amplexograptus confertus* (Lapworth):

4, 13 – the right bank of the Tuloi River, Tuloi Section, loc. LSS-409. Karasa Formation, Darriwilian, *E. dentatus* Zone.

Figs 5, 6, 14–16. *Hustedograptus teretiusculus* (Hisinger):

5, 6 – Former Stretinka Village, Lebed' River, Lebed' Section, loc. S-7546 (=R-254). Karasa Formation, Darriwilian, *H. teretiusculus* Zone; 14 - the right bank of the Biya River, Yurok Section, loc. S-1118-1/3. Tuloi Formation, Floian, *Ps. angustifolius elongatus* Zone; 15 - the right bank of the Biya River, Yurok Section, loc. S-1118-2/8. Tuloi Formation, Floian, *Ps. angustifolius elongatus* Zone; 16 – left bank of the Charysh River, Charysh Section, loc. S-804-6. Bugryshikha Formation, Darriwilian - Sandbian, *H. teretiusculus* – *C. peltifer*, *A. antiquus lineatus* zones.

Figs 7-11. *Undulograptus austrodentatus* (Harris et Keble):

7–10 – vicinity of Maralikha Village, Maralikha Section, loc. S-072. Voskresenka Formation, Darriwilian, *U. austrodentatus* Zone; 11 – vicinity of Maralikha Village, Maralikha Section, loc. S-0523. Voskresenka Formation, Darriwilian, *U. austrodentatus* Zone.

Fig. 12. *Undulograptus sinicus* (Mu et Lee):

12 – vicinity of Maralikha Village, Maralikha section, loc. S-072. Voskresenka Formation, Darriwilian, *U. austrodentatus* Zone.



## Conodonts

### Plate 20

Late Cambrian - Early Ordovician conodonts from the localities in the northern part of Gorny Altai, Kamlak Village, Kamlak Formation.

*Collection of Tatyana Yu. Tolmacheva.*

Figs 1, 2. *Iapetonudus* sp.:

The Kamlak Section, Member 10, Middle Kamlak Subformation, Lower Tremadocian.

Fig. 3. *Cordylodus lindstromi* Druce et Jones:

The Kamlak Section, Member 10, Middle Kamlak Subformation, Lower Tremadocian.

Figs 4, 5. *Cordylodus caseyi* Druce et Jones, emend. Landing:

The Kamlak Section, Members 9-10, Middle Kamlak Subformation, Upper Cambrian.

Figs. 6, 7. *Iapetognathus* sp.:

The Kamlak Section, Member 10, Middle Kamlak Subformation, Lower Tremadocian.

Fig. 8. *Hispidodontus* cf. *H. kulumbense* Tolmacheva et Abaimova:

The Kamlak Section, Members 2, Middle Kamlak Subformation, Upper Cambrian.

Figs 9, 13, 14. *Cordylodus proavus* Müller:

The Kamlak Section, Members 11, Upper Kamlak Subformation, Lower Ordovician.

Fig. 10. *Hirsutodontus simplex* (Druce et Jones, 1971):

The Kamlak Section, Members 2, Middle Kamlak Subformation, Upper Cambrian.

Fig. 11. *Hirsutodontus* sp.:

The Kamlak Section, Members 2, Middle Kamlak Subformation, Upper Cambrian.

Fig. 12. *Phakelodus tenuis* (Müller):

The Kamlak Section, Members 9, Lower Kamlak Subformation, Upper Cambrian.

Figs 15, 16. *Westergaardodina* sp.:

The Kamlak Section, Members 5, Lower Kamlak Subformation, Upper Cambrian.

Figs 17, 19. *Eoconodontus notchpeakensis* (Miller):

The Kamlak Section, Members 9-10, Middle Kamlak Subformation, Upper Cambrian.

Fig. 18. *Furnishina furnishi* Müller:

The Kamlak Section, Members 5, Lower Kamlak Subformation, Upper Cambrian.


Figs 20, 21. *Semiacontiodus* sp.:

The Kamlak Section, Members 9-10, Middle Kamlak Subformation, Upper Cambrian.

Figs 22, 23. *Hispidodontus* cf. *triangularis* (Abaimova, 1975):

The Kamlak Section, Members 9-10, Middle Kamlak Subformation, Upper Cambrian.





## Conodonts

### Plate 21

Late Ordovician conodonts from the localities in the northeastern part of Gorny Altai, Lebed', Biya and Bura rivers, Gur'yanovka Formation.

*Collections of Tatyana Yu. Tolmacheva and Olga T. Obut*

Figs 1–4, 9. *Phragmodus undatus* Branson et Mehl:

1–3, 9 – the Lebed' Section, Member 13, Gur'yanovka Formation, Sandbian; 4 – the Biya Section, Member 20, Gur'yanovka Formation, Sandbian.

Figs 5–8. *Belodina compressa* (Branson et Mehl):

5, 7, 8 – the Lebed' Section, Member 13, Gur'yanovka Formation, Sandbian; 6 – the Biya Section, Member 3, Gur'yanovka Formation, Sandbian.

Figs 10–16. *Scandodus* sp.:

10–14 – the Lebed' Section, Member 13, Gur'yanovka Formation, Sandbian, 15–16 – the Bura Section, Member 10, Gur'yanovka Formation, Sandbian.

Figs 17–19. *Drepanodus* sp.:

the Bura Section, Member 10, Gur'yanovka Formation, Sandbian.

Fig. 20. *Erraticodon* sp.:

the Biya Section, Member 20, Gur'yanovka Formation, Sandbian.

Fig. 21. *Aphelognathus* sp.:

the Lebed' Section, Member 13, Gur'yanovka Formation, Sandbian.

Fig. 22. *Drepanoistodus suberectus* (Branson et Mehl):

the Lebed' Section, Member 13, Gur'yanovka Formation, Sandbian.

Fig. 23. *Panderodus* cf. *P. gracilis* (Branson et Mehl):

the Lebed' Section, Member 13, Gur'yanovka Formation, Sandbian.





## Conodonts

### Plate 22

Late Ordovician conodonts from the locality in the northwestern part of Gorny Altai, Siliceous-terrigenous Body, Katian, the Suetka Section, Member 1. Scale = 100  $\mu\text{m}$ .

*Collections of Tatyana Yu. Tolmacheva and Olga T. Obut.*

Figs 1–3, 6. *Periodon grandis* (Ethington):

1 – Sc element, 2 – Sc element, 3 – Sd element, 4 – M element.

Figs 4, 5. *Histiodella* sp.:

8 – P element, 9 – S element.

Figs 7, 8. *Panderodus* sp.

Figs 9, 13, 14. *Paroistodus? mutatus* (Branson et Mehl):


7 – M element, 11, 12 – S elements.

Fig. 10. *Decoriconus* sp.

Figs 11, 12. *Protopanderodus insculptus* (Branson et Mehl).

Fig. 15. *Belodina compressa* (Branson et Mehl).

Fig. 16. *Belodina* sp.



## Conodonts

### Plate 23

Middle Ordovician conodonts from the Teletskoe Lakeside area, eastern Gorny Altai, Samysh Body, Darriwilian. Specimens illustrated on figs 1, 2, 4–10, 12, 15, 16, 18 - from locality on the right bank of Samysh River. Specimens illustrated on figs 3, 11, 13, 14, 17, 19, 20 – from locality on Nizhniy Turochak Brook, right tributary of the Iogach River. Scale = 200  $\mu\text{m}$ .

*Collections of Olga T. Obut.*

Figs 1–4. *Acodus* sp.

1, 2 – Sd elements, 3, 4 – M elements.

Figs 5, 9, 10. *Drepanoistodus* cf. *arcuatus* Pander:

5, 9 – S elements, 10 – P element.

Figs 6–8, 13 – *Periodon* sp.:

6, 7 – S elements, 8, 13 – M elements.

Fig. 11. ? *Drepanodus* sp.

Figs 12, 18–20. *Drepanoistodus* sp.:


12 – M element, 18–20 – S elements.

Fig. 14. *Eoplacognathus* sp.:

Pa element.

Figs. 15–17. *Parapanderodus striatus* (Graves et Ellison).





*Научное издание*

Николай Валерианович Сенников, Ольга Тимофеевна Обут, Елена Викторовна Лыкова, Александр Владиленович Тимохин, Тарас Викторович Гонга, Ралия Ахатовна Хабибулина, Татьяна Александровна Щербаненко, Татьяна Павловна Киприянова

**ОРДОВИКСКИЕ СЕДИМЕНТАЦИОННЫЕ БАССЕЙНЫ  
И ПАЛЕОБИОТЫ ГОРНОГО АЛТАЯ**

На английском языке

Утверждено к печати Ученым советом  
Института нефтегазовой геологии и геофизики им. А.А. Трофимука СО РАН

Оригинал-макет *Владимировой А.В.*

---

Подписано в печать 21.06.2019. Формат 60×84/8. Усл. п. л. 21,39. Тираж 120 экз. Закз № 127

---

Издательство СО РАН  
630090, Новосибирск, Морской просп., 2  
E-mail: [psb@sibran.ru](mailto:psb@sibran.ru)  
Тел. (383) 330-80-50  
Отпечатано в Издательстве СО РАН  
Интернет-магазин Издательства СО РАН  
<http://www.sibran.ru>